isotopic measurement and could therefore not be used for regression analysis.

Figure 1a shows one example of the difference between the original estimate of Evaristo *et al.*¹ (green triangle) and our revised estimate (orange triangle) of the plant water source (site ID 26). For most of the sites, our estimates differ considerably from those of Evaristo *et al.*¹ (see Extended Data Figs 1, 2 and 3).

Extended Data Figs 1, 2 and 3 also reveal that the data set is extremely heterogeneous in terms of the number of sampled points for plant and groundwater, with sometimes inconsistent data leading to unrealistic values of $\delta^{18}\text{O}_{\text{intercept}}$ and $\delta^2\text{H}_{\text{intercept}}$ for certain sites. Therefore, we applied two non-exclusive criteria to assess the consistency of the calculated intersection values: criterion 1, $\delta^2\text{H}_{\text{intercept}} < \max(\delta^2\text{H}_{\text{plant}})$; and criterion 2, $\delta^2\text{H}_{\text{intercept}} > -200\text{‰}$. The first criterion implies that $m < a$, while the second criterion evaluates whether the plant water source hydrogen isotopic composition value is realistic. Eleven sites failed at least one of the two criteria (IDs 3, 7, 8, 17, 20, 21, 23, 24, 32, 36 and 44). Figure 1b shows one example of an inconsistent data set (site ID 3).

Results of this analysis (summarized in Table 1) show that at 26 sites, where data were consistent, $\delta^2\text{H}_{\text{GW}}$ was statistically different from $\delta^2\text{H}_{\text{intercept}}$ using the non-parametric Wilcoxon rank sum test ($\alpha = 0.05$). In conclusion, rainfall segregation (as defined by Evaristo *et al.*¹) could be observed for only 57% of the sites of the authors' data set and at 74% of the sites with consistent estimates of the intercept as defined by our two criteria.

Online Content Methods, along with any additional Extended Data display items and Source Data, are available in the online version of the paper; references unique to these sections appear only in the online paper.

Mathieu Javaux^{1,2,3}, Youri Rothfuss¹, Jan Vanderborght¹, Harry Vereecken¹ & Nicolas Brüggemann¹

¹Institute of Bio- and Geosciences, IBG-3 Agrosphere, Forschungszentrum Jülich GmbH, Jülich 52425, Germany. email: m.javaux@fz-juelich.de

²Earth and Life Institute, Université catholique de Louvain (UCL), Louvain-la-Neuve 1348, Belgium.

³Department of Land, Air and Water Resources, University of California Davis, Davis, California 95616, USA.

Received 25 January; accepted 6 June 2016.

1. Evaristo, J., Jasechko, S. & McDonnell, J. Global separation of plant transpiration from groundwater and streamflow. *Nature* **525**, 91–94 (2015).

Author Contributions M.J. found the mistake in the original paper. Y.R. performed the statistical analyses. M.J., Y.R., J.V., H.V. and N.B. discussed the results and wrote the manuscript.

Competing Financial Interests Declared none.

doi:10.1038/nature18946

Evaristo *et al.* reply

REPLYING TO M. Javaux, Y. Rothfuss, J. Vanderborght, H. Vereecken & N. Brüggemann *Nature* **536**, <http://dx.doi.org/10.1038/18946> (2016)

In the accompanying Comment¹, Javaux *et al.* correct a mistake in equation (2) of our work²; as they point out¹, the mistake does not impact the central conclusion of our paper that ecohydrological separation is widespread². However, equations (2) and (3) in ref. 2 calculate the source precipitation value of xylem water as the point where the xylem water evaporation line (EL) intersects the local meteoric water line (LMWL). In so doing, Javaux *et al.*¹ note that the mistake affects our finding² that “at 80% of the sites, the precipitation that supplies groundwater recharge and streamflow is different from the water that supplies parts of soil water recharge and plant transpiration”.

There are two key points in our response.

(1) We recognize the mistake now noted in equation (2) and thank Javaux *et al.*¹ for this correction. These authors¹ find that rainfall segregation could be observed at only 74% of the sites (as defined by the two criteria in ref. 1), and not 80% as we originally reported². (2) Our work² presented evidence for ecohydrological separation based on a meta-analysis of isotopic dual liquid water isotope data ($\delta^2\text{H}$ and $\delta^{18}\text{O}$) from 47 studies. This conclusion is supported by studies that analysed water vapour isotope data from the Tropospheric Emissions Spectrometer aboard NASA's Aura satellite³ and by global differences between annual precipitation and groundwater isotope compositions^{4,5}. These global-in-scale lines of evidence support earlier field evidence^{6,7} that ecohydrological separation (defined as plants using water of a character different to that of mobile water found in soils, groundwater and streamflow) is widespread, and is the rule rather than the exception. Ecohydrological separation was calculated using equation (1) in ref. 2.

It must be understood that equation (1) in ref. 2 is independent of the source precipitation analysis, which was calculated using equations (2) and (3) in that paper. Therefore, any issue with equation (2) in our paper, like the one raised by ref. 1, does not affect the ecohydrological separation conclusion.

We hope that this exchange will generate further interest in the use of stable O and H isotopes in plant water relation studies.

J. Evaristo¹, S. Jasechko² & J. J. McDonnell^{1,3,4}

¹Global Institute for Water Security and School of Environment and Sustainability, University of Saskatchewan, Saskatoon, Saskatchewan S7N 3H5, Canada.

email: jaivime.evaristo@usask.ca

²Department of Geography, University of Calgary, Calgary, Alberta T2N 1N4, Canada.

³School of Geosciences, University of Aberdeen, Aberdeen AB34 3FX, UK.

⁴Department of Forest Engineering, Resources and Management, Oregon State University, Corvallis, Oregon 97331, USA.

1. Javaux, M., Rothfuss, Y., Vanderborght, J., Vereecken, H. & Brüggemann, N. Isotopic composition of plant water sources. *Nature* **536**, <http://dx.doi.org/10.1038/nature18946> (2016).
2. Evaristo, J., Jasechko, S. & McDonnell, J. J. Global separation of plant transpiration from groundwater and streamflow. *Nature* **525**, 91–94 (2015).
3. Good, S. P., Noone, D. & Bowen, G. Hydrologic connectivity constrains partitioning of global terrestrial water fluxes. *Science* **349**, 175–177 (2015).
4. Jasechko, S. & Taylor, R. G. Intensive rainfall recharges tropical groundwaters. *Environ. Res. Lett.* **10**, 124015 (2015).
5. Jasechko, S. *et al.* The pronounced seasonality of global groundwater recharge. *Wat. Resour. Res.* **50**, 8845–8867 (2014).
6. Brooks, J. R., Barnard, H. R., Coulombe, R. & McDonnell, J. J. Ecohydrologic separation of water between trees and streams in a Mediterranean climate. *Nat. Geosci.* **3**, 100–104 (2010).
7. Evaristo, J., McDonnell, J. J., Scholl, M. A., Bruijnzeel, L. A. & Chun, K. P. Insights into plant water uptake from xylem-water isotope measurements in two tropical catchments with contrasting moisture conditions. *Hydrol. Process.* <http://dx.doi.org/10.1002/hyp.10841> (in the press).

doi:10.1038/nature18947