

**References**


1. Ozarto, K., Mayer, N., and Sachs, D. H. 1980. Hybridoma Cell Lines Secreting Monoclonal Antibodies to Mouse H-2 and Ia Antigens. *J. Immunology*. **124**:533-540.
2. Long, W. J., Palombo, A., Schofield, T. L., and Eminie, E. A. 1988. Effects of Culture Media on Murine Hybridomas: Definition of Optimal Conditions for Hybridoma Viability, Cellular Proliferation, and Antibody Production. *Hybridoma*. **7**:69-77.
3. Iscove, N. N., and Melchers, F. 1978. Complete Replacement of Serum by Albumin, Transferrin, and Soybean Lipid in Cultures of Lipopolysaccharide-Reactive B Lymphocytes. *J. Exp. Med.* **147**:923-933.
4. Cleveland, W. L., Wood, I., and Erlanger, B. F. 1983. Routine Large-Scale Production of Monoclonal Antibodies in a Protein-Free Culture Medium. *J. Imm. Meth.* **56**:221-234.
5. Antoniadis, H. N., Scher, C. D., and Stiles, C. D. 1979. Purification of Human Platelet-Derived Growth Factor. *Proc. Natl. Acad. Sci. USA* **76**:1809-1813.
6. Helden, C.-H., Westermarck, B., and Wasteson, A. 1979. Platelet-Derived Growth Factor: Purification and Partial Characterization. *Proc. Natl. Acad. Sci. USA* **76**:3722-3726.
7. Maciage, T., Kelly, B., Cerundolo, J., Ilesley, S., Kelley, P. R., Gaudreau, J., and Forand, R. 1980. Hormonal Requirements of Baby Hamster Kidney Cells in Culture. *Cell. Biol. Int. Rep.* **4**:43-50.
8. McKeehan, W.L., and Ham, R. G. 1977. Methods for Reducing the Serum Requirement for Growth *in vitro* of Nontransformed Diploid Fibroblasts. *Dev. Biol. Stand.* **37**:97-108.
9. Hutchings, S.F., and Sato, G.H. 1978. Growth and Maintenance of Hela Cells in Serum-Free Medium Supplemented with Hormones. *Proc. Natl. Acad. Sci. USA* **79**:4074-4077.
10. Wolfe, R.A., Braatz, J.A., Miller, D. A., and Heifetz, A.H. 1988. A New Serum-Free Medium for Monoclonal Antibody Production. *Biotechniques*. **6**:62-67.
11. Glassy, M.C., Peters, R. E., and Mikhailov, A. 1987. Growth of Human-Human Hybridomas in Serum-Free Media Enhances Antibody Secretion. *In Vitro Cell. & Dev. Biol.* **23**:745-749.
12. Higuchi, K. 1977. Cultivation of Mammalian Cell Lines in Serum-Free Chemically Defined Medium. *Cell Biol.* **14**:131-136.
13. Barnes, D., and Sato, G. 1980. Serum-Free Cell Culture: A Unifying Approach. *Cell*. **22**:649-655.
14. Stampfer, M. R., Hackett, A. J., Smith, H. S., Hancock, M. C., Leung, J. P., and Edginton, T. S. 1982. Cold Spring Harbor Conf. Cell Proliferation. **9**:819-829.
15. Hammond, S. L., Ham, R. G., and Stampfer, M. R. 1984. Serum-Free Growth of Human Mammary Epithelial Cells: Rapid Clonal Growth in Defined Medium and Extended Serial Passage with Pituitary Extract. *Proc. Natl. Acad. Sci. USA* **81**:5435-5439.

**CULTURE MEDIA**



Clonetics Corp.'s (San Diego, CA) **serum-free keratinocyte growth medium (KGM™)** is included in the company's EpiPack® kit for culturing normal human epidermal keratinocytes. KGM promotes the growth of a variety of other epithelial cell types, as well, and eliminates the cytotoxicity and enzyme inactivation that can occur with serum-based media. In addition to KGM, the kit contains cultured normal human epidermal keratinocytes (NHEK), which can be used as an *in vitro* model for research in toxicology, cancer, dermatology, immunology, and gene expression. Established correlations between NHEK responses to irritants and those of live animals in the Draize Eye and Skin Irritancy tests offer the possibility of substituting NHEK studies for *in vivo* experiments.

Write in 802 on Reader Service Card



The clear, flat well bottoms of Cel-Cult **multiwell plates** from Sterilin (Middlesex, U.K.) make this product useful not only for tissue culture assay systems but also for other techniques in which small samples must be kept in closed containers for microscopic examination. The plates fit standard microscope multiwell transport stages and are available sterile or non-sterile, with or without lids, and in 24- and 96-well versions.

Write in 803 on Reader Service Card



Origen™ **hybridoma cloning factor (HCF)** spurs the growth of murine hybridoma cell lines. Its manufacturer, Igen (Rockville, MD), reports that Origen HCF is more efficient and easier to use than feeder layers and conditioned media, and that it allows hybridomas to be cloned at lower cell densities. Partially purified and filter-sterilized, Origen HCF is derived from continuous macrophage cell lines. It remains stable for at least six months at -20°C, and is supplied in 50 ml bottles at 10X concentration.

Write in 813 on Reader Service Card



**Tissue culture flasks** from J. Bibby Science Products (Stone, Staffs, U.K.) rely on a canted neck design to maximize cell recovery in applications such as vaccine production. Based on a unique Corning design, Bibby flasks are strong and leak-resistant, and have a shallow profile for economical use of space when large number of flasks must be stored.

Write in 814 on Reader Service Card



Pre-aliquoted transformation media add to the convenience of Ultra-comp™, an *E. coli* **transformation kit** from Invitrogen (San Diego, CA). The kit includes ultra-rich S.O.C. medium, competent *E. coli*, and pre-aliquoted test plasmid sufficient for 10 transformations. Invitrogen offers nine different strains of the organism, selected for insert stability and adaptability to special cloning and analysis strategies. Ultracomp kits are packaged with five aliquots of either 0.2 ml or 1 ml.

Write in 810 on Reader Service Card