Activated Affinity Columns

Unique protein separations require unique solutions. That's why ChromatoChem developed a series of aldehyde activated columns that allow you to immobilize your own protein or peptide ligands on silica. With just one pass of the affinity ligand through the column, you've created your own custom affinity support.

ChromatoChem's new **HIPAC™ Aldehyde Columns**, packed with our high-performance, pH-stable silica, can withstand over 100 cycles—with no evidence of silica erosion. What's more, our unique linker chemistry reduces steric hindrance, to give you:

- ❖ **Faster** Protein Coupling
- ❖ **Rapid** Ligand-Receptor Binding Kinetics
- ❖ **High** Capture Efficiency
- ❖ **Very Low** Nonspecific Binding

So if you have special purification needs, save time and money by creating an affinity column that captures *only* the molecules you're looking for...and nothing else.

Columns are available in analytical and prep-scale sizes. Call us today toll-free for more details.

ChromatoChem, Inc.
2837 Fort Missoula Road
Missoula, MT 59801
Call toll-free: 800/426-7227
Write in No. 341 on Reader Service Card

JAPAN ROUNDUP/

Research scientists in the School of Agriculture at Tokyo University have isolated the structural gene for a bacterial ice-nucleation protein. This protein probably causes frost damage to crops by stimulating the rapid formation of ice crystals at temperatures just below freezing.

Soichi Arai's group isolated the gene for the ice-nucleation protein from a cDNA library cloned from a soil bacterium indigenous to Shizuoka Prefecture. The scientists then expressed this gene in *Escherichia coli*, with the same efficiency as in its natural host.

The protein encoded by this gene contains a 16-amino acid sequence that is repeated 70 times. Molecular modeling of this repeated amino acid segment suggests that it forms a structure with six-fold symmetry that matches the six-fold symmetry of ice crystals. This may be how the protein efficiently stimulates the nucleation of ice crystals.

The research groups of Masaki Suemura and Hiroshi Kikutani at Osaka University Medical School have used the IgE receptor to block the antigen-mediated release of allergens by cells of the immune system *in vitro*. This represents an important first step towards developing a method for using IgE receptors to treat allergic reactions.

In allergic inflammation, circulating IgE antibodies first bind to allergens, such as pollen. These IgE molecules subsequently bind to specific cell surface receptors on mast cells, macrophages, or acidophils. The cross-linking of IgE receptors in the cell membrane induces the release of substances such as histamine and bradykinin. These agents induce the symptoms of an allergic reaction, including tissue inflammation, airways constriction, and capillary dilation.

Suemura and colleagues were able to prevent the allergen-induced release of these substances from macrophages and acidophilic cells *in vitro* by adding large quantities of a soluble fragment of the IgE receptor to the culture medium. These IgE receptor fragments bound the IgE immunoglobulins in solution, before they could bind to cell surface receptors.

Scientists at Kikkoman (Nodashi) are developing methods to employ the enzyme fucose dehydrogenase to quantify fucose levels in urine. Recent studies suggest that there is a good correlation between creatine-fucose levels in the urine and some types of cancer, including liver and skin cancer.

L-fucose is a six-carbon sugar that occurs on many tissue surfaces. It is found in the complex carbohydrate moieties that determine blood groups, for example, and also plays an important role in the clearing of serum proteins by the liver and in the homing of macrophages.

Kikkoman and its subsidiary Seishin Seiyaku Co. (Tokyo) plan to incorporate the enzyme into a medical diagnostic kit. Takara Shuzo Co. (Kyoto) is developing a similar assay.

Prepared by Hal Plotkin and Ken Coleman, *Biotechnology in Japan Newsservice*, Japan Pacific Associates (Palo Alto, CA).