MXL-3 and HLH-30 transcriptionally link lipolysis and autophagy to nutrient availability

Eyleen J. O'Rourke and Gary Ruvkun

Nat. Cell Biol. **15**, 668–676 (2013); published online 21 April 2013; corrected after print 21 November 2014

The original version of this Article did not mention TFEB, the mammalian orthologue of the protein HLH-30, in the abstract. The fourth sentence in the abstract should have read 'Here we identify MXL-3 and HLH-30 (TFEB orthologue) as transcriptional switches coupling lysosomal lipolysis and autophagy to nutrient availability and controlling fat storage and ageing in *Caenorhabditis elegans*.' This has been corrected in all online versions of the Article.

MicroRNAs are transported in plasma and delivered to recipient cells by high-density lipoproteins

Kasey C. Vickers, Brian T. Palmisano, Bassem M. Shoucri, Robert D. Shamburek and Alan T. Remaley

Nat. Cell Biol. **13**, 423–433 (2011); published online 20 March 2011; corrected after print 8 December 2014

In the version of this Article originally published online, incorrect values appearing in Supplementary Table S4 were used to generate Fig. 3a and to calculate the *P* and *R* values. The annotation between Exosome and HDL should have read '*R* = 0.22*', the annotation between Exosome and LDL should have read '*R* = 0.63**' and the annotation between HDL and LDL should have read '*R* = 0.54**', where ***P* < 0.0001 and **P* < 0.001. The amended Fig. 3a is shown below. These errors have been corrected in the online versions of the Article, and in the Supplementary Information.