

at the Pacific east coast and the Indian Ocean west coast. Both coasts have warm water currents of similar strengths flowing north to south, yet, at 3,500 km apart and facing different oceans, they have distinct algal floras.

The seaweed distribution shifts were similar in magnitude on both coasts. The authors warn that temperate Australia is home to many seaweed-dependent marine organisms. Global warming is therefore likely to be driving the rearrangement of whole marine communities — not merely altering the seaweed composition of Australia's shores. AP

**CLIMATOLOGY**

**Feedbacks feeding back**

*J. Geophys. Res.* **116**, D20109 (2011)

Self-reinforcing processes, known as positive feedbacks, amplify climate change, and are responsible for rates of change in the Arctic being among the highest on Earth. Although the major processes involved in these feedbacks are quite well understood, it is not clear how different feedbacks interact.

Yonghua Chen of Columbia University and the Goddard Institute for Space Studies, USA, and co-workers used a climate model to investigate how the amount of thermal radiation emitted towards the surface by the Arctic atmosphere varied with changes in water vapour and cloud properties throughout the seasonal cycle, and how they are projected to increase over the twenty-first century.

The results show that the emitted thermal radiation was most sensitive to water vapour and cloud density in the winter, explaining much of why the largest amplification of surface temperature occurs during this season. The increased levels of water vapour and cloud density expected in coming decades owing to greater greenhouse-gas concentrations will weaken the feedback interaction, suggesting that the current rate of amplified warming in the Arctic, relative to the rest of the Northern Hemisphere, may be reduced in future. AB

**METEOROLOGY**

**The drought child**

*Environ. Res. Lett.* **6**, 044007 (2011)

The El Niño/Southern Oscillation (ENSO) is a coupled atmosphere–ocean circulation occurring over the tropical Pacific Ocean, and is one of the main sources of the Earth's climate variability. Its extremes — the El Niño (warm phase) and La Niña

(cold phase) — both affect rainfall in many parts of the world.


To better understand how ENSO influences droughts, Sergio Vicente-Serrano, of the Pyrenean Institute of Ecology (IPE-CSIC) in Zaragoza, Spain, and co-workers looked globally at the role of the ENSO phases on various drought types associated with hydrological, agricultural and environmental impacts.

The results indicate that the number of regions and months affected, and the total surface area with drought is much higher for El Niño phases than for La Niña phases. In large areas of America and Eastern Europe, ENSO contributes to droughts on short timescales of one to three months at the beginning of the events. In South Africa, Australia and Southeast Asia the effects were more obvious some months later, and at longer timescales. The authors suggest that the large temporal lag between the development of ENSO phenomena and the identification of drought conditions detected in the study may help in forecasting dry conditions in some regions up to one year before their occurrence, potentially improving early warning and adaptation to drought conditions. AB

**METEOROLOGY**

**Sooty cyclones**

*Nature* **479**, 94–97 (2011)



© NASA/JPL/QUIKSCAT SCIENCE TEAM

The development of tropical storms in the Arabian Sea — associated with low-pressure cyclonic airflow — is limited by the interaction of winds at different heights, an effect known as vertical wind shear. The region is strongly influenced by monsoonal circulation patterns, which exhibit strong vertical shear and inhibit cyclone development. However, emissions of anthropogenic aerosols (fine particles suspended in the atmosphere) are thought to have weakened this monsoonal circulation. As such, it seemed plausible that the aerosol-driven circulation modification also affects the intensity of Arabian Sea tropical cyclones.

Amato Evan, of the University of Virginia in Charlottesville, USA, and co-workers investigated the intensity of the pre-monsoon tropical cyclones in the Arabian Sea during the period 1979–2010.

Based on a combination of modelling and observational data, they report an increase in pre-monsoon intensity and demonstrate that this is concurrent with a simultaneous upward trend in anthropogenic black carbon and sulphate aerosol emissions. AB

**BIOGEOGRAPHY**

**In hot water**

*Science* **334**, 652–655 (2011)



MEGAN SAUNDERS

To speak of a steady poleward advance of species is to oversimplify how organisms are responding to a warming world, argues a team led by Michael Burrows of the Scottish Marine Institute in Oban. A more accurate approach is to map spatial changes in isotherms, and then to consider the challenges that organisms face in maintaining their thermal niches.

Burrows and colleagues did this, and then compared the results with a calculation of the seasonal timing of temperatures over the past 50 years, measured in days per decade. They report that both methods show a patchy velocity of climate change on land and seasonal shifts at sea. For example, spring advanced 30–40% faster in the oceans in both hemispheres than on land between 1960 and 2009.

The authors say that certain marine zones are particularly at risk from climate change, notably those near the poles where species may lose their thermal niches entirely, and places such as the Mediterranean Sea, where the European land mass prohibits creatures from tracking thermal niches. AP

*Written by Anna Petherick, Monica Contestabile and Alastair Brown.*