ARTICLE

Received 17 Aug 2010 | Accepted 13 Apr 2011 | Published 17 May 2011

DOI: 10.1038/ncomms1311

Transcription factor IRF8 directs a silencing programme for T_H17 cell differentiation

Xinshou Ouyang¹, Ruihua Zhang¹, Jianjun Yang¹, Qingshan Li¹, Lihui Qin², Chen Zhu³, Jianguo Liu⁴, Huan Ning⁴, Min Sun Shin⁵, Monica Gupta⁶, Chen-Feng Qi⁵, John Cijiang He¹, Sergio A. Lira¹, Herbert C. Morse III⁵, Keiko Ozato⁶, Lloyd Mayer¹ & Huabao Xiong¹

T_H17 cells are recognized as a unique subset of T helper cells that have critical roles in the pathogenesis of autoimmunity and tissue inflammation. Although ROR γ t is necessary for the generation of T_H17 cells, the molecular mechanisms underlying the functional diversity of T_H17 cells are not fully understood. Here we show that a member of interferon regulatory factor (IRF) family of transcription factors, IRF8, has a critical role in silencing T_H17-cell differentiation. Mice with a conventional knockout, as well as a T cell-specific deletion, of the *Irf8* gene exhibited more efficient T_H17 cells. Indeed, studies of an experimental model of colitis showed that IRF8 deficiency resulted in more severe inflammation with an enhanced T_H17 phenotype. IRF8 was induced steadily and inhibited T_H17-cell differentiation during T_H17 lineage commitment at least in part through its physical interaction with ROR γ t. These findings define IRF8 as a novel intrinsic transcriptional inhibitor of T_H17-cell differentiation.

¹ Immunology Institute, Department of Medicine, Mount Sinai School of Medicine, 1 Gustave L. Levy Place, New York, New York 10029, USA. ² Department of Pathology, Mount Sinai School of Medicine, New York, New York 10029, USA. ³ Center for Neurologic Diseases, Brigham and Women's Hospital, Harvard Medical School, Boston, Massachusetts 02115, USA. ⁴ Division of Immunobiology, Saint Louis University School of Medicine, St Louis, Missouri 63104, USA. ⁵ Laboratory of Immunopathology, National Institute of Allergy and Infectious Diseases, Bethesda, Maryland 20892, USA. ⁶ Program in Genomics of Differentiation, Eunice Kennedy Shriver National Institute of Child Health and Human Development, National Institutes of Health, Bethesda, Maryland 20892, USA. Correspondence and requests for materials should be addressed to H.X. (email: huabao.xiong@mssm.edu).

CD4⁺ T helper (T_H) T cell subsets are characterized by the secretion of unique cytokine profiles and have critical roles in orchestrating adaptive immune responses. In addition to T_H1 and T_H2 cells, T_H17 cells have been identified more recently as a third T_H subset mediating inflammatory and autoimmune responses through the production of interleukin (IL)-17A, IL-17F and IL-22 (refs 1–4). T_H17 lineage commitment is initially driven by transforming growth factor (TGF)- β in the presence of IL-6 or IL-21 (refs 5–8), whereas IL-23 serves to expand or maintain T_H17 populations^{2,5,9,10}. The orphan nuclear receptor, RORC, also known as ROR γ t, has been identified as the master transcription factor for T_H17 development¹¹. The differentiation of T_H17 cells is also regulated by several recently described positive and negative feedback loops involving IL-21, IL-23R, IL-10 and IL-27 (refs 6, 7, 12–15), indicating that intrinsic genetic programmes may contribute to the silencing of T_H17 lineage commitment. There is increasing evidence that T_H17 cells are involved in the pathogenesis of various autoimmune/inflammatory diseases, including multiple sclerosis, rheumatoid arthritis, inflammatory bowel diseases and asthma¹⁶. Thus, a more complete understating of the molecular mechanisms involved in the regulation of T_H17 immune responses should provide insights into the pathogenesis and treatment of these and possibly other inflammatory diseases. Several transcription factors, including ROR γ t, ROR α , STAT3 and interferon regulatory factor (IRF)4, have been reported to be important for T_H17-cell differentiation. However, the silencing programme for T_H17-cell differentiation has not been fully examined.

IRF8, a member of the IRF family, is expressed by B cells, dendritic cells (DCs), macrophages^{17–19} and activated T cells^{20,21}, and has been shown to have a diverse roles in the regulation of innate and adaptive immune responses. IRF8 has a DNA-binding domain in the amino (N)-terminal half of the protein and an IRF association domain in the carboxy (C) terminus that is responsible for heterodimerization with other transcription factors²². IRF8 functions as a transcriptional repressor or activator depending on the formation of different heterodimeric DNA-binding complexes with partners that include members of the ETS family and the IRF family²². It is known that IRF8 has critical roles in the differentiation of myeloid cells, promoting monocyte over granulocyte differentiation²³. It is also a crucial regulator of many aspects of DC development, differentiation and function²⁴, thereby having an essential role in the establishment of innate immune responses. Although IRF8 is critical for the regulation of immune cell growth, differentiation and survival²⁵, the direct effects of IRF8 on T-cell activation and differentiation are incompletely understood.

In the present study, we show that mice deficient in IRF8 because of a conventional knockout (KO) or with a T cell-specific conditional deletion exhibited enhanced T_H17-cell differentiation while exhibiting no significant effects on T_H1 or T_H2 cells. In addition, transfer of naive T cells from IRF8-deficient mice induced more severe colitis in *Rag*^{-/-} mice than T cell from normal controls. Furthermore, we report that IRF8 physically interacts with ROR γ t, resulting in inhibition of IL-17 transcription. These findings suggest that IRF8 has a suppressive role in the control of T_H17 differentiation and highlight the importance of intrinsic genetic programmes for the silencing of T_H17-dependent immune responses.

Results

IRF8 deficiency enhances T_H17-cell differentiation. To investigate the function of IRF8 in T cells, we first examined the expression of IRF8 in CD4⁺ T cells from normal or OT-II transgenic mice activated by different stimuli. We found that T-cell antigen receptor (TCR) engagement with anti-CD3 and anti-CD28 antibodies as well as stimulation of OT-II cells resulted in significant induction of IRF8 protein expression, as determined by western blotting (Supplementary Fig. S1a,b). Interestingly, IRF8 protein was more

stably expressed in naive CD4⁺ T cells polarized for 12 to 72 h under T_H17-inducing conditions compared with T_H1- or T_H2-inducing conditions (Supplementary Fig. S1a). To clarify how T_H17-polarizing conditions induce stable IRF8 expression, CD4⁺ cells were stimulated with TGF- β in the absence of TCR activation and the results showed that TGF- β clearly induced IRF8 expression at both 48 and 72 h (Supplementary Fig. S1c). In addition, mitogen-activated protein kinase inhibitors significantly blocked IRF8 protein expression induced by TCR activation (Supplementary Fig. S1d) and STAT3 mutant mice showed impaired IRF8 mRNA expression (Supplementary Fig. S1e). These data indicate that IRF8 is consistently elevated in activated CD4⁺ T cells and TCR signalling cascade is involved in induction of IRF8 expression.

We then assessed the contributions of IRF8 to T_H17 differentiation by studying CD4⁺ T cells from mice deficient in IRF8 due to a conventional KO of the gene (*Irif8*^{-/-} mice). Naive CD4⁺ T cells from *Irif8*^{-/-} or wild-type (WT) littermate mice were primed *in vitro* for 4 days under T_H0 or T_H17 polarizing conditions. The cells were then re-stimulated with phorbol myristate acetate (PMA)/ionomycin and examined for the percentages of IL-17-producing cells by intracellular staining using flow cytometry. Notably, the frequency of IL-17-producing cells generated from *Irif8*^{-/-} T-cell cultures was about threefold greater than cells from WT cultures (Fig. 1a). These observations correlated with enhanced IL-17 secretion by *Irif8*^{-/-} T_H cells generated under T_H17 polarizing conditions as determined by enzyme-linked immunosorbent assay (ELISA; Fig. 1b). In addition, IL-17-producing CD4⁺ T cells were significantly increased among lamina propria lymphocytes isolated from *Irif8*^{-/-} mice as compared with WT littermate controls following *in vitro* activation under T_H17 conditions or at the basal levels (Fig. 1c and Supplementary Fig. S2).

To determine whether the effects of IRF8 deficiency on T_H17 potential were cell type-specific, we next compared CD4⁺ T cells from mice expressing Lck-Cre that were WT (wt/wt) or homozygous (*fl/fl*) for a conditional allele of *Irif8* gene resulting in selective depletion of IRF8 in the T-cell compartment. Quantitative real-time reverse transcription(RT)-PCR (qPCR) analyses revealed dramatically lower levels of *Irif8* transcripts in sorted thymocyte subpopulations (DP, CD4SP, CD8SP) and splenic CD3⁺CD4⁺ cells from *Lck-Cre*⁺*Irif8*^{fl/fl} compared with cells from *Lck-Cre*⁺*Irif8*^{wt/wt} littermate control mice (Supplementary Fig. S3), confirming that the *Irif8* gene was efficiently deleted from *Lck-Cre*⁺*Irif8*^{fl/fl} T cells. Splenic and lymph node CD4⁺ T-cell subsets from mice with IRF8-deficient T cells as well as from mice homozygous for a conventional *Irif8* null allele were normal in number as well as in expression of the T-cell activation markers CD62L, CD44, CD25 and CD69. Expression of FOXP3 in thymic and peripheral lymph node T cells from mice of both genotypes was also similar (Supplementary Fig. S4), indicating that CD4⁺ T cells develop normally in the absence of IRF8. Naive CD4⁺ T cells from *Lck-Cre*⁺*Irif8*^{fl/fl} and *Lck-Cre*⁺*Irif8*^{wt/wt} littermate controls were subjected to T_H17 polarization. As expected, T_H cells from mice with a T cell-specific deficiency in IRF8 showed a remarkable increase in the generation of IL-17-producing cells (Fig. 1d) in association with significantly elevated levels of IL-17 secretion (Fig. 1e). These results excluded the possibility that the effects of IRF8 deficiency on T cells from mice with a conventional KO of the gene could be attributed to the altered activities of other subsets of IRF8-deficient cells, such as B cells, DC or macrophages. T_H17 cells generated from splenic T cells of *Irif8*^{-/-} mice comprised a major portion of the β -TCR⁺ CD4⁺ lymphocyte subset (Supplementary Fig. S5). Indeed, high levels of IRF8 protein expression were detected in CD4⁺ T cells polarized under T_H17 conditions as determined by immunoblot analyses (Fig. 1f). Accordingly, transcript levels of the iconic T_H17 cytokines IL-17A, IL-17F and IL-22 were enhanced in *Irif8*^{-/-} T_H17 cells (Fig. 1g). In contrast, T_H1 or T_H2 differentiation was not noticeably affected in *Irif8*^{-/-} T-cell cultures (Fig. 2a–c). In addition, [³H]-Thymidine incorporation assay showed that the

Figure 1 | Increased IL-17 production in IRF8 deficient T helper cells. (a) Naive CD4⁺ T cells from wild-type (WT) or *Irif8*^{-/-} mice were differentiated under T_H0 and T_H17 polarizing conditions for 4 days. Cells were then re-stimulated with PMA/ionomycin for 5 h, stained for intracellular IL-17 and IFN- γ , and analysed by flow cytometry. Representative fluorescence-activated cell sorting (FACS) dot plots gated on CD4⁺ cells and the percentages of IL-17-producing CD4⁺ cells are shown. Data are from one experiment representative of three independent experiments. Error bars, s.d. (b) The cells prepared in a, T_H1 and T_H2 polarizing conditions were re-stimulated with PMA/ionomycin for 24 h and the supernatants were analysed for IL-17 by ELISA. Data are from one experiment representative of three independent experiments. Error bars, s.d. (c) Wild-type or *Irif8*^{-/-} lamina propria lymphocytes (LPL) were differentiated under T_H17 conditions for 4 days. Cells were then re-stimulated with PMA/ionomycin for 5 h, stained for intracellular IL-17 and analysed by flow cytometry. (d) Naive CD4⁺ T cells from *Lck-Cre*⁺ *Irif8*^{wt/wt} and *Lck-Cre*⁺ *Irif8*^{fl/fl} mice were differentiated under T_H0 and T_H17 polarizing conditions for 4 days and the cells were re-stimulated with PMA/ionomycin for 5 h for staining of IL-17, IFN- γ and FOXP3. Representative FACS dot plots gated on CD4⁺ cells are shown. (e) The cells prepared in d were re-stimulated with PMA/ionomycin for 24 h and the supernatants were analysed for IL-17 by ELISA. Data are from one experiment representative of two independent experiments. Error bars, s.d. (f) Naive CD4⁺ T cells from C57BL/6 mice were differentiated under T_H0 and T_H17 conditions for 4 days. The cells were re-stimulated with PMA/ionomycin for 12 h and IRF8 expression was analysed by western blot. (g) The cells prepared in a were re-stimulated with PMA/ionomycin for 5 h and mRNA expression of indicated genes was determined by qPCR. The data shown were normalized to levels of ubiquitin expression as analysed by qPCR. The results are representative of three independent experiments. Error bars, s.d.

proliferation of CD4⁺ T cells from *Lck-Cre*⁺ *Irif8*^{fl/fl} and *Lck-Cre*⁺ *Irif8*^{wt/wt} mice cultured under T_H17 conditions was comparable (Fig. 2d). Taken together, these results indicate that IRF8 is induced during T-cell activation, and that T_H17 -cell differentiation is enhanced in cells deficient in IRF8.

T_{reg} cells and autocrine cytokines are not altered in *Irif8*^{-/-} mice. To understand whether alterations in T_{reg} cells might contribute to enhanced T_H17 differentiation in IRF8-deficient mice, we analysed FOXP3⁺ CD4⁺ T cells in these mice. There were no significant

differences between the FOXP3⁺ CD4⁺ T-cell populations of WT and *Irif8*^{-/-} mice under T_H17 - or T_{reg} -inducing conditions (Fig. 3a,b). Thus, the more efficient generation of *Irif8*^{-/-} T_H17 cells in response to the combined effects of TGF- β plus IL-6 was not because of alterations in TGF- β -derived T_{reg} suppression. IL-17-producing cells generated from *Irif8*^{-/-} T-cell cultures were greatly increased following stimulation with IL-6 plus TGF- β (Fig. 3c).

To further investigate how IRF8 affects T_H17 differentiation, naive WT and *Irif8*^{-/-} CD4⁺ T cells were subjected to T_H17 differentiation in the presence of IL-6, IL-23 and TGF- β , either alone or

Figure 2 | T_H1 and T_H2 differentiation in *Irf8*^{-/-} CD4⁺ T cells. Naive CD4⁺ T cells of WT or *Irf8*^{-/-} mice were differentiated under T_H1 or T_H2 polarizing conditions for 4 days. Cells were then re-stimulated with PMA/ionomycin for 5 h and stained for intracellular IFN-γ as a T_H1 marker (a) and IL-4 as a T_H2 marker (b) by flow cytometry. Data are from one experiment representative of three independent experiments. Error bars, s.d. The cells prepared in a and b were re-stimulated with PMA/ionomycin for 12 h and the supernatants were analysed for IFN-γ and IL-4 by ELISA (c). Data are from one experiment representative of three independent experiments. Error bars, s.d. Naive CD4⁺ T cells from spleens and lymph nodes of *Lck-Cre*⁺*Irf8*^{wt/wt} and *Lck-Cre*⁺*Irf8*^{fl/fl} mice were prepared and the cells were differentiated under T_H0 and T_H17 polarizing conditions for 3 days. [³H]-Thymidine was added during the last 8 h of culture. Then the cells were collected and were counted with a beta-counter (d). Data are from one experiment representative of two independent experiments. Error bars, s.d.

in various combinations, and then examined for the expression of lineage-specific genes by qPCR. Neither IL-6 nor TGF-β alone induced significant levels of IL-17 or IL-17F transcripts (Fig. 3d). In contrast, dramatic increases in IL-17 and IL-17F transcripts were induced at multiple time points by the combination of IL-23, IL-6 and TGF-β in cultures of both *Irf8*^{-/-} and WT T cells (Fig. 3d). Increased expression of IL-17 was confirmed at the protein level by ELISA (Fig. 3e–g). IL-23 induced low levels of IL-17 and IL-17F transcripts with no significant differences being seen between CD4⁺ T cells of WT and *Irf8*^{-/-} mice (Fig. 3d). This suggests that IRF8 may not target the IL-23 signalling cascade, but may exert a major influence on T_H17 differentiation instead of T_H17 expansion and maintenance.

IL-21, an autocrine cytokine produced by CD4⁺ T follicular helper cells and T_H17 cells^{6,7}, induces T_H17 differentiation in the presence of TGF-β. IRF8-deficient T cells displayed enhanced induction of T_H17-associated molecules following stimulation by TGF-β combined with IL-6 or IL-21 (Figs 3h and 4a). However, production of IL-21 and IL-10 was comparable in WT and *Irf8*^{-/-} T_H17 cells (Fig. 4b–e). These results suggest that an autocrine loop involving either IL-21 or IL-10 is not involved in the functional control of T_H17 differentiation by IRF8.

We next determined whether the induction of T_H17-associated genes may be affected by forced expression of IRF8 in T cells.

Retroviral transduction of IRF8-IRES-GFP into WT naive CD4⁺ T cells significantly decreased the percentage of IL-17-producing cells under T_H17 polarizing conditions (Fig. 5a), and RORγt-positive cells were moderately reduced (Fig. 5a). Similarly, retroviral transduction of IRF8 into the EL4T lymphoma cell line stimulated with PMA/ionomycin resulted in significantly reduced transcripts for IL-17, but had no effect on the expression of interferon (IFN)-γ, IL-4, IL-10 or FOXP3 (Fig. 5b). Taken together, these results demonstrate a direct role for IRF8 in suppressing T_H17-specific gene expression.

IRF8 interacts with RORγt and suppresses IL-17 transcription.

The above findings prompted us to probe the molecular basis for IRF8 control of T_H17-cell differentiation. As many studies have demonstrated a critical role for RORγt in T_H17-cell differentiation both *in vitro* and *in vivo*¹¹, we asked if IRF8 might affect RORγt-mediated IL-17 induction. EL4 cells were transiently transfected RORγt followed by stimulation with PMA/ionomycin. Overexpression of RORγt resulted in significantly increased expression of IL-17 and IL-17F, whereas co-transfection with IRF8 greatly reduced the expression of these genes, suggesting that IRF8 inhibits RORγt-induced expression of IL-17 transcripts (Fig. 6a). Using a 6-kbp IL-17 promoter reporter plasmid, we confirmed that RORγt strongly induced IL-17 promoter reporter activity in 293T cells (Fig. 6b),

Figure 3 | IRF8 inhibits the expression of T_H17-associated genes. Naive CD4⁺ T cells from wild-type and *Irif8*^{-/-} mice were stimulated with plate-bound anti-CD3 and anti-CD28 antibodies in the presence of IL-6 (10 ng ml⁻¹) plus differing concentrations of TGF- β (0.1, 1, 5 ng ml⁻¹) (a), or TGF- β (5 ng ml⁻¹)/IL-6 (10 ng ml⁻¹) with or without retinoic acid (RA, 100 nM) (b). After 4 days of stimulation, IL-17 and FOXP3 intracellular staining was performed and analysed by flow cytometry. (c) The cells prepared in a and b except for the presence of TGF- β (5 ng ml⁻¹) plus various concentrations of IL-6 (5, 10, 50, 100 ng ml⁻¹) were re-stimulated with PMA/ionomycin for 5 h and stained for intracellular IL-17 and IFN- γ and analysed by flow cytometry. (d) Naive CD4⁺ T cells from wild-type and *Irif8*^{-/-} mice were stimulated with IL-6, IL-23, TGF- β or different combinations of these cytokines for various intervals. Total RNA was extracted and analysed by RT-PCR for mRNA expression of IL-17 (top panel) and IL-17F (bottom panel). Data are from one experiment representative of three independent experiments. Error bars, s.d. (e) Naive CD4⁺ T cells from WT and *Irif8*^{-/-} mice were stimulated with the indicated cytokines for 4 days. Cells were re-stimulated with PMA/ionomycin for 5 h and stained for intracellular IL-17 and IFN- γ and analysed by flow cytometry. The results are representative of three independent experiments. (f) Cells were prepared as in e and the culture supernatants were collected after 4 days of stimulation. IL-17 protein secretion was analysed by ELISA. (g) Naive CD4⁺ T cells from WT and *Irif8*^{-/-} mice were prepared and stimulated with TGF- β and IL-6 in the presence of neutralizing anti-IFN- γ (10 μ g ml⁻¹) and anti-IL-4 (10 μ g ml⁻¹) antibodies. At 72 h after the stimulation, IL-17 protein secretion in culture supernatants was analysed by ELISA. Data are the mean $\pm$ s.d. of triplicate cultures. (h) Naive CD4⁺ T cells from wild-type or *Irif8*^{-/-} mice were stimulated with the indicated cytokine combinations for 48 h. Total RNA was extracted and analysed by qPCR for mRNA expression of the indicated genes. Data are from one experiment representative of three independent experiments. Error bars, s.d.

which was shown to be due to a direct effect of ROR γ t on the IL-17 promoter^{26,27}. Co-transfection of IRF8 in these cells suppressed ROR γ t-mediated IL-17 promoter activity in a dose-dependent manner (Fig. 6c). Similar results were observed in EL4 cells (Fig. 6d). CNS2 is a conserved non-coding sequence (CNS) element ~5 kbp upstream of the IL-17 locus that functions as a ROR γ t-dependent enhancer element required for optimal IL-17 transcription^{27,28}. Overexpression of IRF8 significantly suppressed CNS2-enhanced IL-17 promoter activity (Fig. 6e), indicating that IRF8 inhibits IL-17 transcription.

To investigate whether IRF8 can directly bind the CNS2 region of the IL-17 promoter, we co-transfected an IL-17 promoter reporter (containing CNS2) and IRF8 plasmids into 293T cells and performed chromatin immunoprecipitation (ChIP) using an

IRF8-specific antibody. The precipitated chromatin DNA was analysed by qPCR using primers covering the CNS2 region of the IL-17 promoter. The results showed that IRF8 antibody specifically pulled down the CNS2 region sequences (Fig. 6f). To confirm the results, naive WT and *Irif8*^{-/-} CD4⁺ T cells were stimulated under T_H17 polarizing conditions for 60 h and ChIP assay was performed as above. The precipitated chromatin DNA was analysed by PCR using primers covering the CNS2 region of the IL-17 promoter. Similarly, IRF8 antibody specifically pulled down the CNS2 region sequences of T_H17 cells from WT mice but not from *Irif8*^{-/-} mice (Fig. 6g). These results demonstrate that IRF8 bound directly to this region of the IL-17 promoter. Further analyses of the CNS2 sequence revealed several IRF consensus binding sequence elements (Supplementary Fig. S6a). We showed that mutation of a typical IRF-binding site

Figure 4 | IL-21 signalling and IL-10 production during T_H17-cell differentiation in *Irf8*^{-/-} mice. Naive CD4⁺ T cells from WT and *Irf8*^{-/-} mice were prepared, and the cells were stimulated with anti-CD3 and anti-CD28 antibodies in the presence or absence of TGF-β plus IL-21 or IL-6 for 72 h. Intracellular staining for IL-17 and IFN-γ expression was performed and analysed by flow cytometry (**a**). The secretion of IL-21 protein in culture supernatants was analysed by ELISA (**b**). Data are from one experiment representative of three independent experiments. Error bars, s.d. Naive CD4⁺ T cells were stimulated with anti-CD3 and anti-CD28 antibodies in the presence of TGF-β plus IL-6 or IL-21. At 96 h after stimulation, intracellular staining for IL-17 and IL-10 was performed and analysed after re-stimulation with PMA/ionomycin by flow cytometry (**c**). Naive CD4⁺ T cells from spleens and lymph nodes of WT and *Irf8*^{-/-} mice were prepared and the cells were stimulated under T_H0, T_H1, T_H2 and T_H17 polarizing conditions for 4 days. Then the cells were re-stimulated with PMA/ionomycin for 12 h and IL-10 protein secretion was analysed by ELISA (**d**). Data are from one experiment representative of three independent experiments. Error bars, s.d. Naive CD4⁺ T cells were stimulated with anti-CD3 and anti-CD28 antibodies in the presence of TGF-β plus IL-6 or IL-21 for 72 h. IL-10 expression was analysed by RT-PCR (**e**). Data are from one experiment representative of three independent experiments. Error bars, s.d.

in the CNS2 region of IL-17 promoter nullified the inhibition of RORγt-mediated IL-17 promoter activation by IRF8 (Supplementary Fig. S6b). These results provided a molecular basis for understanding the inhibitory effects of IRF8 on IL-17 transcription. It is likely that RORγt cooperates with other known transcrip-

tion factors, such as FOXP3 or RUNX1, or unknown factors in the coordinate regulation of T_H17 differentiation^{26,29,30}. We then co-transfected HA-tagged IRF8 and T7-RORγt plasmids into 293T cells for co-immunoprecipitation. Immunoprecipitation of RORγt resulted in co-precipitation of IRF8 (Fig. 7a), even in the presence of ethidium bromide or DNase I, indicating that IRF8 and RORγt interact with each other without the involvement of DNA. Using confocal microscopy, we also determined that IRF8 and RORγt co-localized in the nucleus of NIH3T3 cells co-transfected with GFP-IRF8 and T7-RORγt constructs (Fig. 7b). Flow cytometric analyses of naive CD4⁺ T cells stimulated under T_H17 polarizing conditions clearly revealed a population of RORγt⁺CD4⁺ cells with the majority of these cells also staining for IRF8 (Fig. 7c). Furthermore, anti-IRF8 antibodies were found to co-immunoprecipitate RORγt from lysates of primary CD4⁺ T cells from WT but not from *Irf8*^{-/-} mice cultured under T_H17 polarizing conditions (Fig. 7d), indicating that endogenous IRF8 and RORγt interact with each other. As shown in Figure 7e, IRF8 is comprised of an N-terminal DNA-binding domain, a flanking internal region, and a C-terminal IRF association domain³¹. To map the binding sites between IRF8 on RORγt, we co-transfected 293T cells with T7-tagged RORγt and Flag-tagged full-length IRF8 or one of a series of C-terminal truncation mutants (1–390, 1–356, 1–305, 1–253, 1–230, 1–190 and 1–154) followed by co-immunoprecipitation experiments. The results showed that IRF8 amino-acid residues between 230 and 190 were important for the physical interaction with RORγt (Supplementary Fig. S7a, Fig. 7f). IRF8 mutant (1–114), which is not bound to RORγt, did not suppress RORγt-mediated IL-17 promoter activation as WT and other binding mutants did (Fig. 7g and Supplementary Fig. S7b), demonstrating that the inhibitory activity of IRF8 on IL-17 transcription is related to its interaction with RORγt. In addition, IRF4, another IRF family member that is required for the generation of T_H17 cells³², was also found in co-immunoprecipitation experiments to interact with RORγt, whereas there was no interaction between IRF1 and RORγt (Supplementary Fig. S8a). Although transcript levels for IRF4 in cells deficient in IRF8 did not change under T_H17 polarizing conditions (Supplementary Fig. S8b), we still cannot exclude possibility that IRF8 and IRF4 mutually influence their activities in T_H17 cells. Thus, our results suggest that protein–protein interactions between IRF8 and RORγt have at least a partial role in IRF8-mediated inhibitory effects on IL-17 transcription.

IRF8 controls T_H17-cell differentiation *in vivo*. To further assess the effects of IRF8 on T_H17-cell differentiation *in vivo*, we performed adoptive transfer experiments using CD4⁺CD62L⁺CD45RB^{hi}CD25⁻ cells from WT and *Irf8*^{-/-} mice to induce colitis in RAG1 KO (*Rag1*^{-/-}) mice. *Irf8*^{-/-} mice did not develop spontaneous colitis during an observation period of 1.5 years (Fig. 8a,b). However, *Rag1*^{-/-} mice reconstituted with *Irf8*^{-/-} naive CD4⁺ T cells began losing weight earlier and lost more weight than mice in the control group. Parallel histological studies of colon sections from *Rag1*^{-/-} mice reconstituted with *Irf8*^{-/-} T cells revealed more severe inflammatory cell infiltrates and significantly higher pathological scores than those observed in sections from mice reconstituted with T cells from WT mice (Fig. 8c–e). In addition, mice reconstituted with *Irf8*^{-/-} cells had a significantly higher percentage of IL-17-producing cells than control mice (Fig. 8f). To determine whether T_{reg} cells from *Irf8*^{-/-} mice could suppress effector T cells, we examined the population of T_{reg} cells in *Rag1*^{-/-} mice after the transfer of naive WT or *Irf8*^{-/-} CD4⁺ T cells. *Rag1*^{-/-} recipients of naive CD4⁺ T cells from mice of either genotype generated a small percentage of CD4⁺FOXP3⁺ and CD4⁺IFN-γ⁺ cells in the mesenteric lymph nodes (Fig. 8f). Furthermore, co-transfers of naive WT CD4⁺ T cells with CD4⁺CD25⁺ cells purified from WT or *Irf8*^{-/-} mice resulted in similar effects on body weight (Supplementary Fig. S9). These results indicate that the effects of IRF8 deficiency on T cell-mediated inflammation

Figure 5 | Transduction of IRF8 inhibits T_H17 -associated genes. (a) Naive $CD4^+$ T cells from C57BL/6 mice were infected with retrovirus encoding IRF8 or empty vector and were activated under T_H17 -inducing conditions for 4 days. The cells were re-stimulated with PMA/ionomycin for 5 h and stained for intracellular IL-17 and RORγt and analysed by flow cytometry. (b) EL4 cells were transduced with retroviruses encoding IRF8 (black column) or GFP (white column) for 48 h and the transduced cells were then stimulated with PMA/ionomycin for various times as indicated. Total RNA was extracted and the transcript levels of T_H17 -associated genes were analysed by qPCR as indicated. The results are normalized to ubiquitin levels. Data are from one experiment representative of three independent experiments. Error bars, s.d.

could not be explained by influences on the function of $CD4^+$ T_{reg} cells. Thus, IRF8 deficiency promotes intestinal inflammation in a T cell-mediated model of colitis, suggesting that IRF8 may have an inhibitory role in the control of T_H17 -mediated immune responses. To further understand the role of IRF8 in T_H17 -cell differentiation *in vivo*, we extended our observations to an infection model, as T_H17 cells have also been proposed to have a role in inflammation against both intracellular and extracellular bacteria^{33,34}. *Staphylococcus aureus* is a Gram-positive bacterium that can induce IL-17 production from $CD4^+$ T cells mainly through *Staphylococcus aureus* enterotoxin A (SEA), and humans deficient in T_H17 cells are highly susceptible to infection with this agent^{35,36}. To better understand the regulatory effects of IRF8 on T_H17 -cell differentiation in a broader sense, we used superantigenic *S. aureus* to induce IL-17 production *in vivo*. Spleen cells from *Lck-Cre⁺ Irf8^{fl/fl}* and *Lck-Cre⁺ Irf8^{wt/wt}* littermate controls immunized with SEA 4 days previously were re-stimulated *in vitro* with SEA for an additional 2 days and then examined for IL-17-producing $CD4^+$ T cells by flow cytometry and for IL-17 secretion by ELISA (Supplementary Fig. S10). IRF8-deficient mice generated significantly more IL-17-producing $CD4^+$ T cells than WT mice, further confirming that IRF8 negatively regulates T_H17 -cell differentiation *in vivo*.

Discussion

T_H17 cells represent a recently defined member of a still growing family of T helper cells. The mechanisms involved in the silencing

programme for this T helper subset remain unclear. Here we demonstrate that IRF8 serves as an intrinsic silencer for T_H17 -cell differentiation. IRF8-deficiency in both conventional and T cell-specific conditional KO mice led to more robust T_H17 -cell differentiation without effects on either T_H1 or T_H2 cell lineages. Furthermore, transfer of IRF8^{-/-} $CD4^+$ $CD45Rb^{hi}$ cells into *Rag1^{-/-}* mice induced more severe colitis than transfer of WT $CD4^+$ $CD45Rb^{hi}$ cells. In addition, mice reconstituted with IRF8^{-/-} cells had a significantly higher percentage of IL-17-producing cells than mice reconstituted with WT cells. In addition, we showed that IRF8 physically interacts with RORγt resulting in suppression of IL-17 transcription. These results suggest that IRF8 negatively regulates the development of T_H17 immune response resulting in the control of inflammation.

Many studies have demonstrated that IRF8 has important functions in myeloid cells²⁴. Macrophages from *Irf8^{-/-}* mice did not produce IL-12 in response to IFN-γ and LPS. IRF8 regulates IL-12 expression by binding to the IL-12 p40 promoter region, acting in synergy with IRF1 to activate IL-12 p40 gene expression³⁷. In addition, IRF8 also induces the expression of other inflammatory proteins expressed by myeloid cells, including iNOS, IL-18 and IL-1 (refs 38–40). IRF8 protein levels are controlled in part by Cbl-mediated ubiquitylation and subsequent proteasomal degradation⁴¹. More recent studies have shown that transcriptional activation of IL-12p40 by IRF8 is enhanced following ubiquitylation by the E3 ubiquitin ligase, TRIM21 (ref. 42). Here, we showed that $CD4^+$ T cells clearly expressed IRF8 protein on TCR engagement and that

Figure 6 | IRF-8 represses IL-17 transcription. (a) IRF8-expressing EL4 cells (black column) or control cells (white column) were transiently transfected with a ROR γ t plasmid for 24 h, and the cells were treated with plate-bound anti-CD3 and anti-CD28 antibodies in the presence or absence of the indicated cytokines for 8 h. qPCR analyses of transcripts of the indicated genes were performed and the results were normalized to the levels of ubiquitin transcripts. ** $P < 0.01$ versus cells transfected with IRF8 (Student's t -test). Data are representative of multiple experiments. (b) 293T cells were co-transfected with an IL-17 promoter reporter construct containing a 6-kbp promoter and increasing doses of a ROR γ t plasmid for 30 h. (c) 293T cells were co-transfected with an IL-17 promoter reporter construct containing a 6-kbp promoter, a ROR γ t plasmid, a FOXP3 plasmid, and increasing doses of an IRF8 plasmid for 30 h. (d) IRF8-expressing EL4 cells (black column) or control cells (white column) were co-transfected with an IL-17 promoter reporter construct containing a 6-kbp promoter plus ROR γ t plasmid, and treated with plate-bound anti-CD3 antibody in the presence or absence of the indicated cytokines for 24 h. Luciferase activities in (b, c, d) were measured and normalized to β -galactosidase activity. Data are mean $\pm$ s.d. of triplicate cultures and are representative of three independent experiments. * $P < 0.05$ versus cells transfected with IRF8 (Student's t -test). (e) IRF8-expressing EL4 cells (black column) or control cells (white column) were co-transfected with an IL-17 promoter reporter construct containing a minimal 1.1-kb promoter plus CNS2, and ROR γ t plasmids. The cells were treated with plate-bound anti-CD3 and anti-CD28 antibodies and IL-6/TGF- β for 24 h. A luciferase assay was performed as described in (b), (c), and (d). * $P < 0.05$ versus cells transfected with IRF8 (Student's t -test). (f) 293T cells were co-transfected with an IL-17 promoter reporter construct (containing CNS2) with either an IRF8 overexpression plasmid or a control vector for 36 h, followed by ChIP analysis. 3 μ g of anti-IRF8 antibody or isotype-matched IgG as control antibody were used in the immunoprecipitation step. qPCR was used to quantify the amount of precipitated DNA with primers flanking the CNS2 region of the IL-17 promoter. Data were normalized to input DNA in each respective sample. (g) Naive CD4 $^{+}$ T cells from wild-type or *Irf8* $^{-/-}$ mice were cultured under T $_{H}17$ -polarizing conditions for 60 h, followed by ChIP assay as described above. 3 μ g of anti-IRF8 antibody or isotype-matched IgG as control antibody were used in the immunoprecipitation step. PCR was used to quantify the amount of precipitated DNA with primers flanking the CNS2 region of the IL-17 promoter.

T $_{H}17$ polarization conditions induced stable IRF8 protein expression. In addition, mitogen-activated protein kinase inhibitors completely blocked IRF8 protein expression induced by TCR activation and STAT3 mutant mice showed impaired IRF8 mRNA expression, indicating that the TCR signalling cascade is involved in induction of IRF8 expression. The percentages of CD4 $^{+}$ T cells in tissues of *Irf8* $^{-/-}$ and WT mice were comparable. Following stimulation under T $_{H}17$ polarizing conditions, however, the percentages of IL-17-producing CD4 $^{+}$ T cells were greatly increased and the expression of T $_{H}17$ signature genes was significantly enhanced for cells from *Irf8* $^{-/-}$ as compared with WT mice. These results suggest that IRF8 is an important transcription factor in controlling CD4 $^{+}$ T cell plasticity by targeting T $_{H}17$ -cell differentiation.

The balance between pathogenic T $_{H}17$ cells and suppressive T $_{reg}$ cells in the immune system depends on the presence of inflammatory cytokines such as IL-6 and IL-21 (ref. 43). The anti-inflammatory cytokine, TGF- β , combined with IL-6 or IL-21 can drive the conversion of a T $_{H}$ cell phenotype from TGF- β -induced

FOXP3-expressing T $_{reg}$ cells to ROR γ t-expressing T $_{H}17$ cells. TGF- β inhibits the expression of STAT4 and GATA3, thereby preventing the differentiation of T $_{H}1$ and T $_{H}2$ cells, respectively, and currently facilitating T $_{H}17$ -cell development. CD4 $^{+}$ T cells from WT and *Irf8* $^{-/-}$ mice yielded similar populations of FOXP3 $^{+}$ cells following stimulation under T $_{H}17$ - or T $_{reg}$ -inducing conditions. In addition, there were no significant differences between WT and *Irf8* $^{-/-}$ T $_{H}17$ cells in the production of the autocrine cytokines, IL-21 and IL-10. These results rule out the possibility that enhanced generation of T $_{H}17$ cells by CD4 $^{+}$ T cells from *Irf8* $^{-/-}$ mice stimulated with TGF- β plus IL-6 was due to alterations in TGF- β -derived T $_{reg}$ suppression or to an autocrine loop involving IL-21 or IL-10.

IRF8 acts as a transcriptional repressor or a transcriptional activator depending on the target DNA sequence and interactions with different partner proteins, including PU.1, E47 and other IRFs²⁵. We demonstrated that IRF8 interacts directly with ROR γ t, resulting in suppression of IL-17 transcription. The association of IRF8 with ROR γ t was not competed by FOXP3, another ROR γ t-binding

Figure 8 | Lack of IRF8 enhances the T_H17 immune response in experimental colitis. WT and *Lrf8*^{-/-} mice were maintained under specific pathogen free (SPF) conditions for up to 18 months. Mice were killed and intestines were removed for histological analysis. Histology of colon tissues (**a**) and disease score (**b**) from age-matched young (15 weeks) and old (17–18 months) WT and *Lrf8*^{-/-} mice (three to four mice in each group). Scale bars, 200 μm. CD4⁺CD45RB^{hi} T cells were purified from spleens and lymph nodes of wild-type or *Lrf8*^{-/-} mice and 5 × 10⁵ cells were injected (i.p.) into recipient *Rag1*^{-/-} mice. Body weight change was monitored every week and mice were killed 7 weeks later. (**c**) Changes in body weight of *Rag1*^{-/-} mice (*n* = 5–6 mice per group) after intraperitoneal transfer of WT or *Lrf8*^{-/-} CD4⁺CD45RB^{hi} T cells were recorded. Data are presented as the mean ± s.d. of the percentage of initial body weight and are representative of two similar experiments. *P < 0.05 versus recipients of WT cells (ANOVA test and Student's *t*-test). Disease scores (**d**) and sections of colons with colitis (**e**) from *Rag1*^{-/-} mice (*n* = 5–6 mice in each group) on day 35 after naive T cell transfer as described in **c**. *P < 0.05 versus recipients of WT cells (Mann-Whitney test). Scale bars, 200 μm. (**f**) The percentage of IL-17, IFN-γ and FOXP3-producing cells from mesenteric lymph nodes of *Rag1*^{-/-} mice in **c** (white column, transfer with WT cells; black column, transfer with *Lrf8*^{-/-} cells). **P < 0.01 versus wild-type cell transferred mice (Student's *t*-test). Data are presented as the mean ± s.d. from four mice in each group. Two independent experiments were performed with similar results.

iCycler PCR with SYBR Green PCR master Mix (Applied Biosystems). Results were normalized based on the expression of ubiquitin. The following primer sets were used: IL-17A, IL-17F, IL-21, IL-23R, RORγt, IFN-γ, IRF8, IRF1, IRF4, T-bet, FOXP3, CCR6, IL-10, IL-4, IL-2 and ubiquitin (Supplementary Table S1).

Transfection and luciferase reporter assay. 293T cells were transiently transfected with an IL-17 promoter luciferase reporter plasmid together with RORγt in the presence of IRF8 plasmid at different concentrations. For each transfection, 2.0 μg of plasmid was mixed with 100 μl of Opti-MEM I medium (without serum and antibiotics) and 4.0 μl of Lipofectamine 2000 reagent. The mixture was incubated at room temperature for 20 min and added to 12-well plates containing cells and complete medium. The cells were incubated for 30 h and collected using reporter lysis buffer (Promega) for determination of luciferase activity. Cells were co-transfected with a β-galactosidase reporter plasmid to normalize experiments for transfection efficiency.

Generation of the mutant IL-17 promoters. A predicted IRF-binding site adjacent to the downstream RORγt-binding site in the CNS2 region of the mouse IL-17A promoter was mutated using QuickChange XL Site-Directed Mutagenesis Kit (Stratagene) according to manufacture's instruction. Two mutations were introduced on the IL-17 promoter regions - 5204 to - 5202 (TGG to CCC) and - 5201 to - 5199 (AAA to CCC) from the transcriptional initiation site (+ 1) using the following primer sets: 5'-GGTTGGAAAAAACCCTTCTGACCCA-3' and 5'-TGGGTCAGAAAACCTTGGGTTTTCCTTCAACC-3' for TGG to CCC; 5'-TGGAAAAAATGGCCGTTTCTGACCCACT-3' and 5'-AGTGGGTCAGAAAACGGGCCATTTTTCCTTCA-3' for AAA to CCC. Mutations were verified by sequencing.

Retroviral transduction of IRF8 in CD4⁺ T cells. To prepare pseudotyped virus human 293 EbnaT cells were seeded at a density of 4 × 10⁶ cells in a 10-cm dish. The

next day, cells were transfected using calcium phosphate with a mixture of 2.5 μg of plasmid pMD.G encoding vesicular stomatitis virus G protein, 7.5 μg of plasmid encoding gag-pol, and 10 μg of a retroviral expression constructs encoding GFP or IRF8. At 48 h after transfection, the viral supernatant was collected, centrifuged at 800g, and used to infect cells. CD4⁺ T-cell transduction was performed as previously described¹¹, sorted naive CD4⁺ T cells were plated as above and cultured for 24 h in the presence of anti-CD3 (1 μg ml⁻¹; 145-2C11, BD Biosciences) and soluble anti-CD28 (2 μg ml⁻¹; 37.51; BD Biosciences). Activated cells were then transduced with fresh retrovirus supernatant by centrifugation for 1.5 h at 2000g in the presence of polybrene (6 μg ml⁻¹; Sigma). After 24 h, the cells were re-transduced using the same procedure and cultured for an additional 24 h and were then stimulated with TGF-β and IL-6. The cells were collected on day 5 or 6 for intracellular cytokine staining.

T-cell proliferation assay. Naive CD4⁺ T cells were purified from spleens and lymph nodes of *Lck-Cre⁺ Irf8^{fl/fl}* and *Lck-Cre⁺ Irf8^{wt/wt}* littermate controls. Cells (1 × 10⁵ per well) were cultured in the absence or presence of anti-CD3 (1 μg ml⁻¹) and anti-CD28 (2 μg ml⁻¹) antibodies for 3 days in 96-well microplates. [³H]-Thymidine was added during the last 8 h of a 72-h culture. The cells were then collected and counted with a beta-counter.

Co-immunoprecipitation and immunoblotting analysis. Cells were washed with cold phosphate-buffered saline and lysed for 15 min on ice in 0.5 ml of lysis buffer (50 mM Tris-HCl, pH 8.0, 280 mM NaCl, 0.5% Nonidet P-40, 0.2 mM EDTA, 2 mM EGTA, 10% glycerol and 1 mM dithiothreitol) containing protease inhibitors. Cell lysates were clarified by centrifugation (4°C, 15 min, 20,000 r.p.m.), aliquots (500 μg) were incubated with 2 μg of normal rabbit IgG for 4 h and 20 μl of protein G-Sepharose was added to the mixture for 2 h. After centrifugation, the supernatant was collected and incubated with 2 μg of anti-Flag antibody overnight at 4°C with gentle rocking, after which immune complexes were collected as described

above with 20 μ l of protein G-Sepharose. After washing five times with lysis buffer, immunoblotting was performed. Anti-Flag (Sigma), anti-IRF8 (Santa Cruz), anti- β -actin (Sigma) and anti-T7 (MBL) antibodies were used according to the manufacturers' instructions. Secondary antibodies were from Santa Cruz.

T-cell-transfer colitis studies and histopathology. T-cell-transfer colitis was performed as previously described^{49,50}. Briefly, purified CD4⁺CD45RB^{hi} T cells from WT and *Irfl8*^{-/-} mice were injected intraperitoneally into *Rag1*^{-/-} recipients (5 $\times$ 10⁵ cells per mouse in 200 μ l sterile PBS per injection). Mice were weighed every week throughout the course of experiments. After 5 weeks, mice were killed and colon tissues were excised. Tissues were fixed in 10% buffered formalin and paraffin embedded. The sections (5 μ m) of tissue samples stained with hematoxylin and eosin. All the slides were read and scored by an experienced pathologist (L.Q.) without previous knowledge of the type of treatment. The degree of inflammation in the epithelium, submucosa and muscularis propria was scored separately as described by Totsuka *et al.*⁴⁹

In vivo stimulation of T_H17 cells by *Staphylococcus aureus*. *Lck-Cre⁺Irfl8^{fl/fl}* mice and *Lck-Cre⁺Irfl8^{wt/wt}* littermate controls were immunized (i.p.) with SEA (10 μ g per mouse) for 4 days. Mice were killed and spleen cells were prepared. The cells were re-stimulated *in vitro* with SEA (10 ng ml⁻¹) for additional 2 days and cells were collected for the analysis of IL-17-producing CD4⁺ T cells by flow cytometry gating on CD4⁺ T cells. The supernatants were collected for the measurement of IL-17 production by ELISA.

Chromatin immunoprecipitation assay. The ChIP procedure was performed using an assay kit following the manufacturer's instruction (Upstate Biotechnology). Briefly, 293T cells were co-transfected with an IL-17 promoter containing CNS2 domain and an IRF8 plasmid for 36 h and 1 $\times$ 10⁷ transfected cells were then crosslinked by 1% formaldehyde for 10 min at 37°C. Nuclei were prepared and subjected to sonication to obtain DNA fragments. Chromatin fractions were pre-cleared with protein A-agarose beads followed by immunoprecipitation overnight at 4°C with 3 μ g of anti-IRF8 or control antibody. Crosslinking was reversed at 65°C for 4 h, followed by proteinase K digestion. DNA was purified and subjected to qPCR. The input DNA was diluted 200 times before PCR amplification. The input and immunoprecipitated DNA were amplified by qPCR using primers (5'-CAGCCCTGGTCCTTAACTG-3' and 5'-TCACTTTCGTTGTCCTTTG-3') encompassing the CNS2 region of the mouse IL-17 promoter.

Cytokine ELISA. Supernatants from cell cultures were collected after activation under various conditions and secreted cytokines in the supernatants were measured by ELISA kits with purified coating and biotinylated detection antibodies: anti-IL-17, anti-22 and anti-IL-21 (R&D systems), anti-IFN- γ , anti-IL-4 and anti-IL-10 (BD Bioscience).

Statistical analysis. Statistical analysis was performed using Student's *t*-test for most of the experiments. For colitis experiments, Man-Whitney test and analysis of variance test were used for comparison of disease score and weight loss. *P* values < 0.05 were considered statistically significant.

References

- Bettelli, E., Korn, T. & Kuchroo, V. K. Th17: the third member of the effector T cell trilogy. *Curr. Opin. Immunol.* **19**, 652–657 (2007).
- Weaver, C. T., Harrington, L. E., Mangan, P. R., Gavrieli, M. & Murphy, K. M. Th17: an effector CD4⁺ T cell lineage with regulatory T cell ties. *Immunity* **24**, 677–688 (2006).
- Zheng, Y. *et al.* Interleukin-22, a T(H)17 cytokine, mediates IL-23-induced dermal inflammation and acanthosis. *Nature* **445**, 648–651 (2007).
- Liang, S. C. *et al.* Interleukin (IL)-22 and IL-17 are coexpressed by Th17 cells and cooperatively enhance expression of antimicrobial peptides. *J. Exp. Med.* **203**, 2271–2279 (2006).
- Veldhoen, M., Hocking, R. J., Atkins, C. J., Locksley, R. M. & Stockinger, B. TGF β in the context of an inflammatory cytokine milieu supports de novo differentiation of IL-17-producing T cells. *Immunity* **24**, 179–189 (2006).
- Korn, T. *et al.* IL-21 initiates an alternative pathway to induce proinflammatory T(H)17 cells. *Nature* **448**, 484–487 (2007).
- Nurieva, R. *et al.* Essential autocrine regulation by IL-21 in the generation of inflammatory T cells. *Nature* **448**, 480–483 (2007).
- Zhou, L. *et al.* IL-6 programs T(H)-17 cell differentiation by promoting sequential engagement of the IL-21 and IL-23 pathways. *Nat. Immunol.* **8**, 967–974 (2007).
- Mangan, P. R. *et al.* Transforming growth factor- β induces development of the T(H)17 lineage. *Nature* **441**, 231–234 (2006).
- Bettelli, E. *et al.* Reciprocal developmental pathways for the generation of pathogenic effector Th17 and regulatory T cells. *Nature* **441**, 235–238 (2006).
- Ivanov, I. I. *et al.* The orphan nuclear receptor ROR γ directs the differentiation program of proinflammatory IL-17⁺ T helper cells. *Cell* **126**, 1121–1133 (2006).
- McGeachy, M. J. *et al.* The interleukin 23 receptor is essential for the terminal differentiation of interleukin 17-producing effector T helper cells in vivo. *Nat. Immunol.* **10**, 314–324 (2009).
- Yang, L. *et al.* IL-21 and TGF- β are required for differentiation of human T(H)17 cells. *Nature* **454**, 350–352 (2008).
- Harrington, L. E. *et al.* Interleukin 17-producing CD4⁺ effector T cells develop via a lineage distinct from the T helper type 1 and 2 lineages. *Nat. Immunol.* **6**, 1123–1132 (2005).
- Stumhofer, J. S. *et al.* Interleukin 27 negatively regulates the development of interleukin 17-producing T helper cells during chronic inflammation of the central nervous system. *Nat. Immunol.* **7**, 937–945 (2006).
- Korn, T., Bettelli, E., Oukka, M. & Kuchroo, V. K. IL-17 and Th17 Cells. *Annu. Rev. Immunol.* **27**, 485–517 (2009).
- Lee, C. H. *et al.* Regulation of the germinal center gene program by interferon (IFN) regulatory factor 8/IFN consensus sequence-binding protein. *J. Exp. Med.* **203**, 63–72 (2006).
- Zhao, J. *et al.* IRF-8/interferon (IFN) consensus sequence-binding protein is involved in Toll-like receptor (TLR) signaling and contributes to the cross-talk between TLR and IFN- γ signaling pathways. *J. Biol. Chem.* **281**, 10073–10080 (2006).
- Tsujimura, H., Tamura, T. & Ozato, K. Cutting edge: IFN consensus sequence binding protein/IFN regulatory factor 8 drives the development of type I IFN-producing plasmacytoid dendritic cells. *J. Immunol.* **170**, 1131–1135 (2003).
- Driggers, P. H. *et al.* An interferon gamma-regulated protein that binds the interferon-inducible enhancer element of major histocompatibility complex class I genes. *Proc. Natl Acad. Sci. USA* **87**, 3743–3747 (1990).
- Barber, S. A., Fultz, M. J., Salkowski, C. A. & Vogel, S. N. Differential expression of interferon regulatory factor 1 (IRF-1), IRF-2, and interferon consensus sequence binding protein genes in lipopolysaccharide (LPS)-responsive and LPS-hyporesponsive macrophages. *Infect. Immun.* **63**, 601–608 (1995).
- Wang, H. & Morse, H. C. III IRF8 regulates myeloid and B lymphoid lineage diversification. *Immunol. Res.* **43**, 109–117 (2009).
- Tamura, T., Nagamura-Inoue, T., Shmeltzer, Z., Kuwata, T. & Ozato, K. ICSBP directs bipotential myeloid progenitor cells to differentiate into mature macrophages. *Immunity* **13**, 155–165 (2000).
- Gabriele, L. & Ozato, K. The role of the interferon regulatory factor (IRF) family in dendritic cell development and function. *Cytokine Growth Factor Rev.* **18**, 503–510 (2007).
- Tamura, T., Yanai, H., Savitsky, D. & Taniguchi, T. The IRF family transcription factors in immunity and oncogenesis. *Annu. Rev. Immunol.* **26**, 535–584 (2008).
- Ichiyama, K. *et al.* Foxp3 inhibits ROR γ -mediated IL-17A mRNA transcription through direct interaction with ROR γ . *J. Biol. Chem.* **283**, 17003–17008 (2008).
- Zhang, F., Meng, G. & Strober, W. Interactions among the transcription factors Runx1, ROR γ and Foxp3 regulate the differentiation of interleukin 17-producing T cells. *Nat. Immunol.* **9**, 1297–1306 (2008).
- Akimzhanov, A. M., Yang, X. O. & Dong, C. Chromatin remodeling of interleukin-17 (IL-17)-IL-17F cytokine gene locus during inflammatory helper T cell differentiation. *J. Biol. Chem.* **282**, 5969–5972 (2007).
- Zhang, Z., Zheng, M., Bindas, J., Schwarzenberger, P. & Kolls, J. K. Critical role of IL-17 receptor signaling in acute TNBS-induced colitis. *Inflamm. Bowel Dis.* **12**, 382–388 (2006).
- Schraml, B. U. *et al.* The AP-1 transcription factor Batf controls T(H)17 differentiation. *Nature* **460**, 405–409 (2009).
- Ozato, K., Tailor, P. & Kubota, T. The interferon regulatory factor family in host defense: mechanism of action. *J. Biol. Chem.* **282**, 20065–20069 (2007).
- Brustle, A. *et al.* The development of inflammatory T(H)-17 cells requires interferon-regulatory factor 4. *Nat. Immunol.* **8**, 958–966 (2007).
- Miossec, P., Korn, T. & Kuchroo, V. K. Interleukin-17 and type 17 helper T cells. *N. Engl. J. Med.* **361**, 888–898 (2009).
- Matsuzaki, G. & Umemura, M. Interleukin-17 as an effector molecule of innate and acquired immunity against infections. *Microbiol. Immunol.* **51**, 1139–1147 (2007).
- Ma, C. S. *et al.* Deficiency of Th17 cells in hyper IgE syndrome due to mutations in STAT3. *J. Exp. Med.* **205**, 1551–1557 (2008).
- Islander, U. *et al.* Superantigenic *Staphylococcus aureus* stimulates production of interleukin-17 from memory but not naive T cells. *Infect. Immun.* **78**, 381–386 (2010).
- Masumi, A., Tamaoki, S., Wang, I. M., Ozato, K. & Komuro, K. IRF-8/ICSBP and IRF-1 cooperatively stimulate mouse IL-12 promoter activity in macrophages. *FEBS Lett.* **531**, 348–353 (2002).
- Xiong, H. *et al.* Complex formation of the interferon (IFN) consensus sequence-binding protein with IRF-1 is essential for murine macrophage IFN- γ -induced iNOS gene expression. *J. Biol. Chem.* **278**, 2271–2277 (2003).
- Kim, Y. M., Im, J. Y., Han, S. H., Kang, H. S. & Choi, I. IFN- γ up-regulates IL-18 gene expression via IFN consensus sequence-binding protein and activator protein-1 elements in macrophages. *J. Immunol.* **165**, 3198–3205 (2000).
- Marecki, S., Riendeau, C. J., Liang, M. D. & Fenton, M. J. PU.1 and multiple IFN regulatory factor proteins synergize to mediate transcriptional activation of the human IL-1 β gene. *J. Immunol.* **166**, 6829–6838 (2001).

41. Xiong, H. *et al.* Ubiquitin-dependent degradation of interferon regulatory factor-8 mediated by Cbl down-regulates interleukin-12 expression. *J. Biol. Chem.* **280**, 23531–23539 (2005).
42. Kong, H. J. *et al.* Cutting edge: autoantigen Ro52 is an interferon inducible E3 ligase that ubiquitinates IRF-8 and enhances cytokine expression in macrophages. *J. Immunol.* **179**, 26–30 (2007).
43. Bettelli, E. & Kuchroo, V. K. IL-12- and IL-23-induced T helper cell subsets: birds of the same feather flock together. *J. Exp. Med.* **201**, 169–171 (2005).
44. Zhou, L. *et al.* TGF-beta-induced Foxp3 inhibits T(H)17 cell differentiation by antagonizing RORgamma function. *Nature* **453**, 236–240 (2008).
45. Leppkes, M. *et al.* RORgamma-expressing Th17 cells induce murine chronic intestinal inflammation via redundant effects of IL-17A and IL-17F. *Gastroenterology* **136**, 257–267 (2009).
46. Seiderer, J. *et al.* Role of the novel Th17 cytokine IL-17F in inflammatory bowel disease (IBD): upregulated colonic IL-17F expression in active Crohn's disease and analysis of the IL17F p.His161Arg polymorphism in IBD. *Inflamm. Bowel Dis.* **14**, 437–445 (2008).
47. O'Connor, W. Jr *et al.* A protective function for interleukin 17A in T cell-mediated intestinal inflammation. *Nat. Immunol.* **10**, 603–609 (2009).
48. De Jager, P. L. *et al.* Meta-analysis of genome scans and replication identify CD6, IRF8 and TNFRSF1A as new multiple sclerosis susceptibility loci. *Nat. Genet.* **41**, 776–782 (2009).
49. Totsuka, T. *et al.* IL-7 is essential for the development and the persistence of chronic colitis. *J. Immunol.* **178**, 4737–4748 (2007).
50. Powrie, F., Leach, M. W., Mauze, S., Caddle, L. B. & Coffman, R. L. Phenotypically distinct subsets of CD4⁺ T cells induce or protect from chronic intestinal inflammation in C. B-17 scid mice. *Int. Immunol.* **5**, 1461–1471 (1993).

Acknowledgments

We thank C. Dong, A. Yoshimura, T. Taniguchi, O. Sakaguchi, M. Ono, H. Wang, J. Yang and G. Lu for providing reagents, technical supports and helpful suggestions. We are

grateful to J. Unkeless, K. Honda and J. Blander for critical reading of the manuscript. This work was supported by the NIH P01 DK072201, a Crohn's and Colitis Foundation of America grant, a grant from the Eli and Edythe L. Broad Foundation, and by the intramural research programme of the NIH, National Institute of Allergy and Infectious Diseases and the Eunice Kennedy Shriver National Institute of Child Health and Human Development.

Author contributions

X.O. and H.X. initiated the project and designed the experiments. X.O. and R.Z. performed the majority of the experiments. J.Y., Q.L. and C.Z. performed experiments. L.Q. did histology analysis. J.L., H.N. performed ChIP assay experiments. M.S.S., C.-F.Q. and H.C.M. provided IRF8 conditional KO mice. M.G. and K.O. generated various IRF8 mutant plasmids. J.C.H., S.A.L., and L.M. provided important reagents and critical discussion. X.O. and H.X. wrote the manuscript.

Additional information

Supplementary Information accompanies this paper at <http://www.nature.com/naturecommunications>

Competing financial interests: The authors declare no competing financial interests.

Reprints and permission information is available online at <http://npg.nature.com/reprintsandpermissions/>

How to cite this article: Ouyang, X. *et al.* Transcription factor IRF8 directs a silencing programme for T_H17-cell differentiation. *Nat. Commun.* **2**:314 doi: 10.1038/ncomms1311 (2011).

License: This work is licensed under a Creative Commons Attribution-NonCommercial-Share Alike 3.0 Unported License. To view a copy of this license, visit <http://creativecommons.org/licenses/by-nc-sa/3.0/>