ARTICLE

Received 8 Aug 2016 | Accepted 4 Oct 2016 | Published 28 Nov 2016

DOI: 10.1038/ncomms13431

OPEN

Borane catalysed ring opening and closing cascades of furans leading to silicon functionalized synthetic intermediates

Chinmoy K. Hazra^{1,2}, Narasimhulu Gandhamsetty^{1,2}, Sehoon Park^{1,2} & Sukbok Chang^{1,2}

The conversion of renewable biomass resources to synthetically valuable chemicals is highly desirable, but remains a formidable challenge in regards to the substrate scope and reaction conditions. Here we present the development of tris(pentafluorophenyl)borane-catalysed conversion of furans via ring-opening and closing cascade processes to afford silicon-functionalized synthetic chemicals under transition metal-free conditions. The furan ring-opening with hydrosilanes is highly efficient (TON up to 2,000) and atom-economical without forming any byproduct to give rise to α -silyloxy-(Z)-alkenyl silanes. Additional equivalents of silane smoothly induce a subsequent $B(C_6F_5)_3$ -catalysed cyclization of initially formed olefinic silane compounds to produce *anti*-(2-alkyl)cyclopropyl silanes, another versatile synthon being potentially applicable in the synthesis of natural products and pharmacophores.

¹Center for Catalytic Hydrocarbon Functionalizations, Institute for Basic Science (IBS), Daejeon 305-701, South Korea. ²Department of Chemistry, Korea Advanced Institute of Science & Technology (KAIST), Daejeon 305-701, South Korea. Correspondence and requests for materials should be addressed to S.C. (email: sbchang@kaist.ac.kr).

Production of furans and their subsequent transformations have received a great attention^{1,2}. This research activity can be attributed to the relevance of furan derivatives to the renewable biomass. Thermal dehydration of glucose or fructose, the most abundant monosaccharides, in the presence of acid catalysts provides important nonpetroleum chemicals including furfural and 5-hydroxymethylfurfural (HMF)^{3–6}. These compounds are considered to be versatile platform chemicals with high potential utility in organic synthesis and industrial applications to give rise to a diverse range of furans and their derivatives. In fact, a number of catalytic transformations of furans or their congeners to valuable chemicals have long been developed in both academia and industry^{7–9}. Among the precedents, however, the ring-opening reactions of furans are relatively less highlighted^{10–12} when compared with the derivatization of furans maintaining the ring skeleton^{13–15}. One notable example of the ring-opening of furans is the production of levulinic acid via an acid-catalysed hydration of HMF (Fig. 1a)^{16,17}. Recently, Gordon and coworkers showed that a sequential catalysis involving ring-opening hydrolysis of chain-extended HMF derivatives followed by hydrode-oxygenation using a combination of acid and metal catalyst can produce linear alkanes¹⁸. In addition, while a borane catalyst system was proved to be applicable in the transformations of carbohydrates, it was also employed for the ring-opening of furan derivatives. For instance, Gagné *et al.* found that B(C₆F₅)₃ catalyses the deoxygenation of carbohydrate-based polyols with hydrosilanes to give chiral-alcohol synthons¹⁹ or hydrocarbons^{20,21} with remarkable chemoselectivity. Njardarson *et al.* employed B(C₆F₅)₃ catalyst for the reduction of 2,5-dihydrofurans using hydrosilanes to obtain silyl(homo)allyl ethers, which was proposed to form via a hydrosilylative C–O bond cleavage²². On the other hand, Ashley *et al.* reported that hydrogenation of furan derivatives under the frustrated Lewis pairs catalysis using (B(C₆Cl₅)(C₆F₅)₂) delivers reduced tetrahydrofuran compounds²³. Despite of such progresses on furan transformation reactions, there have been no developments thus far for the selective conversion of furans to ring-opened products bearing a sp³ C–Si bond at ambient conditions.

Described herein is the development of a boron-catalysed cascade silylative transformation of furans (**I**) involving selective ring-opening and closing processes, thus allowing for the sequential formation of two types of new silylated products of α -silyloxy-(*Z*)-alkenyl silanes (**II**) and *trans*-(2-alkyl)cyclopropyl silanes (**IV**) (Fig. 1b). In this transformation, several aspects are especially notable: First, while the B(C₆F₅)₃-catalysed hydrosilylation of olefins is known²⁴, the sp³ C–O and sp² C=C bonds in the initially ring-opened α -silyloxy-(*Z*)-alkenyl silanes (**II**) remain unreacted. Second, the subsequent ring-closing of **II** is readily enabled simply by additional equivalent of hydrosilane leading to silylcyclopropanes (**IV**) without the need to isolate the ring-opened compounds **II**. Third, the observed high regio and stereoselectivities in both the initial ring-opening and subsequent cyclopropanation processes are an outcome of kinetic differentiation within the borane catalytic cycle. In addition, both types of products obtained from the present furan conversion are synthetically highly valuable compounds possessing sp³ C–Si bonds^{25–27}. Also, a broad range of furan congeners derived from 5-hydroxymethylfurfural are successfully applied for the selective cascade hydrosilylation to yield silicon-functionalized products (Fig. 1c).

Results

Discovery of new cascade transformations of furans. When 2-methylfuran was treated with 1.0 equiv. of PhMe₂SiH in the presence of B(C₆F₅)₃ (5.0 mol%) in dichloromethane solvent, a

new product, α -silyloxy-3-pentenyl silane **Z-2a**, was observed to form in 10 min at –78 °C along with unreacted **1a** in a 1:1 ratio (Fig. 2a; see Supplementary Fig. 1). Subsequent addition of 2.0 equivalents of PhMe₂SiH into the above reaction mixture brought about exclusive formation of silylated cyclopropane as a single stereoisomer (*anti*-**3a**) in 74% yield over 16 h in addition to (PhMe₂Si)₂O byproduct **4** (Fig. 2a; see Supplementary Fig. 2). These results indicate that the kinetic barrier for the B(C₆F₅)₃-catalysed ring-opening hydrosilylation of **1a** to **Z-2a** is much lower than that of the ring-closing process from **Z-2a** to *anti*-**3a**. Consistent with this interpretation, we were able to confirm that **1a** was quantitatively converted to **Z-2a** with 2.0 equivalents of PhMe₂SiH, while the use of 3.0 equiv. of the silane led to the exclusive formation of *anti*-**3a** together with stoichiometric byproduct **4** (Fig. 2b).

To gain more insight into the kinetics of each step of the cascade hydrosilylation processes, we monitored the reaction progress by ¹H NMR spectroscopy (Fig. 2c,d; see Supplementary Fig. 3). A precooled mixture of B(C₆F₅)₃, **1a**, and PhMe₂SiH (1:50:200) in CD₂Cl₂ at –70 °C was found to give **Z-2a** in a quantitative NMR yield over 3.5 h. On further warming up to 25 °C, **Z-2a** started to subsequently undergo a ring-closing process to afford silylated cyclopropane *anti*-**3a** (53% NMR yield in 7 h). These results clearly indicate that the furan transformation proceeds via sequential reductive pathways, which are strongly governed by kinetic factors. Moreover, the initial rates for each process in the hydrosilylation cascade of 2-methylfuran were determined to be $1.14 \times 10^{-4} \text{ M s}^{-1}$ (for the ring-opening at –70 °C) and $3.06 \times 10^{-5} \text{ M s}^{-1}$ (for the cyclopropanation at 25 °C) (Fig. 2d; Supplementary Fig. 3).

Substrate scope of the B(C₆F₅)₃-catalysed cascade silylative reduction of furans. With the preliminary results on the 2-methylfuran conversion in hand, we carried out additional optimization studies (Supplementary Tables 1–3) and investigated the substrate scope of this catalysis. 2-Alkyl or 2-arylfuran derivatives employed in this study were easily prepared via the Pd-catalysed Suzuki–Miyaura cross-coupling reaction of (2-furanyl)boronic acid with the corresponding alkyl or aryl halides (Supplementary Methods, GP1). In the present cascade transformation of furans, several notable features were revealed: (i) only equimolar amounts of hydrosilanes are needed to enable the ring-opening process, thus indicating that the conversion is atom-economical; (ii) the reaction is highly stereoselective in that one isomeric products are formed; (iii) the subsequent cyclopropanation process can be carried out without isolation of ring-opened intermediates, α -silyloxy-(*Z*)-homoallylsilanes; (iv) again, the ring-closing process is highly stereoselective affording *anti*-products exclusively; and (v) the overall procedure is convenient and easy to scale up with high catalyst turnovers (~2,000).

We were pleased to observe that the optimized conditions were readily applicable to a broad range of furan derivatives to give α -silyloxy-(*Z*)-homoallylsilanes (Table 1; Conditions A). Substrates with aliphatic substituents on the furan C-2 were all compatible with the B(C₆F₅)₃-catalysed (2.0 mol%) ring-opening process at ambient temperature. Analytically pure α -silyloxy-(*Z*)-homoallylsilane products (**Z-2a**~**2d**) were obtained in high yields and with excellent stereoselectivities (*Z*/*E*: >99/1, as measured by crude ¹H NMR experiments). The reaction of aryl-substituted furans proceeded with a similar level of selectivity and efficiency irrespective of the electronic properties of the aryl moiety (**Z-2e**~**2k**, *Z*/*E*: >99/1, 83–92%). Gratifyingly, potentially reactive functional groups (**Z-2l**~**2n**) were tolerated under the reaction conditions, while reactions of the furan substrates

Figure 1 | Furans from biomass resources and their transformations. (a) Furan platform chemicals and known post-transformations leading to diverse synthetic valuables. (b,c) Present work; Borane-catalysed ring-opening and closing cascades of furans giving rise to synthetically valuable silicon compounds in reaction with hydrosilanes (Si = SiR'₂R'').

bearing sterically bulky substituents were also smooth (Z-2o~2q). In addition, substrates having multiple furan rings (1r~1t) were readily employed for the current process to afford the corresponding multi-functionalized C-2 or C-3 symmetrical products (2r~2t). Interestingly, the newly generated multiple double bonds are all in (Z)-form, strongly suggesting that the ring-opening process is highly stereoselective.

We next turned our attention to the formation of silylated cyclopropanes based on the above mechanistic insights (Fig. 2). A brief optimization study led us to establish the one-pot reaction conditions that do not need to isolate α -silyloxy-(Z)-homoallylsilanes compounds. The key in this triple cascade hydrosilylation process was the amounts of hydrosilanes employed: while dimethylphenylsilane was most effective, the use of slightly excessive this silane (4.0 equivalents) was found to be optimal for high product yields. Under these conditions, a range of furan substrates were smoothly converted to the desired 2-alkylcyclopropyl silanes with excellent *anti*-diastereoselectivity in the presence of B(C₆F₅)₃ catalyst (Table 1; Conditions B). *Anti*-diastereoselectivity was confirmed by NMR analyses (Supplementary Figs 114–122). Furans substituted with alkyl groups at the 2-position smoothly underwent the triple hydrosilylation cascade with PhMe₂SiH (4.0 equiv.) to afford the corresponding products (*anti*-3a, 3b~3f, diastereomeric ratio;

> 99/1; the diastereoselectivity was measured by ¹H NMR of the crude reaction mixture). Likewise, 2-arylfurans with varying electronic properties were converted to *anti*-(2-arylmethyl)cyclopropyl silane in high yields (*anti*-3g~3m).

Furans bearing multi-substituted phenyl and polyaromatic moieties were also competent substrates for this hydrosilylation cascade (*anti*-3n~3p). Interestingly, multiple furans connected through a benzene core were smoothly converted to the corresponding products bearing bis- or tris-cyclopropyl groups (3q~3r, respectively), still displaying an *anti*-stereochemical relationship in each newly generated cyclopropane. It should be mentioned that Stephan, Hashmi, and coworkers recently showed that a stoichiometric reaction of B(C₆F₅)₃ with 1,6-enynes proceeds via initial cyclopropanation and then formal 1,1-carbaboration²⁸. In addition, Erker *et al.* reported that HB(C₆F₅)₂ can mediate a conversion of allyldimesitylphosphanes to phosphinomethyl-substituted cyclopropane derivatives under frustrated Lewis pair conditions in a stoichiometric manner²⁹.

Structural diversity with other furans and benzofurans. In addition to 2-substituted furans, furan and other regioisomeric derivatives were observed to undergo the selective ring-opening reactions at ambient temperature to give structurally diverse

Figure 2 | Preliminary catalytic reactions and NMR monitoring. (a) Sequential addition of hydrosilane. (b) Control of product distribution by the equivalents of hydrosilane. (c) Ring-opening and closing transformations of 2-methylfuran in one flask. (d) NMR monitoring of the $\text{B}(\text{C}_6\text{F}_5)_3$ -catalyzed cascade conversion of 2-methylfuran with hydrosilane. (**Si** = SiMe_2Ph).

products bearing $\text{sp}^3\text{C}-\text{Si}$ bonds (Table 2). Unsubstituted furan was reacted with PhMe_2SiH under the borane catalysis, leading to (*Z*)- α -hydroxy allylsilane upon treatment of the reaction mixture with methanol solution of K_2CO_3 (**Z-5**, Table 2). 3-Substituted furan was also reactive toward the silylative reduction, furnishing an allylsilane having both silyloxy and aryl moieties at the C1 and C2-positions (**Z-6**). It was interesting to see that disubstituted furan underwent the ring-opening process via slightly modified pathway (see Supplementary Discussion 2). For instance, 2,3-dimethylfuran was reacted under the same conditions to eventually provide γ -silylketone product (**7**) after treating the catalytic reaction mixture with tetrabutylammonium fluoride ($n\text{Bu}_4\text{NF}$) in one pot (**7**, 73% over two steps). In a similar manner, 2,5-dimethylfuran was doubly hydrosilylated to afford the corresponding γ -silylketone (**8**) upon treatment of $n\text{Bu}_4\text{NF}$ (**8**, 75% over two steps). The present procedure was also applicable to the silylative reduction of benzofurans. Reactions of benzofuran or 2-substituted benzofurans with hydrosilane in the presence of borane catalyst selectively proceeded to provide a range of 2-alkyl-substituted phenols (**9**~**12**) having a $\text{sp}^3\text{C}-\text{Si}$ bond in high yields again after treating the methanol solution of K_2CO_3 .

Mechanistic experiments and proposed catalytic pathway. Based on the present observations and precedents^{30,31} regarding the mechanism of $\text{B}(\text{C}_6\text{F}_5)_3$ -mediated hydrosilylation, we propose a cascade catalytic pathway leading to the *Z*-selective homoallylic silane, and subsequently the *anti*-cyclopropyl silanes, as seen in Fig. 3a using 2-methylfuran as a model substrate. On the *in situ* generation of a borane-silane adduct **I**, 2-methylfuran **1a** attacks the silylium species to afford an oxonium species **II** that immediately reacts with the borohydride leading to a partially reduced furan intermediate **III** bearing a $\text{sp}^3\text{C}-\text{Si}$ bond next to an oxygen atom. A subsequent hydrosilylation of **III** is suggested to take place through a key intermediate **IV**, at which a selective

borohydride attack at the α -carbon via a nucleophilic vinylic substitution ($\text{S}_\text{N}\text{V}$) pathway leads to α -silyloxy-(*Z*)-alkenyl silane (**V**)³² (Fig. 3a (top), **1a** to **V**).

An *O*-silyl oxonium species **VI** formed by a reaction of **V** with active species **I** is proposed to be involved in the ring-closing process, which will induce the borohydride nucleophilic attack ($\text{S}_\text{N}2'$ -type mechanism) to give the corresponding silylated cyclopropanes with the release of one equivalent of siloxane (Fig. 3a (bottom), **V** to **anti-3a**). With regard to the oxonium species **VI**, two different types of intermediates would be plausible: (i) **VI** that has C1 and C4 substituents (oxonium/silyl and methyl groups, respectively) at the opposite space and (ii) **VI'** having two groups on the same side. The relative stereochemistry (*anti*-) of two substituents in the resultant cyclopropane product can be reasoned by proposing a nucleophilic substitution of an *in situ* generated oxonium species **VI** (ref. 20) by the borohydride nucleophile to minimize the steric repulsion between substituents³³.

This mechanistic proposal was supported by a series of experiments, including kinetic and isotopic studies. A hydrosilylation of 2-methylfuran (**1a**) with PhMe_2SiD (2.0 equiv.) under standard conditions led to the exclusive incorporation of deuterium at the two positions to give **Z-2a-d₂** (Fig. 3b). A reaction of 2,3-dihydro-5-methylfuran, **13** with 1.0 equiv. of PhMe_2SiD afforded a product **Z-14-d** with a selective deuterium incorporation at the olefinic carbon (Fig. 3c). These results led us to propose a selective and consecutive attack of a borohydride nucleophile following the silylium ion (R_3Si^+) (refs 34–38) transfer and the involvement of a partially reduced dihydrofuran intermediate such as **13** during the ring-opening process. Although the proposed vinylic substitution by the borohydride nucleophile is unknown to our best knowledge, an example of an inversion at the alkenyl configuration was previously reported in a reaction of alkylvinylidonium electrophiles with halide nucleophiles³⁹.

Table 1 | $\text{B}(\text{C}_6\text{F}_5)_3$ catalysed cascade silylative transformation of furans.

Conditions A: Isolated products yields are presented, in which Z/E was determined by ^1H NMR analysis of the crude reaction mixture.

*Z-2a[†]: (3,5-dinitro)benzoate derivative from the corresponding alcohol.

†Isolated as a free hydroxyl compound on treating the reaction mixture with K_2CO_3 solution (MeOH): yields over 2 steps.

‡4.0 equiv. of silane was used.

§4.0 Mol% catalyst was used.

||6.0 Mol% catalyst and 6.0 equiv. of silane were used. (Si = SiMe_2Ph).

Conditions B: In all cases, diastereomeric ratio (dr) was >99/1 determined by ^1H NMR analysis of the crude reaction mixture. (Si = SiMe_2Ph). * Ph_2SiH_2 (3.0 equiv.) was used.

#Conducted at 40 °C.

**10.0 Mol% catalyst and 8.0 equiv. of silane were used.

††20.0 Mol% catalyst and 12.0 equiv. of silane were used.

Table 2 | B(C₆F₅)₃-catalysed silylative ring-opening of derivatives of furan and benzofuran.

 Z-5, 74%* (for Z-5)	 Z-6, 84% (for Z-6)
 7, 73%† (for 7)	 8, 75%† (for 8)
 9, 90%* (for 9)	 10, 88%* (for 10)
 11, 70%* (for 11)	 12, 79%* (for 12)

Isolated product yields are presented.

*O-desilylated derivatives were isolated upon treating the reaction mixture with K₂CO₃ in MeOH at 23 °C; yields over 2 steps.†After treatment of the reaction mixture with *n*Bu₄NF in THF at 23 °C. (Si = SiMe₂Ph, R₃ = 4-TIPSO-C₆H₄, TIPS = triisopropylsilyl). **Figure 3 | Mechanistic experiments and proposed catalytic pathway. (a)** Proposed pathway of the B(C₆F₅)₃-catalysed silylative ring-opening of furans and subsequent cyclopropanation. For simplicity, hydrosilane is shown as Si-H/D (Si = SiMe₂Ph). **(b)** Deuterium labelling experiment. **(c)** A model reaction with 2,3-dihydro-5-methylfuran. **(d)** A test ring-closing reaction of Z-2a with 1.5 equiv. of deuterated hydrosilane. **(e)** Rate-order assessment of silane based on initial rates in a range of initial silane concentrations.

As expected, the subsequent ring-closing process from **V** to the corresponding silylated cyclopropanes was highly stereoselective as proved by an isotope experiment. Indeed, when α -silyloxy-(Z)-alkenyl silane (Z-2a) was allowed to react with PhMe₂SiD (1.5 equivalents), cyclized product (*anti*-3a-d) was obtained in

82% yield as a single isomer with the exclusive incorporation of deuterium at the α -ethyl position (Fig. 3d). A stoichiometric amount of siloxane was also confirmed to be generated during the cyclopropanation process. An initial-rate study for the cyclopropanation of **V** under the B(C₆F₅)₃-catalysed hydrosilylation

Figure 4 | Enrichment and elaboration of products. (i) CsF (5 equiv.), 23 °C, 8 h, DMF, (R' = (tBu)-C₆H₄, R = Me). (ii) TBAF (3 equiv., 1 M in THF), 23 °C, 15 h, (R' = H, R = (2,4,6-triisopropyl)C₆H₂). (iii) K₂CO₃ (2 equiv.), MeOH, 0 → 23 °C, 4 h, (R' = H, R = Me). (iv) Pb(OAc)₄ (1.1 equiv.), C₆H₆, 0 → 23 °C, 2 h. (v) PPh₃ (1.3 equiv.), Diethyl azodicarboxylate (1.3 equiv.), Phthalimide (1.3 equiv.), THF, 0 → 23 °C, 20 h. (vi) VO(acac)₂ (5 mol%), TBHP (2.2 equiv.), CH₂Cl₂, -10 → 0 °C, 5 h. (vii) *p*-Br-C₆H₄CHO (1.2 equiv.), TFA (2.5 equiv.), CH₂Cl₂ (3 ml), -20 °C, 3 h; DNB = 3,5-dinitrobenzoyl. (viii) Ethyl acrylate (10 equiv.), Grubbs-II (5 mol%), 50 °C, 15 h. (ix) BF₃•Et₂O (1.1 equiv.), CH₂Cl₂ (0.14 M), -78 °C, 5 h, (R' = H, R = Me). (x) (COCl)₂ (2.0 equiv.), DMSO (4.0 equiv.), Et₃N (5.0 equiv.), -78 °C, 4 h, (R' = H, R = Me). (xi) B(C₆F₅)₃ (3.0 mol%), PhMe₂SiH (1.5 equiv.), CH₂Cl₂, 0 → 23 °C, 12 h, (R' = H, R = Me). (xii) 2-Methylfuran (**1a**, 1.0 equiv.), B(C₆F₅)₃ (5.0 mol%), Ph₂SiH₂ (3.0 equiv.), CH₂Cl₂, 0 → 23 °C, 8 h. (xiii) (a) **1ac** (1.0 equiv.), B(C₆F₅)₃ (5.0 mol%), Ph₂SiH₂ (3.0 equiv.), CH₂Cl₂, 0 → 23 °C, 10 h; (b) H₂O₂ (20 equiv.), KF (10 equiv.), KHCO₃ (10 equiv.), THF/MeOH (1/1), 23 °C, 16 h. (**Si** = SiPh₂H/SiMe₂Ph).

conditions clearly showed that the reaction was first-order in silane concentration (Fig. 3e; see Supplementary Figs 4 and 5), leading us to propose that the cyclization path involves an O-silylated oxonium ion possessing a borohydride anion⁴⁰.

Synthetic applications. The synthetic utility of the products obtained was demonstrated to be very broad, including a wide range of simple and convenient organic transformations (Fig. 4). α -Silyloxy-(*Z*)-alkenyl silanes were easily desilylated to give allylic- or homoallylic alcohols with retention of double bond stereochemistry (**Z-15** and **Z-16**, respectively). α -Hydroxy homoallylic silane (**Z-2a'**), accessed through the O-desilylation of α -silyloxy-(*Z*)-homoallylic silane, turned out to be also synthetically versatile. α -Silyloxy acetate (**Z-17**) was readily produced by a radical Brook rearrangement in good yield⁴¹. Moreover, **Z-2a'** was smoothly converted to (*Z*)- α -amino silane derivative (**Z-18**) under the Mitsunobu conditions leading to C–N bond formation. Epoxide (**19**) was obtained *syn*-selectively via the Sharpless directed epoxidation of **Z-2a'** (ref. 42), thus proving that α -silyl alcohol works as an effective directing group. An efficient Prins cyclization⁴³ of **Z-2a'** was achieved to furnish a tetrasubstituted pyran ring (*syn*-**20**) with complete stereocontrol. The observed *syn*-stereochemistry was unambiguously confirmed by NMR and X-ray diffraction analyses. Cross metathesis of **Z-2a'** with ethyl acrylate was carried out stereoselectively with Grubbs II catalyst to give **E-21**. Lewis acid-promoted allylic rearrangement⁴⁴ of α -silyloxy-(*Z*)-allylic silane bearing *tert*-butyldimethylsilyloxy group **Z-5''** (*tert*-butyldimethylsilyl derivative of **Z-5**, see Supplementary Methods) proceeded smoothly to afford silyloxy vinylic silane (**E-22**), another versatile synthetic building block. Acylsilane (**E-23**), a valuable organosilicon reagents in organic synthesis⁴⁵, was readily obtained.

Two preparative procedures of cyclopropyl silanes were mild and efficient on a large scale (Supplementary Methods): (i) a stepwise route via α -silyloxy homoallylic silane (**Z-2a**, procedure “xi” in 77%), and (ii) a direct conversion of furan (procedure “xii” in 81% yield). The subsequent oxidation of the C–Si bond under the Tamao–Fleming conditions⁴⁶ proceeded with retention of stereochemistry to furnish the versatile synthons, cyclopropanol (**24**) and its methanesulfonyl derivative (**25**), which are broadly utilized in synthetic and medicinal chemistry⁴⁷. For instance, cascarillic acid⁴⁸, a natural product derived from the bark of the medicinal shrub *Croton eluteria*, can be envisioned to be accessible based on our current procedure.

In summary, we have developed the tris(pentafluorophenyl)-borane-catalysed cascade transformation of furans that are readily available from renewable biomass resources to obtain synthetically valuable silicon-functionalized products such as α -silyloxy-(*Z*)-alkenyl silanes and *anti*-cyclopropyl silanes. Simply by varying the stoichiometry of employed hydrosilanes in presence of B(C₆F₅)₃ catalyst, the product distribution could completely be controlled with high efficiency (TON up to 2,000) and excellent stereoselectivity. The transformation does not require transition metal catalysts, proceeds efficiently on large scale, and is broadly applicable to various types of furans and their derivatives, bringing about a significant structural diversity in the product obtained. A proposed mechanistic pathway involves a series of hydrosilylation cascades, containing a ring-opening and subsequent S_N2'-type ring-closing process, both mediated by a B(C₆F₅)₃ catalyst. The synthetic utility of obtained silicon-functionalized products was demonstrated in a range of post-transformations. It is anticipated that this study will stimulate future developments in the transformations of biomass-derived platform chemicals to synthetic valuables.

Methods

General procedure for the silylative ring opening reaction (conditions A, GP1). In a flame-dried flask bearing a stirring bar, $\text{B}(\text{C}_6\text{F}_5)_3$ (0.01 ~ 0.02 mmol, 2.0 mol%) was dissolved in CH_2Cl_2 (0.4 ~ 0.8 ml). Silane (1.025 ~ 2.050 mmol) was added, and the solution was shaken shortly to make it homogeneous. The corresponding furan derivative (**1a** ~ **1t**, 0.50 ~ 1.0 mmol) was then added and the reaction mixture was stirred at 23 °C for the indicated time (1 ~ 15 h). After quenching the reaction mixture with Et_3N (10.0 ~ 20.0 mol%), the crude reaction mixture was concentrated under reduced pressure and then purified by flash column chromatography on silica gel (using either hexane only or a mixture of hexane/ethyl acetate) to afford the desired products (**Z-2a** ~ **Z-2t**, in all cases *Z/E* > 99/1).

General procedure for the cyclopropanation reaction (conditions B, GP2). In a flame-dried flask bearing a stirring bar, $\text{B}(\text{C}_6\text{F}_5)_3$ (0.025 mmol, 5.0 mol%) was dissolved in CH_2Cl_2 (0.2 ml). Silane (1.5 ~ 2.0 mmol) was added, and the solution was shaken shortly to make it homogeneous. The corresponding furan derivative (**1a** ~ **1t**, 0.50 mmol) was then added at 0 °C and the reaction mixture was stirred at 23 °C for the indicated time (4 ~ 22 h). After quenching the reaction mixture with Et_3N (10.0 ~ 20.0 mol%), the crude reaction mixture was concentrated under reduced pressure and then purified by flash column chromatography on silica gel (using either hexane or a mixture of hexane and ethyl acetate) to afford the desired products (*anti*, **3a** ~ **3r**, in all cases *dr* > 99/1).

Data availability. The authors declare that the data supporting of the findings of this study are available within the article and Supplementary Information files. For the experimental procedures and spectroscopic and physical data of compounds, see Supplementary Methods. For NMR analysis of the compounds in this article, see Supplementary Figs 6–187. The CCDC 1505482 (**Z-2a''**) and CCDC 1505484 (*syn-20'*) contain the supplementary crystallographic data for this paper (Supplementary Tables 4 and 5). These data can be obtained free of charge from The Cambridge Crystallographic Data Centre via http://www.ccdc.cam.ac.uk/data_request/cif. All other data are available from the authors on reasonable request.

References

- Corma, A., Iborra, S. & Vely, A. Chemical routes for the transformation of biomass into chemicals. *Chem. Rev.* **107**, 2411–2502 (2007).
- De Souza, R. O. M. A., Miranda, L. S. M. & Luque, R. Bio(chemo)technological strategies for biomass conversion into bioethanol and key carboxylic acids. *Green Chem.* **16**, 2386–2405 (2014).
- Hu, L. *et al.* Catalytic conversion of biomass-derived carbohydrates into fuels and chemicals via furanic aldehydes. *RSC Adv.* **2**, 11184–11206 (2012).
- Mirzaei, H. M. & Karimi, B. Sulphanilic acid as a recyclable bifunctional organocatalyst in the selective conversion of lignocellulosic biomass to 5-HMF. *Green Chem.* **18**, 2282–2286 (2016).
- Van Putten, R.-J. *et al.* Hydroxymethylfurfural, a versatile platform chemical made from renewable resources. *Chem. Rev.* **113**, 1499–1597 (2013).
- Luterbacher, J. S., Alonso, D. M. & Dumesic, J. A. Targeted chemical upgrading of lignocellulosic biomass to platform molecules. *Green Chem.* **16**, 4816–4838 (2014).
- Ruppert, A. M., Weinberg, K. & Palkovits, R. Hydrogenolysis goes bio: from carbohydrates and sugar alcohols to platform chemicals. *Angew. Chem. Int. Ed.* **51**, 2564–2601 (2012).
- Gallezot, P. Conversion of biomass to selected chemical products. *Chem. Soc. Rev.* **41**, 1538–1558 (2012).
- Cheng, Y.-T., Wang, Z., Gilbert, C. J., Fan, W. & Huber, G. W. Production of *p*-xylene from biomass by catalytic fast pyrolysis using ZSM-5 catalysts with reduced pore openings. *Angew. Chem. Int. Ed.* **51**, 11097–11100 (2012).
- Julis, J. & Leitner, W. Synthesis of 1-octanol and 1,1-dioctyl ether from biomass-derived platform chemicals. *Angew. Chem. Int. Ed.* **51**, 8615–8619 (2012).
- Xia, Q.-N. *et al.* $\text{Pd}(\text{NbOPO}_4)_4$ multifunctional catalyst for the direct production of liquid alkanes from aldol adducts of furans. *Angew. Chem. Int. Ed.* **53**, 9755–9760 (2014).
- Clement, M. J., Corma, A. & Iborra, S. Conversion of biomass platform molecules into fuel additives and liquid hydrocarbon fuels. *Green Chem.* **16**, 516–547 (2014).
- Thananathanachon, T. & Rauchfuss, T. B. Efficient production of the liquid fuel 2,5-dimethylfuran from fructose using formic acid as a reagent. *Angew. Chem. Int. Ed.* **49**, 6616–6618 (2010).
- Cha, H. G. & Choi, K.-S. Combined biomass valorization and hydrogen production in a photochemical cell. *Nat. Chem.* **7**, 328–333 (2015).
- Zeitsch, K. J. *The Chemistry and Technology of Furfural and its Many By-Products* 1st ed., Vol. 13 (Elsevier, 2000).
- Ghorpade, V. M. & Hanna, M. A. Method and apparatus for production of levulinic acid via reactive extrusion. US patent 5,859,263, 1999.
- Timokhin, B. V., Baransky, V. A. & Eliseeva, G. D. Levulinic acid in organic synthesis. *Russ. Chem. Rev.* **68**, 73–84 (1999).
- Sutton, A. D. *et al.* The hydrodeoxygenation of bioderived furans into alkanes. *Nat. Chem.* **5**, 428–432 (2013).
- Mack, D. J., Guo, B. & Njardarson, J. T. Synthesis of allylic and homoallylic alcohols from unsaturated cyclic ethers using a mild and selective C–O reduction approach. *Chem. Commun.* **48**, 7844–7846 (2012).
- Adduci, L. L., Bender, T. A., Dabrowski, J. A. & Gagné, M. R. Chemoselective conversion of biologically sourced polyols into chiral synthons. *Nat. Chem.* **7**, 576–581 (2015).
- McLaughlin, M. P., Adduci, L. L., Becker, J. J. & Gagné, M. R. Iridium-catalyzed hydrosilylative reduction of glucose to hexane(s). *J. Am. Chem. Soc.* **135**, 1225–1227 (2013).
- Adduci, L. L., McLaughlin, M. P., Bender, T. A., Becker, J. J. & Gagné, M. R. Metal-free deoxygenation of carbohydrates. *Angew. Chem. Int. Ed.* **53**, 1646–1649 (2014).
- Scott, D. J., Fuchter, M. J. & Ashley, A. E. Metal-free hydrogenation catalyzed by an air-stable borane: use of solvent as a frustrated Lewis base. *Angew. Chem. Int. Ed.* **53**, 10218–10222 (2014).
- Rubin, M., Schwieter, T. & Gevorgyan, V. Highly efficient $\text{B}(\text{C}_6\text{F}_5)_3$ -catalyzed hydrosilylation of olefins. *J. Org. Chem.* **67**, 1936–1940 (2002).
- Gandhamsetty, N., Joung, S., Park, S.-W., Park, S. & Chang, S. Boron-catalyzed silylative reduction of quinolines: selective sp^3 C–Si bond formation. *J. Am. Chem. Soc.* **136**, 16780–16783 (2014).
- Gandhamsetty, N., Park, S. & Chang, S. Selective silylative reduction of pyridines leading to structurally diverse azacyclic compounds with the formation of sp^3 C–Si bonds. *J. Am. Chem. Soc.* **137**, 15176–15184 (2015).
- Kim, Y. & Chang, S. Borane-catalyzed reductive α -silylation of conjugated esters and amides leaving carbonyl groups intact. *Angew. Chem. Int. Ed.* **55**, 218–222 (2016).
- Hansmann, M. M. *et al.* Cyclopropanation/carboboration reactions of enynes with $\text{B}(\text{C}_6\text{F}_5)_3$. *J. Am. Chem. Soc.* **137**, 15469–15477 (2015).
- Krupski, S., Kehr, G., Daniliuc, C. G. & Erker, G. Cyclopropane formation under frustrated Lewis pair conditions. *Chem. Commun.* **52**, 2695–2697 (2016).
- Parks, D. J. & Piers, W. E. Tris(pentafluorophenyl)boron-catalyzed hydrosilylation of aromatic aldehydes, ketones, and esters. *J. Am. Chem. Soc.* **118**, 9440–9441 (1996).
- Gevorgyan, V., Liu, J.-X., Rubin, M., Benson, S. & Yamamoto, Y. A novel reduction of alcohols and ethers with a HSiEt_3 /catalytic $\text{B}(\text{C}_6\text{F}_5)_3$ system. *Tetrahedron Lett.* **40**, 8919–8922 (1999).
- Bernasconi, C. F. & Rappoport, Z. Recent advances in our mechanistic understanding of $\text{S}_\text{N}2$ reactions. *Acc. Chem. Res.* **42**, 993–1003 (2009).
- Taylor, R. E., Risatti, C. A., Engelhardt, F. C. & Schmitt, M. J. Cyclopropane structural units from homoaldol adducts. *Org. Lett.* **5**, 1377–1379 (2003).
- Parks, D. J., Blackwell, J. M. & Piers, W. E. Studies on the mechanism of $\text{B}(\text{C}_6\text{F}_5)_3$ -catalyzed hydrosilylation of carbonyl functions. *J. Org. Chem.* **65**, 3090–3098 (2000).
- Houghton, A. Y., Hurmalainen, J., Mansikkamäki, A., Piers, W. E. & Tuononen, H. M. Direct observation of a borane–silane complex involved in frustrated Lewis-pair-mediated hydrosilylations. *Nat. Chem.* **6**, 983–988 (2014).
- Oestreich, M., Hermeke, J. & Mohr, J. A unified survey of Si–H and H–H bond activation catalyzed by electron-deficient boranes. *Chem. Soc. Rev.* **44**, 2202–2220 (2015).
- Rendler, S. & Oestreich, M. Conclusive evidence for an $\text{S}_\text{N}2$ -Si mechanism in the $\text{B}(\text{C}_6\text{F}_5)_3$ -catalyzed hydrosilylation of carbonyl compounds: Implications for the related hydrogenation. *Angew. Chem. Int. Ed.* **47**, 5997–6000 (2008).
- Sakata, K. & Fujimoto, H. Quantum chemical study of $\text{B}(\text{C}_6\text{F}_5)_3$ catalyzed hydrosilylation of carbonyl group. *J. Org. Chem.* **78**, 12505–12512 (2013).
- Ochiai, M., Oshima, K. & Masaki, Y. Inversion of configuration in nucleophilic vinylic substitutions of (*E*)- β -alkylvinylidonium tetrafluoroborates with halides. *J. Am. Chem. Soc.* **113**, 7059–7061 (1991).
- Kalkofen, R., Brandau, S., Wibbeling, B. & Hoppe, D. Synthesis of stereohomogeneous cyclopropanecarbaldehydes and cyclopropyl ketones by cycloalkylation of 4-hydroxy-1-alkenyl carbamates. *Angew. Chem. Int. Ed.* **43**, 6667–6669 (2004).
- Brook, A. G. Some molecular rearrangements of organosilicon compounds. *Acc. Chem. Res.* **7**, 77–84 (1974).
- Zhang, W., Basak, A., Kosugi, Y., Hoshino, Y. & Yamamoto, H. Enantioselective epoxidation of allylic alcohols by a chiral complex of vanadium: an effective controller system and a rational mechanistic model. *Angew. Chem. Int. Ed.* **44**, 4389–4391 (2005).
- Zheng, K. *et al.* Completely OH-selective FeCl_3 -catalyzed Prins cyclization: highly stereoselective synthesis of 4-OH-tetrahydropyrans. *J. Am. Chem. Soc.* **134**, 17564–17573 (2012).

44. Kim, A. I., Kimmel, K. L., Romero, A., Smitrovich, J. H. & Woerpel, K. A. Mechanistic studies of the allylic rearrangements of α -silyloxy allylic silanes to silyloxy vinylic silanes. *J. Org. Chem.* **72**, 6595–6598 (2007).
45. Zhang, H.-J., Priebbenow, D. L. & Bolm, C. Acylsilanes: valuable organosilicon reagents in organic synthesis. *Chem. Soc. Rev.* **42**, 8540–8571 (2013).
46. Fleming, I., Henning, R., Parker, D. C., Plaut, H. E. & Sanderson, P. E. J. The phenyldimethylsilyl group as a masked hydroxy group. *J. Chem. Soc. Perkin Trans. 1*, 317–337 (1995).
47. Kulinkovich, O. G. The chemistry of cyclopropanols. *Chem. Rev.* **103**, 2597–2632 (2003).
48. Kumar, P., Dubey, A. & Harbindu, A. Enantio- and diastereocontrolled conversion of chiral epoxides to *trans*-cyclopropane carboxylates: application to the synthesis of cascarillic acid, grenadamide and L-(–)-CCG-II. *Org. Biomol. Chem.* **10**, 6987–6994 (2012).

Acknowledgements

This article is dedicated to Professor Naoto Chatani on the occasion of his 60th birthday. This research was supported by the Institute for Basic Science (IBS-R10-D1) in Korea. We are thankful to Prof. Mu-Hyun Baik for his inspiring discussions and critical reading.

Author contributions

C.K.H., N.G., and S.P. carried out the experiments. C.K.H., N.G. and S.C. designed and directed this project. S.P. and S.C. wrote the manuscript. All authors analysed the data, discussed the results and commented on the manuscript.

Additional information

Supplementary Information accompanies this paper at <http://www.nature.com/naturecommunications>

Competing financial interests: The authors declare no competing financial interests.

Reprints and permission information is available online at <http://npg.nature.com/reprintsandpermissions/>

How to cite this article: Hazra, C. K. *et al.* Borane catalysed ring opening and closing cascades of furans leading to silicon functionalized synthetic intermediates. *Nat. Commun.* **7**, 13431 doi: 10.1038/ncomms13431 (2016).

Publisher's note: Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

This work is licensed under a Creative Commons Attribution 4.0 International License. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in the credit line; if the material is not included under the Creative Commons license, users will need to obtain permission from the license holder to reproduce the material. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>

© The Author(s) 2016