ARTICLE

Received 14 Mar 2011 | Accepted 22 Aug 2011 | Published 20 Sep 2011

DOI: 10.1038/ncomms1488

An energy transduction mechanism used in bacterial flagellar type III protein export

Tohru Minamino^{1,2}, Yusuke V. Morimoto¹, Noritaka Hara¹ & Keiichi Namba¹

Flagellar proteins of bacteria are exported by a specific export apparatus. FliI ATPase forms a complex with FliH and FliJ and escorts export substrates from the cytoplasm to the export gate complex, which is made up of six membrane proteins. The export gate complex utilizes proton motive force across the cytoplasmic membrane for protein translocation, but the mechanism remains unknown. Here we show that the export gate complex by itself is a proton-protein antiporter that uses the two components of proton motive force, $\Delta\psi$ and ΔpH , for different steps of the protein export process. However, in the presence of FliH, FliI and FliJ, a specific binding of FliJ with an export gate membrane protein, FlhA, is brought about by the FliH-FliI complex, which turns the export gate into a highly efficient, $\Delta\psi$ -driven protein export apparatus.

¹ Graduate School of Frontier Bioscience, Osaka University, Suita, Osaka, Japan. ² Precursory Research for Embryonic Science and Technology, Japan Science and Technology Agency, 4-1-8 Honcho, Kawaguchi, Saitama 332-0012, Japan. Correspondence and requests for materials should be addressed to T.M. (email: tohru@fbs.osaka-u.ac.jp) or to K.N. (email: keiichi@fbs.osaka-u.ac.jp).

Proton motive force (PMF), the electrochemical potential difference of protons across a biological membrane, is utilized as the energy source of many biological activities, including protein export, ATP synthesis and flagellar motor rotation. PMF consists of two components: the electric potential difference ($\Delta\psi$) and the proton concentration difference (ΔpH). Although $\Delta\psi$ and ΔpH are equivalent driving forces for proton movement and translocation in physics and have also been believed to have equivalent roles in biological functions, recent experimental data suggest a differential usage of $\Delta\psi$ and ΔpH in biological processes, such as flagellar motor rotation¹ and ATP synthesis².

The bacterial flagellum, which is responsible for motility, consists of at least three substructures: the basal body, the hook and the filament. Flagellar assembly begins with the basal body, followed by the hook and finally the filament. For construction of the flagellum, many of the flagellar component proteins are transported to the distal end of the flagellar structure by the flagellar type III protein export apparatus. The export apparatus consists of three soluble proteins, FliH, FliI and FliJ, and six integral membrane proteins, FlhA, FlhB, FliO, FliP, FliQ and FliR^{3–5}. These components of the export apparatus share substantial sequence and functional similarities with those of the type III secretion system of pathogenic bacteria, which directly inject virulence factors into their host cells⁶. Recently, it has become apparent that FliH, FliI and FliJ are evolutionarily related to components of F_0F_1 -ATP synthase, which utilizes PMF for ATP synthesis^{5,7–9}.

FliI is an ATPase¹⁰ and forms a complex with FliH and FliJ^{11–13}. The rate of the protein export is markedly low in the absence of FliH, FliI and FliJ, although these components are not essential because flagellar formation occasionally occurs in their absence^{13–16}. The FliH–FliI–FliJ complex binds to export substrates and chaperon–substrate complexes in the cytoplasm^{17,18} and delivers them to the docking platform of the export gate made up of the cytoplasmic domains of FlhA and FlhB (FlhA_C and FlhB_C, respectively)^{19,20}. An association of the substrate-bound FliH_X–FliI₆–FliJ ring complex with the FlhA_C–FlhB_C docking platform induces the initial entry of the substrate into the gate^{15,21}. The chemical energy derived from ATP hydrolysis by FliI appears to be required for the dissociation of the FliH_X–FliI₆–FliJ ring complex from the gate²². The export gate complex utilizes PMF as an energy source for the unfolding and translocation of export substrates into the central channel of the flagellum^{15,16}. Interestingly, it has been shown, albeit qualitatively, that $\Delta\psi$ alone is sufficient to support export in wild-type cells¹⁶. Thus, it remains unclear how PMF is used as the energy source.

To clarify the roles of the two components of PMF in flagellar protein export, we collapsed $\Delta\psi$ and ΔpH separately and analysed flagellar protein export in a wild-type strain of *Salmonella enterica* and its ΔfliH – fliI bypass mutant whose second-site P28T mutation in FlhB recovers the export efficiency of some substrates to wild-type levels; therefore, flagellar formation increases markedly in the absence of FliH and FliI¹⁵. We show that the export gate uses the two components of PMF differently for protein export depending on the presence and absence of FliH and FliI. Protein–protein interactions monitored by a pull-down assay using GST affinity chromatography combined with bacterial genetics show that what is required for efficient $\Delta\psi$ -driven protein export of the wild-type export apparatus is not FliH and FliI themselves, but instead a specific interaction of FlhA with FliJ that is brought about by the docking of the FliH–FliI₆ ring complex to the gate.

Results

Effects of $\Delta\psi$ and ΔpH on flagellar protein export. To investigate the role of each component of PMF on flagellar protein export in wild-type cells and a ΔfliH – fliI bypass mutant [ΔfliH – fliI flhB (P28T)], $\Delta\psi$ and ΔpH were collapsed individually by treatment with valinomycin in the presence of KCl¹⁶ and by adding potassium

benzoate²³, respectively. The $\Delta\psi$ and ΔpH were measured using tetramethylrhodamine methyl ester (TMRM)¹ and a ratiometric pH indicator protein, pHluorin²⁴, respectively. Because the hook-capping protein FlgD is one of the export substrates that are secreted at wild-type levels by the ΔfliH – fliI bypass mutant¹⁵, we analysed the secretion level of FlgD as a representative measure of the export activity (Fig. 1). The $\Delta\psi$ and ΔpH measured at different pH values in the presence and absence of benzoate are shown in Figure 2. In wild-type cells, at an external pH of 7.0, the secretion level was significantly reduced by discharging $\Delta\psi$ by 20 μM valinomycin at 150 mM KCl (Fig. 1a left panel) but was not reduced at all by collapsing ΔpH by using potassium benzoate (Fig. 1b left panel). These results confirmed a previous report¹⁶ in a more quantitative manner. In contrast, the secretion by the ΔfliH – fliI bypass mutant was effectively abolished not only by discharging $\Delta\psi$ (Fig. 1a right panel) but also by collapsing ΔpH (Fig. 1b right panel). Interestingly, neither of the remaining ΔpH (0.35, which is equivalent to 24 mV) nor $\Delta\psi$ (81 mV) was effective in driving protein export. The intracellular levels of FlgD were not changed by either treatment. Because the total PMF of the fliH – fliI bypass mutant was the same as that of wild-type cells (Fig. 2d), these results indicate that $\Delta\psi$ and ΔpH are both essential and are used differently for flagellar protein export in the absence of FliH and FliI.

The acidification of the cytoplasm is known to interfere with flagellar protein export¹⁶. This relationship raised the possibility that the export apparatus of the bypass mutant may be more sensitive to low intracellular pH than the apparatus of the wild-type cells. To examine the ΔpH dependence of protein export without lowering the intracellular pH, we varied only the external pH over a range of 6.0–7.5 (Fig. 1c). The intracellular pH did not change significantly over this external pH range (Fig. 2b). In wild-type cells, the secretion level was constant over this external pH range (Fig. 1c left panel). However, the secretion level by the ΔfliH – fliI bypass mutant showed a marked dependence on the external pH and therefore on ΔpH ; it increased on a downward pH shift from 7.0 to 6.0 and almost diminished by an upward shift to 7.5 where ΔpH is nearly zero (Fig. 1c right panel and Fig. 2c). The $\Delta\psi$ did not change significantly over an external pH range of 6.0–7.5 (Fig. 2a). Because the ΔpH component of the PMF was significantly increased by lowering the external pH (Fig. 2c), resulting in an increase in total PMF (Fig. 2d), we conclude that an increase in the ΔpH component facilitates flagellar protein export in the ΔfliH – fliI bypass mutant but not in wild-type cells. We obtained essentially the same results with another fliH – fliI bypass mutant (ΔfliH – fliI flhA (V404M)) (Supplementary Fig. S1).

Solvent-isotope effect on flagellar protein export. We next tested whether the rate of proton transfer would affect the rate of protein export. It has been shown that D_2O affects the high-speed rotation of the proton-driven bacterial flagellar motor at a low load because of a limitation in the rate of proton translocation through the proton channel of the stator²⁵. We therefore analysed this solvent-isotope effect on the flagellar protein export. The PMF did not show any changes from replacing H_2O with D_2O (data not shown). Although the FlgD-secretion level of the wild-type cells was not affected by D_2O , the secretion level of the ΔfliH – fliI bypass mutant was markedly reduced (Fig. 3). Interestingly, an increase in ΔpD by lowering the external pD enhanced the secretion rate of FlgD in the bypass mutant in a manner similar to that of the ΔpH (Fig. 3). This result suggests that the protein export activity is limited by the rate of proton translocation through the export gate in the absence of FliH and FliI.

Effect of the FlhB(P28T) mutation alone on ΔpH dependence. It is thought that the flhB (P28T) mutation in the absence of FliH and FliI increases only the probability of the substrate insertion into the gate¹⁵. To examine the effect of this mutation by itself, we analysed the ΔpH dependence of the protein export by the flhB (P28T)

Figure 1 | Effect of $\Delta\Psi$ and ΔpH on flagellar protein export. (a) Effect of valinomycin in the presence of 150 mM KCl. Immunoblotting, using polyclonal anti-FlgD antibody, of whole-cell proteins and culture supernatant fractions prepared from SJW1103 (wild-type, indicated as WT) and MMH10117 ($\Delta fliH$ -*fliB*(P28T), indicated as $\Delta fliH$ *fliB**) grown at 30°C in a buffer containing 150 mM KCl with or without 20 μM valinomycin at an external pH of 7.0. (b) Effect of potassium benzoate on the secretion level of FlgD (upper panel) and ΔpH change (closed circle) and the quantified secretion level of FlgD (open circle; lower panel) at an external pH of 7.0. Intracellular pH was measured with pHluorin. These data are the average of at least three independent measurements. The experimental errors are less than 10%. (c) Effect of external pH on FlgD secretion over a pH range of 6.0–7.5.

mutant in the presence of FliH and FliI at an external pH of 7.0. The result was essentially the same as that of the wild-type cells; there was no ΔpH dependence (Supplementary Fig. S2). This result confirms that only $\Delta\Psi$ is used for protein export in the presence of FliH and FliI and that, when FliH and FliI are missing, the export gate requires both $\Delta\Psi$ and ΔpH to couple the energy of the proton influx with protein export.

Effect of FliJ deletion on PMF-driven protein export. FliJ is required for an efficient export process, although it is not essential for export, because occasional flagellar formation is observed in its absence¹⁴. FliJ enhances the interaction of the FliT–FliD chaperone-substrate complex with the N-terminal portion of FlhA_C²⁶. Because FlhA is a component of the export gate^{3–5}, there is a possibility that FliJ may have a role in the energy transduction mechanism of the flagellar protein export. We therefore analysed the effect of FliJ deletion in the $\Delta fliH$ -*fliI* bypass mutant at different external pH values. When FliJ was deleted from the $\Delta fliH$ -*fliI* bypass mutant, the secretion level of FlgD was largely reduced, although not to the level of the $\Delta fliH$ -*fliI*-*fliJ* triple null mutant (Fig. 4a). This observation indicates that the FlhB(P28T) mutation can bypass the FliJ defect to some extent. The deletion of FliJ did not change the response of the *fliH*-

fliI bypass mutant to the change in ΔpH ; a downward shift of external pH from 7.0 to 6.0 caused an increase in the secretion level both in the presence and in the absence of FliJ (Fig. 4b). Interestingly, co-overproduction of FliH and FliI in the $\Delta fliH$ -*fliI*-*fliJ* *fliB*(P28T) mutant neither bypassed the motility defect (Fig. 4c) nor recovered the protein export activity (Fig. 4d). The ΔpH dependence of the protein export remained even in the presence of FliH and FliI (Fig. 4d). These results clearly indicate that FliJ is the essential component responsible for the $\Delta\Psi$ -driven protein export by the wild-type export apparatus.

Interaction of FlhA with the soluble proteins. FlhA is composed of an amino-terminal transmembrane domain (FlhA_{TM}) and a carboxy-terminal cytoplasmic domain (FlhA_C). FlhA_C not only forms part of the docking platform for the soluble export components FliI, FliH and FliJ but also is directly involved in the early process of protein export with these soluble components^{11,21}. An in-frame deletion of residues 328–351 of FlhA, which is part of the flexible linker between FlhA_{TM} and FlhA_C²⁷, results in a loss-of-function phenotype²⁸. It has been shown that FliJ binds to the flexible linker²⁶, raising the possibility that the loss-of-function phenotype of the FlhA(Δ328–351) mutant could result from its impaired

Figure 2 | pH dependence of PMF components. pH dependence of (a) $\Delta\psi$, (b) intracellular pH, (c) ΔpH and (d) total PMF. $\Delta\psi$ was measured in 10 mM potassium phosphate, 0.1 mM EDTA, and 10 mM sodium lactate at an external pH of 6.0, 6.5, 7.0 or 7.5 with (closed bar) or without (open bar) 20 mM potassium benzoate using TMRM. More than 100 cells were measured. Intracellular pH was measured with pHluorin. At least six independent experiments were conducted. Vertical bars indicate standard deviations.

ability to interact with FliJ. However, it is not yet known whether the linker of FlhA is also involved in the interaction with FliH and FliI. To test for this possibility, we performed pull-down assays by GST affinity chromatography. We used GST as a negative control. His-FLAG-FlhA copurified with GST-FliH, GST-FliI and GST-FliJ, but not with GST alone (Fig. 5a left panels). His-FLAG-FlhA bound to FliJ with an affinity much higher than its affinity to FliH and FliI. The deletion of residues 328–351 abolished the interaction with FliJ but not with FliH or FliI (Fig. 5a right panels). These results indicate that residues 328–351 of FlhA are the binding target of FliJ but are not the binding target of FliH and FliI.

The P28T mutation of FlhB overcomes the FliH–FliI–FliJ defects to some extent (Fig. 4a). To investigate whether the P28T mutation also suppresses the FlhA($\Delta 328$ –351) defect, we transformed a

Figure 3 | Solvent-isotope effects on flagellar protein export. Levels of FlgD secreted by the wild-type (WT) and $\Delta fliH fliB(P28T)$ mutant ($\Delta fliH fliB^*$) cells grown in a buffer containing either H₂O or D₂O at external pH levels of 6.0 and 7.0. The level of FlgD secretion was analysed by immunoblotting with the polyclonal anti-FlgD antibody.

$\Delta fliH fliB(P28T) \Delta fliA$ mutant strain with pTrc99A, pMM108 (His-FLAG-FlhA) or pMM108-1 (His-FLAG-FlhA($\Delta 328$ –351)) and analysed the motility of the resulting transformants in soft agar plates (Fig. 5b). His-FLAG-FlhA restored the motility of the $\Delta fliH fliB(P28T) \Delta fliA$ mutant, whereas His-FLAG-FlhA($\Delta 328$ –351) did not. This result indicates that residues 328–351 of FlhA are critical for the FlhA function even in the $\Delta fliH fliB$ bypass mutant. Together with the binding of FliJ to residues 328–351 of FlhA (Fig. 5a), these results suggest that a specific conformation of the FlhA linker must be brought about by the proper binding of FliJ to residues 328–351 of FlhA to turn the export gate into an efficient, $\Delta\psi$ -driven protein export apparatus.

Characterization of a *fliJ*($\Delta 13$ –24) mutant. It has been shown that in-frame deletion of residues 13–24 in FliJ considerably reduced the export activity¹⁴. This raised the possibility that the deletion may affect the interaction with the linker region of FlhA. To test this possibility, we analysed the interaction between FlhA and FliJ($\Delta 13$ –24) by pull-down assays. His-FLAG-FlhA did not copurify with GST-FliJ($\Delta 13$ –24) (Fig. 6a). Unlike wild-type FliJ, FliJ($\Delta 13$ –24) did not restore motility of the $\Delta fliH fliB(P28T) \Delta fliA$ mutant (Fig. 6b). These results indicate that the reduced export activity of the *fliJ*($\Delta 13$ –24) mutant results from the impaired ability of FliJ to interact with the linker of FlhA.

To understand the role of the specific interaction between FliJ and the flexible linker of FlhA in more detail, nine gain-of-function mutants were isolated from the *fliJ*($\Delta 13$ –24) mutant. Genetic mapping and DNA sequencing revealed that the second-site mutations are located in the *fliH* and *fliI* genes (Supplementary Fig. S3). These pseudorevertants significantly improved motility although not as well as that of the wild type (Fig. 6c). Most of the flagellar proteins were detected in the culture supernatant of the pseudorevertants (Fig. 6d). In one of the pseudorevertants, MM104H-3 (*fliJ*($\Delta 13$ –24) *fliH*($\Delta 96$ –97)), flagellar proteins were secreted at effectively wild-type levels (Fig. 6d lanes 5 and 7). Consistently, the flagellar hook basal bodies, often with filaments attached, were observed by electron microscopy (data not shown).

To test whether these second-site FliH and FliI mutations also suppress the *fliJ*-null mutant, we analysed the motility of the $\Delta fliH fliI fliJ$ triple null mutant transformed with pMMH1001 (FliH + FliI), pMMH104-3 (FliH($\Delta 96$ –97) + FliI) or pMMH104-6 (FliH + FliI(L244R)). These plasmids did not restore the motility of the $\Delta fliH fliI fliJ$ triple null mutant at all (Supplementary Fig. S4a), indicating that the second-site mutations do not suppress the *fliJ*-null mutation.

To test the effect of each second-site mutation by itself on the motility, we isolated the second-site *fliH* and *fliI* mutants and analysed their motility. The motility was normal (Supplementary Fig. S4b). As these second-site mutations show no phenotype, they do not appear to affect the function of the FliH and FliI proteins.

Figure 4 | Effects of FliJ deletion on flagellar protein export.

(a) Immunoblotting, using polyclonal anti-FlgD, of whole-cell proteins and culture supernatant fractions prepared from the wild type (WT), a $\Delta fliH$ - $fliI$ double null mutant ($\Delta fliH$), a $fliH$ - $fliI$ bypass mutant ($\Delta fliH fliB^*$), a $fliH$ - $fliI$ - $fliJ$ triple null mutant ($\Delta fliH fliJ$) and a $fliH$ - $fliI$ - $fliJ fliB(P28T)$ ($\Delta fliH fliJ fliB^*$). (b) Effect of external pH on FlgD secretion by the $\Delta fliH fliB^*$ and $\Delta fliH fliJ fliB^*$ mutant strains. (c) Motility assays of MMH1001 ($\Delta fliH$) and MMH1017 ($\Delta fliH fliB^*$) transformed with pTrc99A (V) or pMMH1001 (FliH + FliI) in soft agar. The plates were incubated at 30 °C for 24 h. (d) Effect of external pH on FlgD secretion by the $\Delta fliH$ - $fliI$ - $fliJ fliB(P28T)$ mutant strain in the presence and absence of FliH and FliI. Immunoblotting, using polyclonal anti-FlgD antibody, of culture supernatant fractions prepared from MMH1017 carrying pTrc99A or pMMH1001 grown at an external pH of 6.0 and 7.0.

Interaction of FliJ($\Delta 13$ –24) with FliH and FliI. The $fliJ(\Delta 13$ –24) mutant caused six extragenic suppressors in FliH and three extragenic suppressors in FliI, raising the possibility that the deletion also affects the interaction with FliH and FliI. Therefore, we analysed the interaction of FliJ($\Delta 13$ –24) with FliH and FliI by pull-down assays. FliJ($\Delta 13$ –24) retained the ability to interact with FliH at wild-type levels and with FliI to an appreciable degree, although not at wild-type levels (Fig. 7a). This observation indicates that the deletion also weakens the FliJ–FliI interaction but not the FliJ–FliH interaction. The suppressor mutations in FliH and FliI did not significantly increase their binding affinity for FliJ($\Delta 13$ –24) (Fig. 7b). To test whether the same mutations influence the interaction of FliJ($\Delta 13$ –24) with FlhA, we analysed the interaction between FliJ($\Delta 13$ –24)

Figure 5 | Effect of a deletion of residues 328–351 of FlhA on the interaction with the soluble proteins.

(a) Pull-down assays by GST affinity chromatography. The mixture of the soluble fractions (L) prepared from a $\Delta fliH$ - $fliI$ mutant expressing GST (first row), GST-FliH (second row), GST-FliI (third row) or GST-FliJ (fourth row) with those from a $\Delta fliH$ mutant transformed with pMM108 (N-His-FLAG-FlhA, FlhA) or pMM108-1 (N-His-FLAG-FlhA($\Delta 328$ –351), FlhA($\Delta 328$ –351)) were loaded onto a GST column. After extensive washing, the proteins were eluted with a buffer containing 10 mM reduced glutathione. The eluted fractions were analysed by CBB staining for GST fusion proteins (upper panels) and immunoblotting with polyclonal anti-FlhA antibody (lower panels). (b) Effect of a $\Delta fliH$ - $fliI$ bypass mutation, FliB(P28T), on the FlhA($\Delta 328$ –351) mutants. Motility of NH0004 ($\Delta flhA \Delta fliH$ - $fliI fliB(P28T)$) transformed with pTrc99A (V), pMM108 (FlhA) or pMM108-1 (FlhA($\Delta 328$ –351)) in soft agar plate. Plates were incubated at 30 °C for 23 h.

and FlhA in the presence of the suppressor variants of the FliH–FliI complex. The suppressor mutation significantly improved the interaction of FliJ($\Delta 13$ –24) with FlhA (Fig. 7c).

Figure 6 | Effects of a deletion of residues 13-24 in FliJ on the FliJ-FlhA interaction. (a) Pull-down assays by GST affinity chromatography. The eluted fractions of GST-FliJ or GST-FliJ(Δ13-24) were analysed by CBB staining (upper panels), whereas the eluted FlhA protein was detected by immunoblotting with polyclonal anti-FlhA antibody. (b) Motility of a *ΔfliH-fliJ flhB(P28T)* (*ΔfliH fliJ flhB**) mutant transformed with pTrc99AFF4 (V), pMM404 (FliJ) or pRCJ104 (FliJ(Δ13-24)) in soft agar. The plates were incubated at 30 °C for 7 h. (c) Motility of pseudorevertants from the *fliJ(Δ13-24)* mutant: wild type (WT), *fliJ(Δ13-24)* mutant and its pseudorevertants (*fliJ(Δ13-24)*, *fliH(Δ96-97)* (labelled as *fliH**) and *fliJ(Δ13-24)*, *fliL(L244R)* (labelled as *fliL**)). (d) Secretion analysis of FlgD, FlgE and FliC. Immunoblotting, using polyclonal anti-FlgD, anti-FlgE or anti-FliC antibody of whole cell (Cell) and culture supernatant (Sup) fractions prepared from the above strains.

Because FliH and FliI directly bind to FlhA_C²¹, it is also possible that extragenic suppressor mutations in FliH or FliI might strengthen the FliH-FlhA or FliI-FlhA interaction, respectively, so that the FliJ-FlhA interaction becomes less critical. To test this

Figure 7 | Effect of the FliH and FliI suppressor mutations on the interaction of FliJ(Δ13-24) with FlhA. (a) Interactions of FliJ with FliH and FliI. Pull-down assays by GST affinity chromatography. The mixture of the soluble fractions (L) prepared from a *ΔflhDC-cheW* mutant expressing GST (left), GST-FliJ (middle) or GST-FliJ(Δ13-24) (right) with those from the *ΔflhDC-cheW* mutant producing His-FliH or His-FliI were loaded onto a GST column. After extensive washing, proteins were eluted with a buffer containing 10 mM reduced glutathione. (b) Interaction of GST-FliJ(Δ13-24) with the wild-type FliH-FliI complex (left) and its suppressor mutant variants (middle, FliH(Δ96-97) + FliI; right, FliH + FliI(L244R)). (c) Interactions of GST-FliJ(Δ13-24) with FlhA in the presence of a suppressor mutant variant of the FliH-FliI complex (FliH + FliI(L244R)). (d) Interaction of GST-FlhA_C with the suppressor mutant variants of FliH and FliI (FliH*, FliH(Δ96-97); FliI*, FliI(L244R)). The eluted fractions were analysed by CBB staining for GST fusion proteins (upper panels) and immunoblotting with polyclonal anti-FliH, anti-FliI or anti-FlhA_C antibody (lower panels).

possibility, we performed pull-down assays. FliH(Δ96-97) and FliI(L244R) both copurified with GST-FlhA_C at the wild-type levels (Fig. 7d), ruling out this possibility. Therefore, we suggest that the pseudorevertants with the second-site mutations are compensat-

ing for the *FliJ*($\Delta 13$ –24) mutation by altering the binding interface between *FliJ* and the linker of *FlhA*.

Discussion

The translocation of proteins across biological membranes requires protein transport machineries that convert biological energies into the mechanical work required for unidirectional protein transport. Nearly all the protein transport systems utilize free energy derived from cycles of nucleoside triphosphate binding and hydrolysis. In addition, PMF contributes to the driving force for protein transport across the bacterial cytoplasmic membrane, the mitochondrial inner membrane and so on²⁹. However, the mechanisms of protein translocation across biological membranes by PMF remain obscure.

It is thought that $\Delta\psi$ drives the translocation of negatively charged residues of preproteins across the membrane by an electrophoretic mechanism^{30–32}. In contrast, PMF functions on the membrane components of the protein translocation machinery, altering an interaction with *SecA* ATPase, which uses the energy of ATP binding and hydrolysis to drive preprotein translocation into and across the cytoplasmic membrane^{33–35}. Recently, it has been shown that the *SecD*/*SecE* complex utilizes PMF to function in the later ATP-independent stages of protein translocation³⁶. In mitochondria, where $\Delta\psi$ is the dominant driving force because of its high-level generation during aerobic respiration, the inside negative potential can exert an electrophoretic effect on positively charged polypeptides that are inserted into the membrane. $\Delta\psi$ also activates an inner membrane protein channel of the mitochondrial presequence translocase³⁷. Interestingly, it has been shown that $\Delta\psi$ can catalyse protein unfolding during protein translocation³⁸. Thus, PMF appears to have multiple roles in protein translocation across the biological membranes.

In this study, we have shown that $\Delta\psi$ alone is sufficient for flagellar protein export in wild-type cells, but both $\Delta\psi$ and ΔpH become essential in the absence of *FliH* and *FliI* (Fig. 1). These results indicate that the two components of PMF have distinct roles in the flagellar protein export process in the absence of *FliH* and *FliI*. This indication suggests that a step of the $\Delta\psi$ -dependent export process is different from that powered by ΔpH . The replacement of H_2O by D_2O reduced the level of *FlgD* secretion in the absence of *FliH* and *FliI* but did not reduce it in the wild-type cells (Fig. 3). Lowering the external pD value from 7.0 to 6.0 increased the secretion levels of *FlgD* in the absence of *FliH* and *FliI* (Fig. 3), indicating that the rate of proton transfer through the export gate determines the overall rate of flagellar protein export by the export gate alone. Taken together, we propose that the flagellar type III export gate complex made up of the six membrane proteins is intrinsically a proton–protein antiporter that requires both $\Delta\psi$ and ΔpH as the energy source. Proton transport could be an intrinsic aspect of the mechanism, which is not rate determining in the wild-type export apparatus. This possibility suggests a specific structural mechanism for the essential requirement of both components of the PMF to drive protein export, although the necessary structural information is currently not available.

The deletion of *FliJ* in addition to the deletion of *FliH* and *FliI* further reduced the level of export activity by the export gate but did not alter the ΔpH -dependent characteristics of the *fliH*-*fliI* bypass mutant (Fig. 4b). The export gate still requires ΔpH even in the presence of *FliH* and *FliI* when *FliJ* is missing (Fig. 4d). An increase in the ΔpH component of PMF significantly facilitated flagellar protein export in the *$\Delta fliH$ - $\Delta fliI$ - $\Delta fliJ$ *flhB*(P28T) mutant (Fig. 4b). Therefore, *FliJ* appears to be required for the full activation of the export gate complex for efficient, $\Delta\psi$ -driven protein export. The *FliJ*–*FlhA* interaction with residues 328–351 of *FlhA* in its linker region and residues 13–24 of *FliJ* was found to be critical for efficient protein export (Figs. 5 and 6). Because the *fliJ*($\Delta 13$ –24) suppressor mutations in *FliH* or *FliI* significantly improved the impaired ability of *FliJ*($\Delta 13$ –24) to interact with the linker of *FlhA* (Figs. 6 and 7), we*

propose that *FliJ* requires support of *FliH* and *FliI* to efficiently and properly interact with the linker of *FlhA* to switch the conformation of the export gate complex into the $\Delta\psi$ -driven mode, which appears to be much more efficient than the export gate made up of the six membrane proteins by themselves. The *flhB*(P28T) mutation restored motility of the *$\Delta fliH$ - $\Delta fliI$ - $\Delta fliJ$* triple null mutant to some degree (Fig. 4) but not that of the *$\Delta fliH$ - $\Delta fliI$ *flhA*($\Delta 328$ –351) mutant (Fig. 5), indicating that the linker region of *FlhA* is essential for flagellar protein export. We propose that a specific conformation of the *FlhA* linker region brought about by the proper binding of *FliJ* is critical for the efficient, $\Delta\psi$ -driven energy transduction mechanism of flagellar protein export.*

It has been proposed that *FliJ* is not only a general chaperone to prevent premature aggregation of export substrates in the cytoplasm¹⁴ but also a chaperone escort protein that promotes the cycling of substrate-specific chaperones, *FlgN* and *FliT*³⁹. The present study, however, has made it clear that *FliJ* is directly involved in a critical energy-transduction process with *FlhA*. *FliJ* facilitates *FliI* ring formation by binding to the centre of the ring, thereby stimulating the ATPase activity^{9,39}. ATP hydrolysis by *FliI* promotes the disassembly of the *FliI* ring, presumably for the rapid turnover of the process of substrate delivery and insertion to the export gate^{15,22}. Because *FliJ* binds to the linker of *FlhA* to fully activate the $\Delta\psi$ -driven protein export by the export gate (Fig. 5), we propose that *FliJ* is a key protein that switches the export gate, which by itself is a low-efficient proton–protein antiporter, into a highly efficient, $\Delta\psi$ -driven protein export apparatus. Thus, ‘export gate-activator switch’ would be a more appropriate name for *FliJ*.

Why is $\Delta\psi$ alone sufficient for protein export in wild-type cells? It has been shown that the flagellar export apparatus is evolutionarily related to F_0F_1 -ATP synthase, which utilizes PMF and ATP for ATP synthesis and proton pumping, respectively⁵. Although the structural mechanism of the flagellar type III protein export apparatus remains obscure, that of F_0F_1 -ATP synthase is well characterized: the α/β subunits form the $\alpha_3\beta_3$ hexameric ring as the core domain of F_1 ; the γ subunit forms the central stalk connecting F_1 with F_0 and is rotated within the central hole of the $\alpha_3\beta_3$ ring by the PMF-driven rotation of the F_0 ring in the membrane; the b and δ subunits form the peripheral stalk connecting F_1 with F_0 to function as the stator⁴⁰. In many of these aspects, the export apparatus shows similarities with F_0F_1 -ATP synthase: *FliI* forms a homo-hexamer ring to exert its ATPase activity^{41,42}; the structures of *FliI* and *FliJ* show extensive structural similarity to the α/β subunits⁸ and the γ subunit⁹, respectively; *FliJ* binds to the centre of the *FliI* ring in a manner similar to the antiparallel α -helical-coiled coil formed by the N- and C-terminal regions of the γ -subunit penetrating into the central cavity of the $\alpha_3\beta_3$ ring⁹; the amino-acid sequences of the N- and C-terminal regions of *FliH* are homologous to those of the b and δ subunits, respectively⁷; the export gate complex made up of the six integral membrane proteins is powered by PMF across the cytoplasmic membrane^{15,16}. Taken together, it may be possible that the export gate complex utilizes free energy derived from ATP hydrolysis by the *FliH*_x-*FliI*₆-*FliJ* ring complex for efficient proton translocation through the proton channel of the gate complex. Because the export gate by itself functions as a proton–protein antiporter, the proton translocation mechanism of the gate may also be similar to that of F_0F_1 -ATP synthase. However, because the present study has clearly shown that one of the important roles of the *FliH*–*FliI* ring complex is to bring *FliJ* to the *FlhA* linker region to switch the export gate from a low-efficient proton–protein antiporter to a highly efficient, $\Delta\psi$ -driven protein export apparatus, the structural similarity between the two systems in their entire assemblies may be used solely to achieve distinct mechanisms for their distinct functions. The detailed structural information of the entire export apparatus is essential for the mechanistic understanding of the flagellar protein export process in detail.

Table 1 | Strains and plasmids used in this study.

Strains and plasmids	Relevant characteristics	Source or reference
<i>Salmonella</i>		
SJW1103	Wild type for motility and chemotaxis	44
SJW1368	$\Delta(\text{cheW-flhD})$; master operon mutant	49
SJW104	<i>fliJ</i> ($\Delta 13-24$)	14
MKM40	$\Delta fliJ$	20
MMHI001	$\Delta fliH-fliI$	42
MMHIJ001	$\Delta fliH-fliI-fliJ$	This study
MMHIJ0117	$\Delta fliH-fliI-fliJ flhB$ (P28T)	This study
MM104H-xx series	<i>fliJ</i> ($\Delta 13-24$) <i>fliH</i> pseudorevertants from SJW104	This study
MM104I-xx series	<i>fliJ</i> ($\Delta 13-24$) <i>fliI</i> pseudorevertants from SJW104	This study
NH001	$\Delta flhA$	50
NH004	$\Delta fliH-fliI flhB$ (P28T) $\Delta flhA$	50
<i>Plasmids</i>		
pTrc99A	Expression vector	GE Healthcare
pGEX-6p-1	Expression vector	GE Healthcare
pGKK1702	pGEX-6p-1/GST-FliI	22
pMM104	pET19b/His-FlhA _C	11
pMM108	pTrc99A/His-FLAG-FlhA	28
pMM108	pTrc99A/His-FLAG- FlhA($\Delta 328-351$)	28
pMMHA003	pET19b/His-FlhA _C 38K	28
pMMHA1001	pGEX-6p-1/GST-FlhA _C	20
pMM306	pTrc99A/His-FliH	11
pMM404	pTrc99AFF4/FliJ	11
pMMJ1001	pGEX6p-1/GST-FliJ	20
pMM1702	pTrc99A/His-FliI	11
pMMHI001	pTrc99AFF4/FliH + FliI	15
pTIH001	pGEX6p-1/GST-FliH	21
pRCJ104	pTrc99AFF4/FliJ($\Delta 13-24$)	14
pMMHI104-3	pTrc99AFF4/FliH($\Delta 96-97$) + FliI	This study
pMMHI104-6	pTrc99AFF4/FliH + FliI(L244R)	This study
pMMJ1002	pGEX6p-1/GST-FliJ($\Delta 13-24$)	This study
pYVM001	pKK223-3/pHluorin(M153R)	47

Methods

Bacterial strains and media. *Salmonella* strains and plasmids used in this study are listed in Table 1. L-broth (LB) and soft tryptone agar plates were prepared as described in refs 11, 43. Motility medium contained 10 mM potassium phosphate, 0.1 mM EDTA and 10 mM sodium lactate. The pH value of each motility medium was adjusted to the desired pH by the addition of HCl or KOH in the presence or absence of potassium benzoate. Ampicillin was added to LB to a final concentration of 100 $\mu\text{g ml}^{-1}$.

DNA manipulations and transductional crosses. DNA manipulations were performed as described before²⁸. P22-mediated transductional crosses were performed with p22HT_{int} as described in ref. 44.

Construction of the *Salmonella* deletion mutants. A $\Delta fliH-fliI-fliJ$ triple null mutant and a $\Delta fliH-fliI-fliJ flhB$ (P28T) mutant were constructed using the λ -red homologous recombination system⁴⁵.

Motility assay. Fresh colonies were inoculated on soft tryptone agar plates and incubated at 30 °C. At least five independent experiments were conducted.

Secretion assay. The cells were grown with shaking in 5 ml of LB at 30 °C until the cell density had reached an OD₆₀₀ of ~1.0 and were washed twice with the motility buffer at pH 7.0. To test the effect of $\Delta\psi$ on flagellar protein export, we used valinomycin as described in ref. 16. To test the effects of ΔpH on the export, the cells were resuspended in the 5-ml motility buffer at pH 6.0, 6.5, 7.0 or 7.5 in the presence and absence of potassium benzoate and incubated at 30 °C for 1 h. To test the D₂O effect, the cells were resuspended in the 5-ml motility buffer prepared in D₂O and incubated at 30 °C for 1 h. Cultures were centrifuged to obtain cell pellets and culture supernatants. Cell pellets were resuspended in the SDS-loading buffer, followed by normalization to a cell density to yield a constant amount of cells.

Proteins in the culture supernatants were precipitated using 10% trichloroacetic acid, suspended in the Tris/SDS loading buffer and heated at 95 °C for 5 min. After SDS-PAGE, immunoblotting with polyclonal anti-FlgD, anti-FlgE and anti-FliC antibodies was performed as described before⁴³. Detection was performed with an ECL immunoblotting detection kit (GE Healthcare). At least three independent secretion assays were performed.

Measurement of the membrane potential. The membrane potential was measured as described by Lo *et al.*¹. Cells were suspended in motility medium plus 10 mM EDTA for 20 min. Cells were washed with motility medium and incubated in motility medium containing 0.1 μM TMRM (Invitrogen) for 10 min at room temperature. To observe bacterial cell bodies and epifluorescence of TMRM, we used an inverted fluorescence microscope (IX-71, Olympus) with a $\times 150$ oil immersion objective lens (UApo150XOTIRFM, NA 1.45, Olympus) and an electron-multiplying charged-coupled device camera (C9100-02, Hamamatsu Photonics), as described before⁴⁶. Calculation of the membrane potential was performed as described in ref. 1, with minor modifications. More than 100 wild-type cells and a *fliH-fliI* bypass mutant were measured at various external pH values with or without 20 mM potassium benzoate.

Intracellular pH measurement. Intracellular pH measurements of wild-type cells and a $\Delta fliH-fliI$ bypass mutant transformed with pYVM001, which encodes pHluorin(M153R) on pKK223-3 (ref. 47), were recorded at various external pH values with or without potassium benzoate, as described previously⁴⁸. The fluorescence-excitation spectra of the wild-type and $\Delta fliH-fliI$ bypass mutant cells expressing pHluorin(M153R) were recorded on a fluorescence spectrophotometer (RF-5300PC, Shimadzu). The 410/470-nm excitation ratios were calculated and converted to pH values based on the calibration curve. At least six independent measurements were recorded.

Pull-down assays by GST affinity chromatography. The soluble fractions prepared from SJW1364 ($\Delta flhDC-cheW$) expressing GST-FliH, GST-FliI, GST-Flj, GST-FliJ($\Delta 13-24$) or GST-FlhA_C were mixed with those from the $\Delta flhDC-cheW$ mutant transformed with pMM108 (His-FLAG-FlhA), pMM108-1 (His-FLAG-FlhA($\Delta 328-351$)), pMM306 (His-FliH), pMM1702 (His-FliI), pMM406 (His-FliJ), pMMHI001 (FliH + FliI), pMMHI104-3 (FliH($\Delta 96-97$) + FliI) or pMMHI104-6 (FliH + FliI(L244R)) were loaded onto a glutathione Sepharose 4B column (bed volume, 1 ml; GE Healthcare). After the column was extensively washed with 25 ml of PBS (8 g of NaCl, 0.2 g of KCl, 3.63 g of Na₂HPO₄·12H₂O, 0.24 g of KH₂PO₄, pH 7.4, per litre), proteins were eluted with 50 mM Tris-HCl, pH 8.0, and 10 mM reduced glutathione. Fractions containing GST, GST-FliH, GST-FliI, GST-FliJ, GST-FliJ($\Delta 13-24$) or GST-FlhA_C were identified by SDS-PAGE with Coomassie brilliant blue (CBB) staining and immunoblotting with polyclonal anti-FlhA_C, anti-FliH or anti-FliI antibody. Detection was performed with an ECL immunoblotting detection kit (GE Healthcare). At least three independent pull-down assays were performed.

References

- Lo, C. J., Leake, M. C., Pilizota, T. & Beryy, R. M. Nonequivalence of membrane voltage and ion-gradient as driving forces for the bacterial flagellar motor at low load. *Biophys. J.* **93**, 294–302 (2007).
- Wiedenmann, A., Dimroth, P. & von Ballmoos, C. Functional asymmetry of the F₀ motor in bacterial ATPsynthases. *Mol. Microbiol.* **72**, 479–490 (2009).
- Macnab, R. M. How bacteria assemble flagella. *Annu. Rev. Microbiol.* **57**, 77–100 (2003).
- Minamino, T. & Namba, K. Self-assembly and type III protein export of the bacterial flagellum. *J. Mol. Microbiol. Biotechnol.* **7**, 5–17 (2004).
- Minamino, T., Imada, K. & Namba, K. Mechanisms of type III protein export for bacterial flagellar assembly. *Mol. Biosyst.* **4**, 1105–1115 (2008).
- Cornelis, G. R. The type III secretion injectisome. *Nat. Rev. Microbiol.* **4**, 811–825 (2006).
- Pallen, M. J., Bailey, C. M. & Beatson, S. A. Evolutionary links between FliH/YscL-like proteins from bacterial type III secretion systems and second-stalk components of the FoF1 and vacuolar ATPases. *Protein Sci.* **15**, 935–941 (2006).
- Imada, K., Minamino, T., Tahara, A. & Namba, K. Structural similarity between the flagellar type III ATPase FliI and F₁-ATPase subunits. *Proc. Natl Acad. Sci. USA* **104**, 485–490 (2007).
- Ibuki, T., Imada, K., Minamino, T., Kato, T., Miyata, T. & Namba, K. Common architecture between the flagellar protein export apparatus and F- and V-ATPases. *Nat. Struct. Mol. Biol.* **18**, 277–282 (2011).
- Fan, F. & Macnab, R. M. Enzymatic characterization of FliI: an ATPase involved in flagellar assembly in *Salmonella typhimurium*. *J. Biol. Chem.* **271**, 31981–31988 (1996).
- Minamino, T. & Macnab, R. M. Interactions among components of the *Salmonella* flagellar export apparatus and its substrates. *Mol. Microbiol.* **35**, 1052–1064 (2000).
- Minamino, T. & Macnab, R. M. FliH, a soluble component of the type III flagellar export apparatus of *Salmonella*, forms a complex with FliI and inhibits its ATPase activity. *Mol. Microbiol.* **37**, 1494–1503 (2000).

13. González-Pedrajo, B., Fraser, G. M., Minamino, T. & Macnab, R. M. Molecular dissection of *Salmonella* FlhH, a regulator of the ATPase FlhI and the type III flagellar protein export pathway. *Mol. Microbiol.* **45**, 967–982 (2002).
14. Minamino, T., Chu, R., Yamaguchi, S. & Macnab, R. M. Role of FlhI in flagellar protein export in *Salmonella*. *J. Bacteriol.* **182**, 4207–4215 (2000).
15. Minamino, T. & Namba, K. Distinct roles of the FlhI ATPase and proton motive force in bacterial flagellar protein export. *Nature* **451**, 485–488 (2008).
16. Paul, K., Erhardt, M., Hirano, T., Blair, D. F. & Hughes, K. T. Energy source of flagellar type III secretion. *Nature* **451**, 489–492 (2008).
17. Thomas, J., Stafford, G. P. & Hughes, C. Docking of cytosolic chaperone-substrate complexes at the membrane ATPase during flagellar type III protein export. *Proc. Natl Acad. Sci. USA* **101**, 3945–3950 (2004).
18. Imada, K., Minamino, T., Kinoshita, M., Furukawa, Y. & Namba, K. Structural insight into the regulatory mechanisms of interactions of the flagellar type III chaperone FlhT with its binding partners. *Proc. Natl Acad. Sci. USA* **107**, 8812–8817 (2010).
19. Minamino, T., Gonzalez-Pedrajo, B., Kihara, M., Namba, K. & Macnab, R. M. The ATPase FlhI can interact with the type III flagellar protein export apparatus in the absence of its regulator, FlhH. *J. Bacteriol.* **185**, 3983–3988 (2003).
20. Minamino, T., Yoshimura, S. D., Morimoto, Y. V., González-Pedrajo, B., Kami-ike, N. & Namba, K. Roles of the extreme N-terminal region of FlhH for efficient localization of the FlhH-FlhI complex to the bacterial flagellar type III export apparatus. *Mol. Microbiol.* **74**, 1471–1483 (2009).
21. Minamino, T. *et al.* Role of the C-terminal cytoplasmic domain of FlhA in bacterial flagellar type III protein export. *J. Bacteriol.* **192**, 1929–1936 (2010).
22. Kazetani, K., Minamino, T., Miyata, T., Kato, T. & Namba, K. ATP-induced FlhI hexamerization facilitates bacterial flagellar protein export. *Biochem. Biophys. Res. Commun.* **388**, 323–327 (2009).
23. Minamino, T., Imae, Y., Oosawa, F., Kobayashi, Y. & Oosawa, K. Effect of intracellular pH on rotational speed of bacterial flagellar motors. *J. Bacteriol.* **185**, 1190–1194 (2003).
24. Miesenböck, G., Angelis, D. A. D. & Rothman, J. E. Visualizing secretion and synaptic transmission with pH-sensitive green fluorescent proteins. *Nature* **394**, 192–195 (1998).
25. Chen, X. & Berg, H. C. Solvent-isotope and pH effects on flagellar rotation in *Escherichia coli*. *Biophys. J.* **78**, 2280–2284 (2000).
26. Bange, G., Kümmerer, N., Engel, C., Bozkurt, G., Wild, K. & Sinning, I. FlhA provides the adaptor for coordinated delivery of late flagella building blocks to the type III secretion system. *Proc. Natl Acad. Sci. USA* **107**, 11295–11300 (2010).
27. Saijo-Hamano, Y. *et al.* Structure of the cytoplasmic domain of FlhA and implication for flagellar type III protein export. *Mol. Microbiol.* **76**, 260–268 (2010).
28. Saijo-Hamano, Y., Minamino, T., Macnab, R. M. & Namba, K. Structural and functional analysis of the C-terminal cytoplasmic domain of FlhA, an integral membrane component of the type III flagellar protein export apparatus in *Salmonella*. *J. Mol. Biol.* **343**, 457–466 (2004).
29. Wickner, W. & Schekman, R. Protein translocation across biological membrane. *Science* **310**, 1452–1456 (2005).
30. Cao, G., Kuhn, A. & Dalbey, R. E. The translocation of negatively charged residues across the membrane is driven by the electrochemical potential: evidence for an electrophoresis-like membrane transfer mechanism. *EMBO J.* **14**, 866–875 (2004).
31. Delgado-Partin, V. M. & Dalbey, R. E. The proton motive force, acting on acidic residues, promotes translocation of amino-terminal domains of membrane proteins when the hydrophobicity of the translocation signal is low. *J. Biol. Chem.* **273**, 9927–9934 (1998).
32. Schuenemann, T. A., Delgado-Nixon, V. M. & Dalbey, R. E. Direct evidence that proton motive force inhibits membrane translocation of positively charged residues within membrane proteins. *J. Biol. Chem.* **274**, 6855–6864 (1999).
33. Nouwen, N., de Kruijff, B. & Tommassen, J. *prfA* suppressors in *Escherichia coli* relieve the proton electrochemical gradient dependency of the translocation of wild-type precursors. *Proc. Natl Acad. Sci. USA* **93**, 5953–5957 (1996).
34. Nishiyama, K., Fukuda, A., Morita, K. & Tokuda, H. Membrane deinsertion of SecA underlying proton motive force-dependent stimulation of protein translocation. *EMBO J.* **18**, 1049–1058 (1999).
35. Mori, H. & Ito, K. Biochemical characterization of a mutationally altered protein translocation: proton motive force stimulation of the initiation phase of translocation. *J. Bacteriol.* **185**, 405–412 (2003).
36. Tsukazaki, T. *et al.* Structure and function of a membrane component SecDF that enhances protein export. *Nature* **474**, 235–238 (2011).
37. van der Laan, M., Hutu, D. P. & Rehling, P. On the mechanism of preprotein import by the mitochondrial presequence translocase. *Biochim. Biophys. Acta* **1803**, 732–739 (2010).
38. Huang, S., Ratliff, K. S. & Matouschek, A. Protein unfolding by the mitochondrial membrane potential. *Nat. Struct. Biol.* **9**, 301–307 (2002).
39. Evans, L. D., Stafford, G. P., Ahmed, S., Fraser, G. M. & Hughes, C. An escort mechanism for cycling of export chaperones during flagellum assembly. *Proc. Natl Acad. Sci. USA* **103**, 17474–17479 (2006).
40. Yoshida, M., Muneyuki, E. & Hisabori, T. ATP synthase—a marvellous rotary engine of the cell. *Nat. Rev. Mol. Cell Biol.* **2**, 669–677 (2001).
41. Claret, L., Susannah, C. R., Higgins, M. & Hughes, C. Oligomerisation and activation of the FlhI ATPase central to bacterial flagellum assembly. *Mol. Microbiol.* **48**, 1349–1355 (2003).
42. Minamino, T. *et al.* Oligomerization of the bacterial flagellar ATPase FlhI is controlled by its extreme N-terminal region. *J. Mol. Biol.* **360**, 510–519 (2006).
43. Minamino, T. & Macnab, R. M. Components of the *Salmonella* flagellar export apparatus and classification of export substrates. *J. Bacteriol.* **181**, 1388–1394 (1999).
44. Yamaguchi, S., Fujita, H., Sugata, K., Taira, T. & Iino, T. Genetic analysis of H2, the structural gene for phase-2 flagellin in *Salmonella*. *J. Gen. Microbiol.* **130**, 255–265 (1984).
45. Datsenko, K. A. & Wanner, B. L. One-step inactivation of chromosomal genes in *Escherichia coli* K-12 using PCR products. *Proc. Natl Acad. Sci. USA* **97**, 6640–6645 (2000).
46. Morimoto, Y. V., Nakamura, S., Kami-ike, N., Namba, K. & Minamino, T. Charged residues in the cytoplasmic loop of MotA are required for stator assembly into the bacterial flagellar motor. *Mol. Microbiol.* **78**, 1117–1129 (2010).
47. Morimoto, Y. V., Kojima, S., Namba, K. & Minamino, T. M153R mutation in a pH-sensitive green fluorescent protein stabilizes its fusion proteins. *PLoS One* **6**, e19598 (2011).
48. Morimoto, Y. V., Che, Y.-S., Minamino, T. & Namba, K. Proton-conductivity assay of plugged and unplugged MotA/B proton channel by cytoplasmic pHluorin expressed in *Salmonella*. *FEBS Lett.* **584**, 1268–1272 (2010).
49. Ohnishi, K., Ohto, Y., Aizawa, S.-I., Macnab, R. M. & Iino, T. FlgD is a scaffolding protein needed for flagellar hook assembly in *Salmonella typhimurium*. *J. Bacteriol.* **176**, 2272–2281 (1994).
50. Hara, N., Namba, K. & Minamino, T. Genetic characterization of conserved charged residues in the bacterial flagellar type III export protein FlhA. *PLoS One* **6**, e22417 (2011).

Acknowledgements

We acknowledge M. Kihara for continuous support and encouragement, and M. Kinoshita-Minamino and Y. Abe for technical assistance. Y.V.M. and N.H. are research fellows of the Japan Society for the Promotion of Science. This work was supported in part by Grants-in-Aid for Scientific Research (22570161 and 23121516 to T.M., and 21227006 to K.N.) and Targeted Proteins Research Program from the Ministry of Education, Culture, Sports, Science and Technology of Japan and by Takeda Science Foundation (to T.M.).

Author contributions

T.M. and K.N. conceived and designed the experiments. T.M., Y.V.M. and N.H. conducted the experiments. T.M. and Y.V.M. analysed the data. T.M. and K.N. wrote the paper.

Additional information

Supplementary Information accompanies this paper at <http://www.nature.com/naturecommunications>

Competing financial interests: The authors declare no competing financial interests.

Reprints and permission information is available online at <http://ngp.nature.com/reprintsandpermissions/>

How to cite this article: Minamino, T. *et al.* An energy transduction mechanism used in bacterial flagellar type III protein export. *Nat. Commun.* **2**:475 doi: 10.1038/ncomms1488 (2011).

License: This work is licensed under a Creative Commons Attribution-NonCommercial-Share Alike 3.0 Unported License. To view a copy of this license, visit <http://creativecommons.org/licenses/by-nc-sa/3.0/>