ARTICLE

Received 12 Apr 2012 | Accepted 27 Jul 2012 | Published 4 Sep 2012

DOI: 10.1038/ncomms2023

Extracellular ATP mediates mast cell-dependent intestinal inflammation through P2X7 purinoceptors

Yosuke Kurashima^{1,2,3}, Takeaki Amiya^{1,3,4}, Tomonori Nochi¹, Kumiko Fujisawa^{1,3}, Takeshi Haraguchi⁵, Hideo Iba⁵, Hiroko Tsutsui⁶, Shintaro Sato^{1,3}, Sachiko Nakajima⁷, Hideki Iijima⁷, Masato Kubo^{8,9}, Jun Kunisawa^{1,4} & Hiroshi Kiyono^{1,2,3,4}

Mast cells are known effector cells in allergic and inflammatory diseases, but their precise roles in intestinal inflammation remain unknown. Here we show that activation of mast cells in intestinal inflammation is mediated by ATP-reactive P2X7 purinoceptors. We find an increase in the numbers of mast cells expressing P2X7 purinoceptors in the colons of mice with colitis and of patients with Crohn's disease. Treatment of mice with a P2X7 purinoceptor-specific antibody inhibits mast cell activation and subsequent intestinal inflammation. Similarly, intestinal inflammation is ameliorated in mast cell-deficient *Kit^{W-sh/W-sh}* mice, and reconstitution with wild-type, but not *P2x7^{-/-}* mast cells results in susceptibility to inflammation. ATP-P2X7 purinoceptor-mediated activation of mast cells not only induces inflammatory cytokines, but also chemokines and leukotrienes, to recruit neutrophils and subsequently exacerbate intestinal inflammation. These findings reveal the role of P2X7 purinoceptor-mediated mast cell activation in both the initiation and exacerbation of intestinal inflammation.

¹ Division of Mucosal Immunology, Department of Microbiology and Immunology, The Institute of Medical Science, The University of Tokyo, 108-8639, Japan. ² Graduate School of Medicine, The University of Tokyo, 113-0033, Japan. ³ Core Research for Evolutional Science and Technology (CREST), Japan Science and Technology Agency (JST), Tokyo 102-0075, Japan. ⁴ Department of Medical Genome Science, Graduate School of Frontier Science, The University of Tokyo, Chiba 277-8561, Japan. ⁵ Division of Host-Parasite Interaction, Department of Microbiology and Immunology, The Institute of Medical Science, The University of Tokyo, 108-8639, Japan. ⁶ Department of Microbiology, Hyogo College of Medicine, Nishinomiya 663-8501, Japan. ⁷ Department of Gastroenterology and Hepatology, Osaka University Graduate School of Medicine, 565-0871, Japan. ⁸ Research Center for Allergy and Immunology, RIKEN, Yokohama Institute, Tsurumi, Yokohama, Kanagawa, 230-0045, Japan. ⁹ Division of Molecular Pathology, Research Institute for Biological Sciences, Tokyo University of Sciences, Chiba 278-0022, Japan. Correspondence and requests for materials should be addressed to J.K. (email: kunisawa@ims.u-tokyo.ac.jp) or to H.K. (email: kiyono@ims.u-tokyo.ac.jp).

Both active and quiescent immunity occur simultaneously to achieve immunological homeostasis in the harshest of environments—namely, the intestine. Aberrant immune responses in the gut lead to the development of intestinal immune diseases such as colitis and food allergies^{1,2}. Mast cells (MCs) are generally recognized as major effector cells of type 1 allergic diseases, as well as of inflammation, host defenses, innate and adaptive immune responses and homeostatic responses^{3–5}. Histological analyses of patients with, and murine models of, colitis have implicated the involvement of MCs in intestinal inflammation^{4,6}, but the factors responsible for MC activation are not fully understood.

Several lines of evidence have demonstrated that release of extracellular ATP and ADP from injured, dying or activated cells acts as a danger signal by modulating various cellular functions via the activation of P2 purinoceptors^{7,8}. P2 purinoceptors comprise P2X (P2X_{1–7}) and P2Y receptors (P2Y_{1, 2, 4, 6, 11–14}). P2X_{1–7} receptors are ATP-gated ion channels and specific for ATP, whereas P2Y receptors are G protein-coupled receptors that are specific for ADP, UTP and ATP^{7,8}.

Stimulation by ATP or ADP through the P2 purinoceptors of macrophages and dendritic cells (DCs) results in the production of inflammatory cytokines; this can lead to the development of asthma, contact hypersensitivity or graft-versus-host disease^{9–11}. MCs also express several P2 purinoceptors and release histamine, cytokines and chemokines upon nucleotide stimulation¹². Although MCs are thought to be involved in intestinal inflammation, it is unclear whether extracellular nucleotides are required for this process.

Here, we used a newly established anti-MC monoclonal antibody (mAb) to identify activated MCs and found that extracellular ATP mediates MC activation through P2X₇ purinoceptors to initiate and amplify intestinal inflammation. Consequently, obstruction of the ATP-P2X₇ purinoceptor cascade could be used to inhibit gut inflammatory diseases.

Results

Activated MCs in intestinal inflammation. Using a 2,4,6-trinitrobenzene sulphonic acid (TNBS)-induced colitis model, we first examined whether MCs were involved in intestinal inflammation. To assess MC activation *in vivo*, we established an mAb (clone: 5A9) specific for CD63, a marker of activated MCs¹³. We confirmed that our anti-CD63 mAb was reactive specifically to MCs activated by immunoglobulin (Ig)E plus relevant allergen or a calcium ionophore, and not to naïve and CD63-knockdown MCs (Supplementary Fig. S1). In the colons of TNBS-treated mice, increased numbers of CD63⁺-activated MCs were noted until day 3 post administration; the numbers then gradually decreased and reached a basal level on day 6 (Fig. 1a,b), indicating that MC activation was associated with the initiation phase of colitis development, as previously reported in a murine model and in patients with inflammatory bowel disease^{6,14}. It has generally been accepted that the mechanistic basis of ulcerative colitis (UC) and Crohn's disease (CD) are different. Indeed, the pathogenic cytokines involved in the development of UC and CD are different² and the genetic polymorphisms specific for UC and CD are also different¹⁵. In addition, the cytokines required for the development of MCs differ between humans (stem cell factor) and mice [interleukin (IL)-3 and stem cell factor]⁴. Thus, these different pathological environments may have led to differences in the requirement for, and involvement of, MCs in the development of inflammation. Therefore, we analysed MC numbers in both UD and CD patients, although we focused on the TNBS-induced colonic inflammation model. We detected increased numbers of MCs in the colons of patients with CD or UC (Fig. 1c,d). Thus, increased numbers of MCs in the colon is a characteristic of intestinal inflammation.

To directly show the involvement of MCs in the development of intestinal inflammation, we used MC-deficient *Kit^{W-sh/W-sh}* mice. We

confirmed that immunological and inflammatory symptoms induced by TNBS treatment were identical in *Kit^{W-sh/+}* heterozygous and *Kit^{+/+}* homozygous mice; however, inflammatory symptoms, such as body weight loss, massive inflammatory cell infiltration and colon shortening, were restored in *Kit^{W-sh/W-sh}* mice but not in *Kit^{W-sh/+}* heterozygous and *Kit^{+/+}* homozygous mice (Fig. 1e–h). Similarly, our histological and immunological analyses revealed that destruction of the colonic epithelial layer and infiltration by inflammatory cells—especially neutrophils, which were stained neutral pink and had lobulated nuclei,—were reduced in *Kit^{W-sh/W-sh}* mice (Fig. 1f,h,i). Moreover, inflammatory signs were ameliorated in *Kit^{W-sh/W-sh}* mice when we used other well-known inflammatory bowel disease models, such as the dextran sodium sulphate (DSS) colitis model (Fig. 2a–c). As the use of *Kit^{W-sh/W-sh}* mice as an MC-deficient model is controversial^{16,17}, we also used the MC-specific enhancer-mediated toxin receptor-mediated conditional cell knockout (TRECK) system (Mas-TRECK mice)¹⁸. We confirmed that specific depletion of MCs ameliorated the inflammation in this DSS-induced colitis model (Fig. 2d–h). Our data indicate that activated MCs participate in the aggravation of intestinal inflammation.

Establishment of an inhibitory MC-specific mAb. IgE plus a relevant allergen induces MC activation; however, *Rag-1^{-/-}* and *Tcrβ^{-/-} δ^{-/-}* mice showed inflammatory responses comparable to those in TNBS-induced intestinal inflammation (Supplementary Fig. S2a–d)¹⁹ and had increased numbers of CD63⁺-activated MCs in their colons (Supplementary Fig. S2e), suggesting that T and B cells are not involved in MC activation during colitis. We also found no increase in CD63 expression on MCs after stimulation with IL-18 and IL-33, which are known to be involved in colitis (Supplementary Fig. S2f)^{20,21}.

We next tried to establish an anti-MC mAb that could ameliorate activated MC-mediated intestinal inflammation. We immunized rats with purified murine-activated colonic MCs, established hybridomas, performed flow cytometry to select hybridomas that produced mAbs that preferentially recognized colonic MCs and examined the hybridomas' ability to inhibit ovalbumin-induced food allergy²² or TNBS-induced intestinal inflammation (Supplementary Fig. S3). Among 2,000 clones, we obtained an anti-MC mAb (designated clone 1F11; rat IgG2b) that was strongly reactive to colonic MCs (Fig. 3a; Supplementary Fig. S3). In addition to colonic MCs, 1F11 mAb bound efficiently to peritoneal cavity-, lung- and bone marrow (BM)-derived MCs, but not to skin MCs (Fig. 3a). When tested with other immunocompetent cells in the colon, 1F11 mAb was weakly reactive to some CD3⁺ T cells, CD11c⁺ DCs and F4/80⁺ macrophages, but was not reactive to Gr-1⁺ granulocytes, IgA⁺ plasma cells or epithelial cells (ECs) (Fig. 3b).

To show the inhibitory function of 1F11 mAb in intestinal inflammation, mice were given 1F11 mAb (0.5 mg day⁻¹ in a single dose) for 3 days, beginning 1 day before intrarectal administration of TNBS. 1F11 mAb treatment reduced the intestinal inflammation (Fig. 3c–g) and decreased the number of CD63⁺-activated MCs in 1F11 mAb-treated mice (Fig. 3h).

Targeting P2X₇ receptors reduces intestinal inflammation. Mass spectrometry analyses of immunoprecipitants of MC cell lysates with 1F11 mAb showed that the P2X₇ purinoceptor is recognized by 1F11 mAb (Supplementary Fig. S4a). The specificity of 1F11 mAb for the P2X₇ purinoceptor was confirmed by its specific reactivity to cells transfected with P2X₇ receptors but not with other types of P2X receptor (for example, P2X₁ and P2X₄; Supplementary Fig. S4b). MCs derived from *P2x7^{-/-}* mice, however, were not recognized by 1F11 mAb (Supplementary Fig. S4c). Western blot and flow cytometric analysis showed that, among the several variants of P2X₇ purinoceptors²³, 1F11 mAb bound to variant a (full-length; Supplementary Fig. S4d,e). In contrast, variant c (possessing

Figure 1 | Role of activated intestinal MCs in the development of intestinal inflammation. (a) CD63 expression on colonic MCs was examined with flow cytometry. Cells were gated on c-kit⁺ and FcεR1α⁺ cells. (b) The percentage of CD63⁺ MCs in all c-kit⁺ FcεR1α⁺ MCs was determined with flow cytometry at various time points after TNBS administration ($n=3$ for day 6, $n=5$ for day 3, $n=7$ for intact, ETOH, day 1 and 2, $n=14$ for day 4). Control mice were analysed 4 days after ETOH administration (ETOH; $n=7$). Data are shown as means $\pm$ s.e.m. (c) Colonic tissue sections from a healthy volunteer (HV) and UC and CD patients were stained with 4',6-diamidino-2-phenyl indole (blue) and MC tryptase (red) or haematoxylin and eosin (H&E) (bottom). Scale bars, 100 μ m. (d) Tryptase-positive MCs were counted in the fields of the tissue sections (four fields for each section). Data are means $\pm$ s.e.m. ($n=6$). (e) Body weight changes were monitored after TNBS administration to *Kit*^{W-sh/W-sh} MC-deficient mice (*Kit*^{W-sh/W-sh} ETOH; open triangles; $n=4$, *Kit*^{W-sh/W-sh} TNBS; closed triangles; $n=9$), *Kit*^{+/+} control mice (*Kit*^{+/+} ETOH; open diamonds; $n=4$, *Kit*^{+/+} TNBS; closed diamonds; $n=13$) and *Kit*^{W-sh/W+} control mice (*Kit*^{W-sh/W+} ETOH; open squares; $n=4$, *Kit*^{W-sh/W+} TNBS; closed squares; $n=11$). Data are shown as percentages of baseline weights and are means $\pm$ s.e.m., * $P<0.0001$ (two-tailed Student's *t*-test); ** $P=0.0024$ (two-tailed Student's *t*-test). (f) The colon was isolated 4 days after TNBS treatment for H&E staining. Data are representative of at least three independent experiments. Scale bars, 100 μ m. (g) Colon length was measured 4 days after colitis induction. ETOH, closed column; TNBS, open column. * $P<0.0001$ (two-tailed Student's *t*-test), ** $P=0.0024$ (two-tailed Student's *t*-test). Data are shown as means $\pm$ s.e.m. (h) The percentage of CD11b⁺ Gr-1^{high} cells in the colonic lamina propria was calculated, as measured with flow cytometry. ETOH, closed column; TNBS, open column. * $P=0.0003$ (two-tailed Student's *t*-test), ** $P=0.0029$ (Welch's *t*-test) and *** $P<0.0001$ (Welch's *t*-test). Data are shown as means $\pm$ s.e.m. (i) Colonic mononuclear cells were isolated 4 days after TNBS administration and stained with anti-CD11b and anti-Gr-1 antibodies. CD11b⁺ Gr-1^{high} cells were sorted and then stained with May-Giemsa stain. Scale bar, 20 μ m. Data are representative of three experiments.

the ATP-binding portion but lacking the C-terminal region) was detected by western blot, but its surface expression was not detected by flow cytometry because of its defect in extracellular expression (Supplementary Fig. S4d,e)²⁴. In addition, neither western blot nor flow cytometry detected variant d (lacking the ATP-binding portion; Supplementary Fig. S4d,e). These data strongly suggest that 1F11 mAb recognizes P2X7 receptors, specifically the ATP-binding portion. We also confirmed that 1F11 mAb had similar reactivity to that of a commercially available anti-P2X7 mAb (clone: Hano43; Supplementary Fig. S4f,g).

To evaluate whether 1F11 mAb directly affects MCs during ATP-mediated activation, we treated MCs with ATP in the presence of 1F11 mAb *in vitro*. 1F11 mAb treatment reduced the number of CD63⁺-activated MCs induced by ATP in a dose-dependent manner (Fig. 4a). High concentrations of extracellular ATP increased the

cell permeability of the MCs¹². Thus, uptake of Lucifer yellow was observed in ATP-stimulated MCs but was substantially impaired in 1F11 mAb-treated and *P2x7*^{-/-} MCs (Fig. 4b,c).

As many cell types (MCs, T cells and DCs) express P2X7 receptors (Fig. 3b), we then asked whether the P2X7 receptors on MCs were responsible for the MC-mediated intestinal inflammation *in vivo* by analysing MC-deficient *Kit*^{W-sh/W-sh} mice reconstituted with *P2x7*^{+/+} or *P2x7*^{-/-} MCs. We confirmed that reconstituted MCs were present in the colon and maintained P2X7 expression (Supplementary Fig. S5). Like wild-type mice, *Kit*^{W-sh/W-sh} mice reconstituted with *P2x7*^{+/+} MCs showed severe inflammatory responses when treated with TNBS. However, these inflammatory responses were ameliorated when *Kit*^{W-sh/W-sh} mice were reconstituted with *P2x7*^{-/-} MCs; the amelioration included inhibition of neutrophil infiltration and MC activation (Figs 1 and 5a–f).

Figure 2 | Impaired DSS-induced colitis in MC-deficient mice. *Kit^{W-sh/W-sh}* MC-deficient, *Kit^{+/+}* control mice and Mas-TRECK transgenic (Tg) mice were subjected to DSS-induced colitis. **(a)** Body weight changes are shown as percentages of the baseline value and are means $\pm$ s.e.m. ($n = 22$ for *Kit^{+/+}*; $n = 25$ for *Kit^{W-sh/+}*; $n = 10$ for *Kit^{W-sh/W-sh}*). * $P < 0.01$, ** $P = 0.0207$ and *** $P = 0.0004$ (two-tailed Student's *t*-test). **(b,c)** Eleven days after DSS treatment, colon tissue and haematoxylin and eosin (H&E)-stained tissue sections were examined. Data are representative of at least three independent experiments. **(d)** Mas-TRECK Tg mice and their wild-type (WT) littermates were subjected to DSS-induced colitis. For diphtheria toxin (DT) treatment, mice were injected intraperitoneally with 250 ng of DT for 5 consecutive days (black arrows) and then with 150 ng every other day (red arrows). **(e)** Body weight changes are shown as percentages of the baseline value and are means $\pm$ s.e.m. ($n = 6$ for Tg; $n = 10$ for WT), * $P = 0.0107$, ** $P = 0.0037$ (two-tailed Student's *t*-test). **(f)** Representative images of whole colons 10 days after DSS treatment. **(g)** Representative images of H&E staining. Scale bars, 100 μ m. **(h)** Representative flow cytometric data of infiltrated c-kit⁺ Fc ϵ R1 α ⁺ MCs in the colon.

We next analysed whether the MCs in UC or CD patients expressed P2X7. Although increased number of MCs were observed in the colons of both UC and CD patients (Fig. 1c,d), P2X7 purinoceptors were expressed by the MCs in CD patients but not by those in UC patients or healthy volunteers (Fig. 5g,h). Thus, it is likely that P2X7 purinoceptor-mediated MC activation also occurs in the human colon, especially in CD patients.

To examine whether ATP was extracellularly released at high concentrations at inflammatory sites, we next measured ATP release from inflammatory colonic tissues. An elevated level of ATP release from the colon tissue was noted in TNBS-treated mice (Fig. 6a). In addition, intrarectal administration of non-hydrolyzable ATP (adenosine 5'-O-(3-thio) triphosphate and O-(4-benzoyl)benzoyl adenosine 5'-triphosphate) led to MC activation in the colonic tissue, similar to the effect of TNBS treatment (Fig. 6b). In contrast, intrarectal administration of other P2Y receptor agonists did not increase colonic MC activation (Fig. 6b). These findings indicate that inflammatory stimuli induce the extracellular release of ATP, which in turn leads to P2X7-dependent MC activation in the colon and subsequent exacerbation of intestinal inflammation.

P2X7 signalling activates the caspase-1 inflammasome to induce the production of IL-1 β and IL-18 (ref. 25). IL-1 β production is also mediated by MC proteases, such as chymases²⁶. We therefore examined whether MCs produced IL-1 β via P2X7 receptor activation, and if so whether this production was caspase-1-dependent. IL-1 β production was decreased when P2X7-deficient MCs were stimulated with ATP, whereas substantial amounts of IL-1 β were produced in caspase-1-deficient MCs (Supplementary Fig. S6), indicating that IL-1 β production was P2X7-dependent but caspase-1-independent. In line with this finding, body weight changes were noted in *Kit^{W-sh/W-sh}* mice reconstituted with *caspase-1^{-/-}*

MCs (Fig. 5a). These results suggest that MC-dependent inflammation through P2X7 purinoceptors is not dependent on caspase-1-mediated IL-1 β or IL-18 production.

An autocrine loop of ATP conversion mediates MC activation. In addition to ATP, other nucleotides (for example, extracellular ADP) act as signals to induce inflammatory responses²⁷. We confirmed that MCs are activated by high concentrations of ADP and ATP (Fig. 7a,b). Extracellular ATP is hydrolysed by ectonucleoside triphosphate diphosphohydrolases (CD39) to ADP and AMP; it is then further hydrolysed by ecto-5'-nucleotidase (CD73) to adenosine, which has anti-inflammatory functions²⁷. Colonic MCs expressed CD39 but not CD73 (Supplementary Fig. S7a,b), indicating that MCs can convert ATP to ADP but not to adenosine. We therefore examined the involvement of ADP-reactive P2Y purinoceptors and found that P2Y1 and P2Y12 were highly expressed on colonic MCs (Fig. 7c). However, inhibitors of P2Y1 and P2Y12 receptors, as well as knockdown of the P2Y12 receptor, had no effect on the induction of CD63⁺-activated MCs (Fig. 7d,e; Supplementary Fig. S8a). Similarly, intestinal inflammation, as well as activation of colonic MCs, was unaffected in clopidogrel (a P2Y12 receptor inhibitor)-treated mice (Supplementary Fig. S8b–d). These data indicate that although P2Y1 and P2Y12 were expressed on MCs neither P2Y1 nor P2Y12 purinoceptors mediate ADP-dependent CD63⁺ MC induction.

It is generally accepted that P2X7 purinoceptors specifically recognize ATP⁷, but we found that they were also involved in ADP-mediated MC activation. Indeed, no activation was noted in *P2x7^{-/-}* MCs when they were stimulated with ADP (Fig. 7f), leading us to hypothesize that ADP promotes ATP release from MCs and their subsequent stimulation. To test this hypothesis, we measured the expression of pannexin-1, connexin 43 and connexin 32, which

Figure 3 | Amelioration of colitis by treatment with intestinal MC-reactive 1F11 mAb. (a) MCs in the colonic lamina propria, peritoneal cavity (PerC), skin and lung, as well as BM-derived MCs, were stained with 1F11 mAb. Control staining with rat IgG2b is shown in grey. (b) Cells were isolated from colonic lamina propria and epithelium. CD3⁺ T cells, IgA⁺ plasma cells (PCs), F4/80⁺ macrophages, CD11c⁺ DCs, Gr1⁺ granulocytes and ECs were gated and their reactivity to 1F11 mAb examined. Control staining with rat IgG2b is shown in grey. (c) C57BL/6 mice were treated with TNBS and their body weights were monitored for 4 days; 0.5 mg of 1F11 or the control mAb was intraperitoneally administered. Data from 9 (EtOH; open squares), 19 (TNBS with control mAb; closed triangles) and 12 (TNBS with 1F11 mAb; closed diamonds) mice. * $P=0.0066$ (Welch's t -test). Data are shown as percentages of baseline weights and are means $\pm$ s.e.m. (d,e) Representative images of haematoxylin and eosin staining and colon tissue from 1F11 mAb-treated mice. Scale bars, 100 μ m. (f) Colon length was measured 4 days after TNBS administration. * $P=0.0445$; ** $P=0.0073$ (two-tailed Student's t -test). (g) Neutrophils (CD11b⁺ Gr-1^{high}) were quantified as percentages and numbers of cells. Data are shown as means $\pm$ s.e.m. ($n=6$), * $P<0.0001$, ** $P=0.0047$ (two-tailed Student's t -test). (h) Percentage of CD63⁺ MCs in all c-kit⁺ FcεR1α⁺ MCs was determined with flow cytometry. Data are shown as means $\pm$ s.e.m. ($n=6$) * $P=0.0202$; ** $P=0.0284$ (two-tailed Student's t -test).

are ATP-releasing hemichannels, during cell activation^{28,29}. The hemichannels were rarely expressed on the colonic MCs (Fig. 7g), and no inhibitory effect was observed when the MCs were treated with ADP in the presence of hemichannel inhibitors (flufenamic acid and carbenoxolone). However, cell activation was inhibited by P2X7 antagonists [oxidized ATP (OxATP), pyridoxal-phosphate-6-azophenyl-2',4'-disulfonate and 4,4'-diisothiocyanatostilbene-2,2'-disulfonic acid disodium salt hydrate] (Fig. 7h). To further exclude the possibility that ADP triggers ATP release, we stimulated MCs with another P2Y ligand (UTP); we found that UTP did not induce MC activation (Fig. 7b).

We then tested whether ADP was converted to ATP by ATP-converting enzymes such as ecto-adenylate kinase, ATP synthase and nucleoside diphosphokinase³⁰. To test the involvement of these enzymes, we used inhibitors of ecto-adenylate kinase (diadenosine pentaphosphate; AD2P5), ATP synthase (oligomycin; oligo) and nucleoside diphosphokinase (UDP), and we found that inhibition of ecto-adenylate kinase and ATP synthase, but not nucleoside diphosphokinase, reduced ADP- as well as ATP-dependent MC activation (Fig. 7h,i). Neither AD2P5 nor oligo inhibited MC activation induced by the crosslinking of IgE with relevant allergen (Fig. 7i). Among the adenylyl kinases, adenylyl kinase 1 (AK1) and AK2 were expressed in colonic MCs, and the expression of AK2 was much higher than that of AK1 (Supplementary Fig. S9a). As with AD2P5 treatment, knockdown of AK2, but not AK1, led to the

inhibition of both ADP- and ATP-mediated MC activation (Supplementary Fig. S9b). These results indicate that P2X7 purinoceptors have an important role in the activation of MCs by ATP, including ATP derived from ADP by the action of ecto-enzymes such as ATP synthase and AK2.

Neutrophil infiltration by MC-derived mediators. Evaluation of MC activation on the basis of CD63 expression is an important criterion¹³; however, degranulation is not absolutely associated with cytokine production³¹. Therefore, we measured MC production of an array of inflammatory cytokine, chemokine and lipid mediators to additionally elucidate the role of P2X7 purinoceptor-mediated MC activation in the development of intestinal inflammation. Stimulation of MCs with ATP induced the production of inflammatory cytokines such as IL-6, tumour necrosis factor (TNF)α and oncostatin M³²; this induction was not observed in $P2x7^{-/-}$ MCs or in wild-type MCs treated with 1F11 mAb (Fig. 8a,b).

We showed that neutrophil infiltration into the colon was mediated by MC activation (Fig. 1h,i), and a previous study suggested that neutrophil infiltration is a potential target in colitis treatment³³. Consistent with these findings, ATP stimulation induced MCs, but not $P2x7^{-/-}$ MCs, to produce leukotrienes (LTs; LT C4/D4/E4), which are associated with the translocation of 5-lipoxygenase (5-LO) into the nucleus—an important step for LT synthesis in MCs³⁴ (Fig. 8c,d). Also, chemokine gene array analysis demonstrated that

Figure 4 | Inhibition of *in vitro* ATP-mediated MC activation by 1F11 mAb. (a) BM-derived MCs, pretreated with various concentrations of 1F11 mAb (0, 1, 10 $\mu\text{g ml}^{-1}$) for 15 min, were stimulated with 0.5 mM ATP for 30 min. Cells were stained with an anti-CD63 mAb for flow cytometric analysis. Data are representative of three independent experiments. (b) BM-derived MCs pretreated with various concentrations of 1F11 mAb or control rat IgG2b (0, 10 $\mu\text{g ml}^{-1}$) for 15 min were stimulated with 0.5 mM ATP for 30 min in the presence of 1 mg ml^{-1} Lucifer yellow (LY). (c) LY uptake was determined by using flow cytometry and fluorescence microscopy. Scale bar, 100 μm . Data are representative of three individual experiments.

ATP stimulation of MCs induced the expression of chemokines, including CCL2, CCL7 and CXCL2 (Fig. 8e–g), and 1F11 mAb treatment or P2X7 deficiency resulted in decreased CCL2 production from MCs activated by ATP but not by IgE plus allergen (Fig. 8g). Furthermore, *Kit^{W^{sh}/W^{sh}}* mice showed decreased levels of both CCL2 and IL-1 β in the colon tissue, but the production levels of these molecules recovered when the mice were reconstituted with wild-type MCs (Supplementary Fig. S10a). As neutrophils express the corresponding chemokine receptors, it is likely that ATP-dependent MC activation induced inflammatory neutrophil infiltration into the colon from the peripheral blood (Supplementary Fig. S10b,c), given the high level of TNF α production by the neutrophils (Supplementary Fig. S10d). These results indicate that ATP-dependent MC activation has key roles in the induction of inflammatory responses (by inducing inflammatory cytokines) and in the exacerbation of inflammatory responses (by inducing LTs and chemokines to recruit TNF α -producing neutrophils to the colon).

Discussion

Here, we showed that MCs have a critical role in the severity of colitis through their interaction with ATP and P2X7 purinoceptors. These interactions not only induce MC-mediated inflammatory responses but also exacerbate them by promoting neutrophil infiltration. Indeed, MC-deficient mice reconstitution with wild-type, but not *P2x7^{-/-}*, MCs resulted in neutrophil infiltration and severe inflammatory responses, together with increased production of IL-1 β , LTs and CCL2 (Figs 5 and 8, and Supplementary Fig. S10). *Kit^{W^{sh}/W^{sh}}* mice spontaneously show elevated levels of neutrophils in their spleens³⁵; however, we confirmed that the neutrophil levels

were the same as those in the colons of *Kit^{+/+}*, *Kit^{W^{sh}/+}* and *Kit^{W^{sh}/W^{sh}}* mice under naïve conditions (Fig. 1h,i). To exclude the possible involvement of other immunological defects in *Kit^{W^{sh}/W^{sh}}* mice, such as the involvement of the *Corin* gene, which is associated with type II transmembrane serine protease³⁵, we further confirmed the amelioration of intestinal inflammation in conditional MC-deficient mice (Fig. 2d–h). These findings strongly suggest that P2X7 on MCs has a pivotal role in the development of murine and human intestinal inflammation.

P2X7 purinoceptors are expressed on T cells, DCs, macrophages and ECs^{9–11,25,36}. In a recent study, ATP/P2X7-mediated signaling inhibited the generation and function of regulatory T cells and ATP stimulation led to their conversion into Th17 cells via an IL-6-dependent pathway; thus, the P2X7 antagonist OxATP inhibited colitis³⁷. In that study, ATP/P2X7-mediated regulation of regulatory T cells was involved in the chronic phase of intestinal inflammation, which takes about 4 weeks for disease development³⁷. Similarly, ATP-mediated DC activation occurs in the chronic phase of intestinal inflammation through the preferential induction of Th17 cells, although whether this is mediated by P2X7 remains to be seen³⁸. In contrast, ATP/P2X7-mediated MC activation in our model was important in the development of T-cell-independent acute colitis, which occurs within 1 week. Thus, our study and those of others^{37,38} complement each other by reflecting the complicated pathological aspects and kinetics of the acute and chronic phases of intestinal inflammation mediated by ATP and P2X7.

We also found that the expression level of P2X7 receptors differed depending on the tissue and animal species. First, colonic MCs expressed high levels of P2X7, but skin MCs did not

Figure 5 | Inhibitory targeting of P2X7 purinoceptors on MCs leads to amelioration of colonic inflammation. *Kit*^{W-sh/W-sh} MC-deficient mice reconstituted with *P2x7*^{+/+}, *P2x7*^{-/-} or *caspase-1*^{-/-} BM-derived MCs were applied to a TNBS-induced colitis model. **(a)** Body weight changes were monitored in TNBS-treated *Kit*^{W-sh/W-sh} mice reconstituted with *P2x7*^{+/+} (closed squares; *n* = 9), *P2x7*^{-/-} (open squares; *n* = 7) or *caspase-1*^{-/-} (open triangles; *n* = 4). BM-derived MCs were used for TNBS treatment, and *P2x7*^{+/+} BM-derived MC-reconstituted *Kit*^{W-sh/W-sh} mice receiving EtOH served as controls (closed triangles; *n* = 3). **P* = 0.0264 (two-tailed Student's *t*-test), ***P* = 0.0058 (two-tailed Student's *t*-test). Data are shown as percentages of baseline weights and are means ± s.e.m. **(b)** Representative images of haematoxylin and eosin staining are shown. Scale bars represent 100 μm. **(c)** Representative images of whole colons are shown. **(d)** Colon length was measured 4 days after TNBS administration. Data are shown as means ± s.e.m. (*n* = 3 for *P2x7*^{+/+} EtOH, *n* = 9 for *P2x7*^{+/+} TNBS, *n* = 7 for *P2x7*^{-/-} TNBS). **P* < 0.001 (two-tailed Student's *t*-test). **(e)** Representative flow cytometric data of infiltrated neutrophils (CD11b⁺Gr-1^{high}) in the colon from three individual experiments. **P* = 0.00741, ***P* = 0.0009 (two-tailed Student's *t*-test). Data are shown as means ± s.e.m. **(f)** The percentage of CD63⁺ MCs in all c-kit⁺ FcεRIα⁺ MCs was determined with flow cytometry. Data are shown as means ± s.e.m. (*n* = 3–9). **P* = 0.007 (Welch's *t*-test), ***P* = 0.0234 (Welch's *t*-test). **(g)** Colonic tissue sections from a healthy volunteer (HV) and from UC and CD patients were stained with 4',6-diamidino-2-phenyl indole (blue), MC tryptase (red) and P2X7 (green). Scale bars, 100 μm. **(h)** Cells expressing both P2X7 and MC tryptase were counted in the fields of the tissue sections (four fields for each section). Data are means ± s.e.m. (*n* = 6).

(Fig. 3a). Second, in contrast to MCs, some macrophages (for example, microglia and RAW264.7 cells) expressed higher levels of P2X7 than did colonic macrophages (Fig. 3b and data not shown). Third, among the several types of immunocompetent cell in the colon, MCs expressed the highest levels of P2X7 (Fig. 3a,b). Fourth, we found P2X7 expression on human colonic ECs, but not on murine colonic ECs (Figs 3b and 5g). In addition, as reported previously³⁶, P2X7 expression on ECs was downregulated in the colons of CD patients; instead, CD patients showed increased numbers of P2X7⁺ MCs in their colons (Fig. 5g,h). It is important to note that, like murine MCs, human lung MCs express functional P2X7 (ref. 39). Therefore, although we must recognize the similarities and differences between mouse and human intestinal inflammation and MC distribution, ATP/P2X7-mediated MC activation seems to have a major role in the development of intestinal inflammation.

We found elevated levels of extracellular ATP in the colons of TNBS-treated mice (Fig. 6a). This high ATP concentration was most likely achieved by a combination or cascade of several ATP production pathways (for example, cell injury or lysis⁷, pattern recognition receptor-mediated activation of monocytes⁴⁰ and commensal bacteria³⁸). In our tissue culture system, we detected elevated release of ATP (40 μM) in the inflamed colon compared with the control (Fig. 6); however, at least 100 μM ATP was required for MC activation

in vitro in the single cell culture system (Fig. 7b). This disparity likely reflects the differences in the culture conditions. Unlike in the single cell culture system, the concentration of secreted ATP in the tissue culture system could have been diluted in the culture medium, or ATP could have been consumed rapidly by activated inflammatory cells in the tissue. Alternatively, a lack of commensal bacteria-derived ATP in the tissue culture system as a result of the inclusion of antibiotics may have reduced the ATP level. Another possibility is that the abundant endogenous ATP-degrading enzymes (for example, CD39) in the colon tissue may have degraded some of the ATP. In support of this idea, a suppressive role for CD39 in intestinal inflammation has been reported⁴¹.

We found that ADP-reactive P2Y1 and P2Y12 receptors were expressed on colonic MCs (Fig. 7c), but inhibition or knockdown of these receptors did not suppress the CD63 expression (Fig. 7d,e; Supplementary Fig. S8a). In previous studies, stimulation of MCs with ADP (0.05–50 μM) has led to calcium influx via the P2Y1- but not the P2Y12-mediated pathway⁴², whereas our results indicate that CD63 expression required a higher concentration of ADP and was not suppressed by a P2Y1 inhibitor (Fig. 7b,d). This finding indicates that P2Y purinoceptors are not involved in the induction of CD63⁺-activated MCs that is mediated by high concentrations of ADP. However, we found that adenylate kinase and ATP synthase converted ADP back to ATP, which subsequently induced P2X7

Figure 6 | Enhanced ATP production in intestinal inflammation and MC activation induced by non-hydrolyzable ATP. (a) The concentration of ATP released from the colon tissue of mice receiving intrarectally administered phosphate-buffered saline, vehicle or TNBS was measured. Data are shown as means $\pm$ s.e.m. ($n = 3-7$). $*P = 0.0004$, $**P = 0.0447$ (two-tailed Student's t -test). **(b)** CD63 expression of colonic MCs was measured with flow cytometry after intrarectal administration of vehicle ($n = 14$), TNBS ($n = 5$), non-hydrolyzable ATP (adenosine 5'-O-(3-thio) triphosphate (ATP γ S); $n = 9$ or O-(4-benzoyl)benzoyl adenosine 5'-triphosphate (BzATP); $n = 10$) or UDP-glucose ($n = 6$), or in intact mice ($n = 7$). Data are shown as means $\pm$ s.e.m. $*P = 0.0002$ (two-tailed Student's t -test), $**P = 0.0135$ (Welch's t -test) and $***P = 0.0238$ (Welch's t -test). NS, not significant.

purinoceptor-dependent MC activation. A similar conversion of ADP to ATP has been reported for endothelial cells²⁷. Among adenylate kinases, AK2 was highly expressed on MCs and had a pivotal role in the conversion of ADP to ATP (Supplementary Fig. S9a,b). As another P2Y ligand (UTP) did not induce MC activation (Fig. 7b), our findings suggest that ADP could be converted into ATP by AK2 and ATP synthase, and that this ATP subsequently activates MCs through P2X7 purinoceptors. In addition, colonic MCs do not express ecto-5'-nucleotidase (CD73), an enzyme that degrades ADP into adenosine, which has anti-inflammatory effects in intestinal inflammation⁴³. Therefore, our study indicates that MCs express CD39, adenylate kinases and ATP synthase, but not CD73, to preferentially convert ADP to ATP for the exacerbation of inflammatory responses through P2X7 purinoceptors.

Here, we showed that colitis aggravated by P2X7-mediated activation of MCs was independent of the inflammasome pathway, and that P2X7-mediated activation of MCs promoted TNF α production by effector cells to further promote intestinal inflammation⁴⁴. Our findings also suggest that MCs exacerbate inflammation by recruiting neutrophils to produce abundant TNF α , but less IL-10 than is produced by other cells (for example, eosinophils, DCs and macrophages; Supplementary Fig. S10d). This neutrophil recruitment was mediated by the production of IL-1 β , LTs and chemokines, which are potential targets for the treatment of colitis. Mice with experimentally induced colitis that lack CXCR2 or 5-LO (a key enzyme for converting arachidonic acid to LTs), as well as mice treated with inhibitors of CCR2, CXCR2 or 5-LO, show reduced inflammation and less neutrophil recruitment in their colons^{33,45,46}. Moreover, given that ATP promotes neutrophil migration⁴⁷, it is possible that P2X7-dependent LT and chemokine production, as well as ATP generation via AK2 and ATP synthase from MCs, could amplify neutrophil infiltration of the colon. These data collectively indicate that MCs are key factors in the induction of intestinal inflammation and also recruit neutrophils to heighten inflammatory responses. P2X7-dependent MC activation could, therefore, be a target for the treatment of intestinal inflammation.

Methods

Mice and human samples. Female C57BL/6 mice were purchased from CLEA Japan. Rag1^{-/-} and P2x7^{-/-} mice were obtained from Jackson Laboratory (Bar Harbor, ME, USA). MC-deficient Kit^{W-sh/W-sh} mice were obtained from Dr H. Suto (Atopy Research Center, Juntendo University, Japan). For the conditional MC-deficient analysis, Mas-TRECK tg mice were injected intraperitoneally with 250 ng of diphtheria toxin for 5 consecutive days and then with 150 ng every other day¹⁸. *Caspase-1*^{-/-} mice were backcrossed with C57BL/6 mice; F5 mice were used for this experiment⁴⁸. All mice were maintained under specific-pathogen-free conditions at the Experimental Animal Facility of the Institute of Medical Science, the University of Tokyo. All experiments were approved by the Animal Care and Use Committee of the University of Tokyo.

MC reconstitution was performed as described previously⁴⁹. Briefly, BM-derived MCs were obtained from P2x7^{+/+}, P2x7^{-/-} or *caspase-1*^{-/-} mice as described previously²². BM-derived MCs (5×10^6) were intravenously transferred to Kit^{W-sh/W-sh} mice at two time points (0 and 14 days). The reconstituted mice were used 3 months after the last transfer.

Colon specimens from UC and CD patients and healthy volunteers were obtained by endoscopic biopsy at Osaka University Hospital. All subjects provided written informed consent, and the study protocol was approved by the Ethics Committee of Osaka University Graduate School of Medicine (no. 08243) and the Institute of Medical Science, The University of Tokyo (no. 20-67-0331).

Experimental colitis. For TNBS-induced colitis, anaesthetized mice (18–22 g) were sensitized with 2.5% TNBS (Sigma-Aldrich) together with acetone and olive oil⁵⁰. After 1 week, after a 3-h fast, the mice were given 100 μ l of 2.5% TNBS in 50% ethanol via a flexible feeding tube that maintained their heads in a vertical position for 10 min. The control group received only 50% ethanol. Weight changes were recorded daily, and tissues were collected for histological analysis and isolation of mononuclear cells from the colonic lamina propria. For mAb treatment, mice were injected intraperitoneally with 0.5 mg of mAb (1F11 or an isotype control) 1 day before being given TNBS/EtOH intrarectally. mAb administration was continued for 3 days. For P2Y12 inhibition with clopidogrel sulphate, (Wako, Osaka, Japan), mice received clopidogrel (0.5 mg ml⁻¹) in their drinking water from 3 days before intrarectal administration of TNBS/EtOH until the end of the study⁵⁰. For DSS-induced colitis, mice were given 3.5% DSS (Wako, for C57BL/6) or 2.5% DSS (MP Biomedicals, Illkirch, France, for Mas-TRECK tg mice) in their drinking water for 5 days and their body weights were monitored daily⁵⁰. In some experiments, non-hydrolysable ATP (adenosine 5'-O-(3-thio) triphosphate and O-(4-benzoyl)benzoyl adenosine 5'-triphosphate) or UDP-glucose (0.25 mg in 50% EtOH) was intrarectally administered and the effects were analysed 2 days later.

In vitro MC stimulation and inhibition. BM-derived MCs (2.5×10^5) were cultured with various concentrations of adenosine, ADP, ATP, UTP or anti-DNP-IgE with DNP-human serum albumin. Adenosine-3-phosphate 5-phosphosulfate (0.25 mM), carbenoxolone (10 μ M), flufenamic acid (100 μ M), pyridoxal-phosphate-6-azophenyl-2',4'-disulfonate (100 μ M), 4,4'-diisothiocyanatostilbene-2,2'-disulfonic acid disodium salt hydrate (100 μ M), O α ATP (0.5 mM), AD2P5 (1 mM), oligo (10 or 100 μ M) or UDP (100 μ M) was added to the cells for the inhibition assay^{27,28,40,51}. All reagents were purchased from Sigma-Aldrich (St Louis, MO, USA, purity was $\geq 95\%$), 5-LO (BD Pharmingen, Franklin Lakes, NJ, USA) was stained after permeabilization with 0.2% Triton-X100 for 10 min; nuclei were stained with 4',6-diamidino-2-phenyl indole.

Cell preparation and flow cytometry. ECs and lamina propria mononuclear cells were isolated from the colon, as described previously⁵². For flow cytometric analysis, cells were incubated with 5 μ g ml⁻¹ of an anti-CD16/32 antibody (10 μ g ml⁻¹, Fc block, BD Pharmingen) for 5 min and stained for 30 min at 4°C with fluorescence-labeled Abs specific for c-kit (0.2 μ g ml⁻¹), Gr-1 (0.4 μ g ml⁻¹), CD4 (1 μ g ml⁻¹), CD11b (0.2 μ g ml⁻¹), CD11c (0.4 μ g ml⁻¹), CD39 (0.4 μ g ml⁻¹), CD45 (0.4 μ g ml⁻¹), IgA (10 μ g ml⁻¹), B220 (0.4 μ g ml⁻¹; BD Pharmingen), CCR3 (2 μ g ml⁻¹), CXCR2 (4 μ g ml⁻¹; R&D Systems, Minneapolis, MN, USA), Fc ϵ RI α (0.4 μ g ml⁻¹), CD73 (0.4 μ g ml⁻¹), TLR2 (10 μ g ml⁻¹; eBioscience, San Diego, CA, USA), F4/80 (20 μ g ml⁻¹), CCR2 (10 μ g ml⁻¹), P2X7 (Hano43; 2 μ g ml⁻¹, Serotec, UK) or CCR1 (10 μ g ml⁻¹, Abnova, Taiwan). Flow cytometric analysis and cell sorting were performed by using FACSCalibur and FACSAria (BD Biosciences, Franklin Lakes, NJ, USA), respectively. Sorted cells were stained with May-Giemsa stain in some experiments. Colonic MCs and BM-derived MCs were prepared as described elsewhere²².

Establishment of an anti-P2X7 mAb (1F11) and an anti-CD63 mAb. The procedure used to establish MC-specific mAbs is shown as a flowchart in Supplementary Figure S3. Briefly, c-kit⁺ Fc ϵ RI α ⁺ MCs were obtained as described previously²² from the colons of mice that exhibited allergic diarrhoea. Purified colonic MCs (10^6 cells) were injected into the footpads of Sprague Dawley rats seven times, as described previously⁵³. Lymphocytes were isolated from the spleen and inguinal lymph nodes and fused with P3X63-AG8.653 myeloma cells (CRL-1580; American Type Culture Collection, Manassas, VA, USA). The reactivity of each hybridoma to the colonic MCs was examined by means of flow cytometry. To identify antigens

Figure 7 | The ecto-adenylate kinase pathway mediates ADP-dependent MC activation through P2X7 purinoceptors. (a) BM-derived MCs treated with ADP or ATP at 0.5 mM for 30 min and examined for CD63 expression. (b) BM-derived MCs treated with IgE plus relevant allergen or various concentrations of ADP, ATP or UTP for the analysis of CD63 expression. Data are representative of four experiments. (c) Expression of mRNA encoding each P2Y receptor in colonic lamina propria lymphocytes (LPLs) and MCs was analysed by quantitative reverse transcription (RT)-PCR ($n=3$). (d,e) BM-derived MCs pre-treated with 0.25 mM P2Y1 inhibitor (adenosine-3-phosphate 5-phosphosulfate (A3P5PS)) (d) or 0.01 mM P2Y12 inhibitor (MRS2353) (e), stimulated with ADP and examined for CD63 expression ($n=3$). (f) BM-derived MCs from $P2x7^{-/-}$ mice stimulated with ATP or ADP; CD63 expression was determined with flow cytometry. Data are representative of four experiments. (g) Expression of pannexin-1 (Panx1), connexin-43 (Connx43) and Connx32 on colonic MCs, macrophages (Mac), CD4⁺ T cells (CD4), DCs, IgA⁺ cells (IgA) and ECs was measured by quantitative RT-PCR ($n=4$). (h) BM-derived MCs were pretreated with inhibitors of P2X receptors [OxATP, 0.5 mM; pyridoxal-phosphate-6-azophenyl-2',4'-disulfonate (PPADS); 4,4'-diisothiocyanatostilbene-2,2'-disulfonic acid (DIDS)], connexins [flufenamic acid (FFA)], Panx-1 [carbenoxolone (CBX)], ecto-adenylate kinase [diadenosine pentaphosphate (AD2P5)], ATP synthase (oligomycin) or nucleoside diphosphokinase (UDP) and subsequently stimulated with 0.25 mM ADP. CD63 expression was determined with flow cytometry. ($n=3$) $*P<0.01$, $**P<0.05$ (two-tailed Student's t -test). All data are shown as means $\pm$ s.e.m. (i) BM-derived MCs were treated with AD2P5, oligomycin or UDP and stimulated with 0.5 mM ATP or IgE plus allergen. CD63 expression was determined with flow cytometry ($n=5$). $*P<0.0001$ (two-tailed Student's t -test), $**P=0.0008$ (two-tailed Student's t -test) and $***P=0.0008$ (Welch's t -test). NS, not significant.

recognized by the mAbs, immunoprecipitation was performed with the mAbs, followed by Liquid chromatography-mass spectrometry analysis, as described previously⁵³. Antigen specificity was confirmed by transfecting CHO cells with plasmids that encoded the murine P2X7 receptor and CD63.

Measurements of membrane permeability and inflammatory mediators.

To assess membrane permeability, BM-derived MCs were washed twice with phosphate-buffered saline (PBS) and incubated with 1 mg ml⁻¹ Lucifer yellow (Sigma-Aldrich) containing 250 μ M sulfapyrazole (Sigma-Aldrich). The MCs were then stimulated with 0.5 mM ATP (Sigma-Aldrich) for 15 min, as described elsewhere¹². In the inhibition assay, 1 or 10 μ g ml⁻¹ of 1F11 mAb or the control antibody (Rat IgG2b) was added before ATP stimulation. The fluorescence signal of Lucifer yellow was determined by using fluorescence microscopy (BZ9000, Keyence, Osaka, Japan) and flow cytometry.

To measure the production of cytokines, chemokines and LTs from MCs, BM-derived MCs (2.5×10^5) were stimulated with 2.5 mM ATP for 30 min, after which the supernatants were collected. Chemokine and cytokine production was detected with an inflammatory cytokine kit (BD Pharmingen). For IL-1 β measurement, BM-derived MCs from wild-type, $P2x7^{-/-}$ and $caspase-1^{-/-}$ mice

were stimulated with 0.1 μ g ml⁻¹ of LPS for 4 h, followed by ADP or ATP stimulation. LT C4/D4/E4 production was detected by use of an enzyme-linked immunosorbent assay (GE Healthcare Bio-Science, NJ, USA). For ATP, cytokine and chemokine measurements from the colon tissue, the colon tissues were isolated from mice 2 days after intrarectal administration of TNBS. Released ATP was measured by culturing colon tissues at 100 mg of tissue per 100 μ l of RPMI1640 medium for 3 h and using a luminescence ATP detection system (PerkinElmer, Norwalk, CT, USA).

Immunoprecipitation and western blotting. Cell lysates obtained from BM-derived MCs or CHO transfectants (mouse P2X7 variants a, c and d and flag-mP2X7s, cloned from C57BL/6 mice) were analysed by western blotting and immunoprecipitation with 1F11 mAb or the control Ab. Membranes were probed with an anti-flag and a polyclonal rabbit anti-P2X7 antibody (Sigma-Aldrich).

Histology. Colonic tissues were fixed in 4% paraformaldehyde and embedded in paraffin. Tissue sections (5 μ m) were stained with haematoxylin and eosin solution, as described previously²². For the detection of MCs and P2X7

Figure 8 | Critical role of the intestinal MC-associated ATP-P2X7 purinoceptor pathway for induction of neutrophil infiltration. *P2x7*^{+/+} and *P2x7*^{-/-} BM-derived MCs were treated with 0.25 mM ATP (+) or left untreated (-). **(a)** Production of IL-6 (left panel; isotype mAb-treated MC, closed column; 1F11 mAb-treated MC, open column; *P2x7*^{+/+}, grey column; and *P2x7*^{-/-}, beige column) and TNFα (right panel) in culture supernatant (*P2x7*^{+/+}, closed column; *P2x7*^{-/-}, open column) was determined after 24 h stimulation. ND, not detected. Data are shown as means ± s.e.m. (*n* = 3). **(b)** Oncostatin M mRNA expression was measured 30 min after stimulation of *P2x7*^{+/+} (closed column) and *P2x7*^{-/-} (open column) MCs with ATP. Data are shown as means ± s.e.m. (*n* = 3). **(c)** LT C4/D4/E4 production from ATP-stimulated (+) or -unstimulated (-) *P2x7*^{+/+} (closed column) or *P2x7*^{-/-} BM-derived MCs (open column) in culture supernatants was measured by using enzyme-linked immunosorbent assay (ELISA). Data are shown as means ± s.e.m. (*n* = 3). ND, not detected. **(d)** *P2x7*^{+/+} and *P2x7*^{-/-} BM-derived MCs were stimulated with 0.5 mM ATP. Cells were fixed and stained with an anti-5LO antibody (red) and 4',6-diamidino-2-phenyl indole (blue). Scale bar, 10 μm. Data are representative of two experiments. **(e)** Representative data of a chemokine gene array are shown. Increased levels of each chemokine are shown as a heat map. **(f)** mRNA expression of CCL2 (left), CCL7 (middle) and CXCL2 (right) was measured by using quantitative reverse transcription-PCR. Data are shown as means ± s.e.m. (*n* = 3). **(g)** CCL2 production was enumerated by using ELISA 24 h after stimulation of BM-derived MCs with ATP or IgE plus antigen (IgE + Ag). Isotype mAb-treated MC, closed column; 1F11 mAb-treated MC, open column; *P2x7*^{+/+} MC, grey column; and *P2x7*^{-/-} MC, beige column). Data are shown as means ± s.e.m. (*n* = 3).

expression in human specimens, colonic tissue sections were stained with antibodies for MC tryptase and P2X7 purinoceptors (Alomone Laboratories, Jerusalem, Israel).

shRNA plasmid construction and lentiviral transduction. For the construction of shRNA expression lentivirus vector plasmids, a series of oligonucleotide pairs were synthesized, as listed below. Each oligo pair was annealed and cloned into pmU6⁵⁴. Each mU6-shRNA cassette was then subcloned into the ΔU3 sequence of the 3'-LTR of the lentivirus vector plasmid pLCG to generate pLCG-shCD63 (sense: 5'-TTTGATCTCTGTGCATCAACATAGCTTCCTGTCACATGTTGATGTCAGCAAGAATCTTTTGG-3', antisense: 5'-AATTCAAAAAAGATCTTGTGCTGCAACAACATAGTCAGGAAGCTATGTTGATGTCAGCAAGAAT-3'), pLCG-shP2Y12 (sense: 5'-TTTGATCTACTAATGATTTCAACTGCTTCCTGTACAGTTAGAAATCATCTAGATCTTTTGG-3', antisense: 5'-AATTCAAAAAAGATCTACTAA TGATTCTAACTGTGACAGGAAGCAGTTAGAATCATTAGTAGAT-3') and pLCG-shAK1 (sense: 5'-TTTGCGAGAAGATGTACAGAAATGCTTCCTGTGCA CATTCTGTACAATCTTCTCGCTTTTGG-3', antisense: 5'-AATTCAAAAAA GCGAGAAGATGTGACAGAAATGTGACAGGAAGCATTCTGTACAATCTT CTG-3') and pLCG-shAK2 (sense: 5'-TTTGGAGCTAATTGAGAAGAATTGC TTCTGTGACAAATCTTCTCAATTAGCTCCATTTTGG-3', antisense: 5'-AATTCAAAAAATGGAGCTAATTGAGAAGAATTGTGACAGGAAGCA ATTCTTCTCAATTAGCTCC-3').

To obtain lentivirus-encoding green fluorescent protein (as a reporter gene) and shRNA for CD63, 293FT cells (6 × 10⁵) were transfected with pLP1, pLP2, pLP-VSVG and pLCG-shRNA by using Lipofectamine 2000 (Invitrogen, Carlsbad, CA, USA) as per the manufacturer's protocol (Invitrogen). After 24- and 48-h incubations, lentivirus-encoding shRNA was collected.

BM-derived MCs (1 × 10⁶) or MC/9 cells were transduced with shRNA expression lentivirus stock in the presence of 8 μg ml⁻¹ Polybrene (Sigma-Aldrich)⁵⁵.

After 24 h, the cells were washed and green fluorescent protein-positive cells were sorted by FACS Aria and used for subsequent experiments.

Quantitative real-time-PCR. Total RNA was prepared by using TRIzol (Invitrogen) and reverse transcribed by use of Superscript VILO (Invitrogen), as described. Quantitative reverse transcription-PCR was performed with the LightCycler 480 II (Roche, Mannheim, Germany) and the Universal Probe Library (Roche). Primer sequences are listed in Supplementary Table S1.

Statistical analysis. Statistical analysis was performed by using the unpaired two-tailed Student's *t*-test and Welch's *t*-test. The data are presented as means ± s.e.m.

References

- Abraham, C. & Cho, J. H. Inflammatory bowel disease. *N Engl. J. Med.* **361**, 2066–2078 (2009).
- Strober, W. & Fuss, I. J. Proinflammatory cytokines in the pathogenesis of inflammatory bowel diseases. *Gastroenterology* **140**, 1756–1767 (2011).
- Alvarez-Errico, D., Lessmann, E. & Rivera, J. Adapters in the organization of mast cell signaling. *Immunol. Rev.* **232**, 195–217 (2009).
- Bischoff, S. C. Role of mast cells in allergic and non-allergic immune responses: comparison of human and murine data. *Nat. Rev. Immunol.* **7**, 93–104 (2007).
- Galli, S. J., Borregaard, N. & Wynn, T. A. Phenotypic and functional plasticity of cells of innate immunity: macrophages, mast cells and neutrophils. *Nat. Immunol.* **12**, 1035–1044 (2011).
- He, S. H. Key role of mast cells and their major secretory products in inflammatory bowel disease. *World J. Gastroenterol.* **10**, 309–318 (2004).
- Surprenant, A. & North, R. A. Signaling at purinergic P2X receptors. *Annu. Rev. Physiol.* **71**, 333–359 (2009).

8. Erlinge, D. P2Y receptors in health and disease. *Adv. Pharmacol.* **61**, 417–439 (2011).
9. Wilhelm, K. *et al.* Graft-versus-host disease is enhanced by extracellular ATP activating P2X7R. *Nat. Med.* **16**, 1434–1438 (2010).
10. Weber, F. C. *et al.* Lack of the purinergic receptor P2X(7) results in resistance to contact hypersensitivity. *J. Exp. Med.* **207**, 2609–2619 (2010).
11. Muller, T. *et al.* A potential role for P2X7R in allergic airway inflammation in mice and humans. *Am. J. Respir. Cell Mol. Biol.* **44**, 456–464 (2011).
12. Sudo, N. *et al.* Extracellular ATP activates mast cells via a mechanism that is different from the activation induced by the cross-linking of Fc receptors. *J. Immunol.* **156**, 3970–3979 (1996).
13. Furuno, T., Teshima, R., Kitani, S., Sawada, J. & Nakanishi, M. Surface expression of CD63 antigen (AD1 antigen) in P815 mastocytoma cells by transfected IgE receptors. *Biochem. Biophys. Res. Commun.* **219**, 740–744 (1996).
14. Rijnsierse, A., Koster, A. S., Nijkamp, F. P. & Kraneveld, A. D. Critical role for mast cells in the pathogenesis of 2,4-dinitrobenzene-induced murine colonic hypersensitivity reaction. *J. Immunol.* **176**, 4375–4384 (2006).
15. Kaser, A., Zeissig, S. & Blumberg, R. S. Inflammatory bowel disease. *Annu. Rev. Immunol.* **28**, 573–621 (2010).
16. Feyerabend, T. B. *et al.* Cre-mediated cell ablation contests mast cell contribution in models of antibody- and T cell-mediated autoimmunity. *Immunity* **35**, 832–844 (2011).
17. Otsuka, A. *et al.* Requirement of interaction between mast cells and skin dendritic cells to establish contact hypersensitivity. *PLoS One* **6**, e25538 (2011).
18. Sawaguchi, M. *et al.* Role of mast cells and basophils in IgE responses and in allergic airway hyperresponsiveness. *J. Immunol.* **188**, 1809–1818 (2012).
19. Fiorucci, S. *et al.* Importance of innate immunity and collagen binding integrin $\alpha 1\beta 1$ in TNBS-induced colitis. *Immunity* **17**, 769–780 (2002).
20. Yoshimoto, T. & Nakanishi, K. Roles of IL-18 in basophils and mast cells. *Allergol. Int.* **55**, 105–113 (2006).
21. Pastorelli, L. *et al.* Epithelial-derived IL-33 and its receptor ST2 are dysregulated in ulcerative colitis and in experimental Th1/Th2 driven enteritis. *Proc. Natl Acad. Sci. USA* **107**, 8017–8022 (2010).
22. Kurashima, Y. *et al.* Sphingosine 1-phosphate-mediated trafficking of pathogenic Th2 and mast cells for the control of food allergy. *J. Immunol.* **179**, 1577–1585 (2007).
23. Nicke, A. *et al.* A functional P2X7 splice variant with an alternative transmembrane domain 1 escapes gene inactivation in P2X7 knock-out mice. *J. Biol. Chem.* **284**, 25813–25822 (2009).
24. Smart, M. L. *et al.* P2X7 receptor cell surface expression and cytolytic pore formation are regulated by a distal C-terminal region. *J. Biol. Chem.* **278**, 8853–8860 (2003).
25. Di Virgilio, F. Liaisons dangereuses: P2X₇ and the inflammasome. *Trends Pharmacol. Sci.* **28**, 465–472 (2007).
26. Mizutani, H., Schechter, N., Lazarus, G., Black, R. A. & Kupper, T. S. Rapid and specific conversion of precursor interleukin 1 β (IL-1 β) to an active IL-1 species by human mast cell chymase. *J. Exp. Med.* **174**, 821–825 (1991).
27. Yegutkin, G. G. Nucleotide- and nucleoside-converting ectoenzymes: important modulators of purinergic signalling cascade. *Biochim. Biophys. Acta.* **1783**, 673–694 (2008).
28. Locovei, S., Bao, L. & Dahl, G. Pannexin 1 in erythrocytes: function without a gap. *Proc. Natl Acad. Sci. USA* **103**, 7655–7659 (2006).
29. Kang, J. *et al.* Connexin 43 hemichannels are permeable to ATP. *J. Neurosci.* **28**, 4702–4711 (2008).
30. Burrell, H. E. *et al.* Human keratinocytes release ATP and utilize three mechanisms for nucleotide interconversion at the cell surface. *J. Biol. Chem.* **280**, 29667–29676 (2005).
31. Foger, N. *et al.* Differential regulation of mast cell degranulation versus cytokine secretion by the actin regulatory proteins Coronin1a and Coronin1b. *J. Exp. Med.* **208**, 1777–1787 (2011).
32. Salamon, P. *et al.* Human mast cells release oncostatin M on contact with activated T cells: possible biologic relevance. *J. Allergy Clin. Immunol.* **121**, 448–455 e5 (2008).
33. Bento, A. F. *et al.* The selective nonpeptide CXCR2 antagonist SB225002 ameliorates acute experimental colitis in mice. *J. Leukoc. Biol.* **84**, 1213–1221 (2008).
34. Luo, M., Jones, S. M., Peters-Golden, M. & Brock, T. G. Nuclear localization of 5-lipoxygenase as a determinant of leukotriene B₄ synthetic capacity. *Proc. Natl Acad. Sci. USA* **100**, 12165–12170 (2003).
35. Nigrovic, P. A. *et al.* Genetic inversion in mast cell-deficient *Wsh* mice interrupts corin and manifests as hematopoietic and cardiac aberrancy. *Am. J. Pathol.* **173**, 1693–1701 (2008).
36. Cesaro, A. *et al.* Amplification loop of the inflammatory process is induced by P2X7R activation in intestinal epithelial cells in response to neutrophil trans epithelial migration. *Am. J. Physiol. Gastrointest Liver Physiol.* **299**, G32–G42 (2010).
37. Schenk, U. *et al.* ATP inhibits the generation and function of regulatory T cells through the activation of purinergic P2X receptors. *Sci. Signal* **4**, ra12 (2011).
38. Atarashi, K. *et al.* ATP drives lamina propria T_H17 cell differentiation. *Nature* **455**, 808–812 (2008).
39. Wareham, K., Vial, C., Wykes, R. C., Bradding, P. & Seward, E. P. Functional evidence for the expression of P2X₁, P2X₄ and P2X₇ receptors in human lung mast cells. *Br. J. Pharmacol.* **157**, 1215–1224 (2009).
40. Piccini, A. *et al.* ATP is released by monocytes stimulated with pathogen-sensing receptor ligands and induces IL-1 β and IL-18 secretion in an autocrine way. *Proc. Natl Acad. Sci. USA* **105**, 8067–8072 (2008).
41. Friedman, D. J. *et al.* From the Cover: CD39 deletion exacerbates experimental murine colitis and human polymorphisms increase susceptibility to inflammatory bowel disease. *Proc. Natl Acad. Sci. USA* **106**, 16788–16793 (2009).
42. Feng, C., Mery, A. G., Beller, E. M., Favot, C. & Boyce, J. A. Adenine nucleotides inhibit cytokine generation by human mast cells through a Gs-coupled receptor. *J. Immunol.* **173**, 7539–7547 (2004).
43. Louis, N. A. *et al.* Control of IFN- α A by CD73: implications for mucosal inflammation. *J. Immunol.* **180**, 4246–4255 (2008).
44. Rijnsierse, A., Koster, A. S., Nijkamp, F. P. & Kraneveld, A. D. TNF- α is crucial for the development of mast cell-dependent colitis in mice. *Am. J. Physiol. Gastrointest. Liver Physiol.* **291**, G969–G976 (2006).
45. Cuzzocrea, S. *et al.* 5-Lipoxygenase modulates colitis through the regulation of adhesion molecule expression and neutrophil migration. *Lab. Invest.* **85**, 808–822 (2005).
46. Waddell, A. *et al.* Colonic eosinophilic inflammation in experimental colitis is mediated by Ly6C^{high} CCR2⁺ inflammatory monocyte/macrophage-derived CCL11. *J. Immunol.* **186**, 5993–6003 (2011).
47. Chen, Y. *et al.* Purinergic signaling: a fundamental mechanism in neutrophil activation. *Sci. Signal* **3**, ra45 (2010).
48. Kuida, K. *et al.* Altered cytokine export and apoptosis in mice deficient in interleukin-1 β converting enzyme. *Science* **267**, 2000–2003 (1995).
49. Tsai, M., Grimbaldston, M. A., Yu, M., Tam, S. Y. & Galli, S. J. Using mast cell knock-in mice to analyze the roles of mast cells in allergic responses *in vivo*. *Chem. Immunol. Allergy* **87**, 179–197 (2005).
50. Wirtz, S., Neufert, C., Weigmann, B. & Neurath, M. F. Chemically induced mouse models of intestinal inflammation. *Nat. Protoc.* **2**, 541–546 (2007).
51. Ransford, G. A. *et al.* Pannexin 1 contributes to ATP release in airway epithelia. *Am. J. Respir. Cell Mol. Biol.* **41**, 525–534 (2009).
52. Kunisawa, J. *et al.* Sphingosine 1-phosphate dependence in the regulation of lymphocyte trafficking to the gut epithelium. *J. Exp. Med.* **204**, 2335–2348 (2007).
53. Nochi, T. *et al.* A novel M cell-specific carbohydrate-targeted mucosal vaccine effectively induces antigen-specific immune responses. *J. Exp. Med.* **204**, 2789–2796 (2007).
54. Yu, J. Y., DeRuiter, S. L. & Turner, D. L. RNA interference by expression of short-interfering RNAs and hairpin RNAs in mammalian cells. *Proc. Natl Acad. Sci. USA* **99**, 6047–6052 (2002).
55. Haraguchi, T., Ozaki, Y. & Iba, H. Vectors expressing efficient RNA decoys achieve the long-term suppression of specific microRNA activity in mammalian cells. *Nucleic Acids Res.* **37**, e43 (2009).

Acknowledgements

We thank Drs T. Kitamura and J. Kitaura (The University of Tokyo) for discussions and for providing reagents; Dr H. Suto (Juntendo University) for providing *Kit^{W-sh/W-sh}* mice; Dr S. Nakae (The University of Tokyo) for advice on analysing *Kit^{W-sh/W-sh}* mice; Drs S. Ohmi, M. Oyama, H. Hata and C. Takamura (The University of Tokyo) for protein analysis; and Dr A. Uozumi, I. Ishikawa and M. Mejima (The University of Tokyo), Drs F. Ishikawa and Y. Saito (RCAI) for technical advice. This work was supported by grants from: the Ministry of Education, Science, Sports and Technology of Japan (Grant-in Aid for Scientific Research S (23229004 to H.K.), for Young Scientists A (22689015 to J.K.), for Scientific Research on Innovative Areas (23116506 to J.K.), for Scientific Research on Priority Areas (19059003 to H.K.), for Challenging Exploratory Research (24659217 to J.K.), Leading-edge Research Infrastructure Program (J.K. and H.K.); the Young Researcher Overseas Visits Program for Vitalizing Brain Circulation of the Japan Society for the Promotion of Science (JSPS; J.K., H.K., Y.K., T.N.), and for JSPS Fellows (021-07124 to Y.K.); the Ministry of Health and Welfare of Japan (J.K. and H.K.); the Global Center of Excellence Program of the Center of Education and Research for Advanced Genome-based Medicine (H.K.); the Program for Promotion of Basic and Applied Research for Innovations in Bio-oriented Industry (BRAIN to J.K.); and the Yakult Bio-Science Foundation (J.K.).

Author contributions

Y.K. conducted the research, performed experiments and wrote the manuscript; T.A. and K.F. performed gene expression and animal experiments; T.N. conducted the mAb experiment; H.T., H. Iba, T.H., M.K. and S.S. contributed to the experimental design and data analysis; S.N. and H. Iijima obtained clinical samples and J.K. and H.K. supervised the project and wrote the manuscript. JK should be contacted for material requests.

Additional information

Supplementary Information accompanies this paper at <http://www.nature.com/naturecommunications>

Competing financial interests: The authors declare no competing financial interests.

Reprints and permission information is available online at <http://npg.nature.com/reprintsandpermissions/>

How to cite this article: Kurashima, Y. *et al.* Extracellular ATP mediates mast cell-dependent intestinal inflammation through P2X7 purinoceptors. *Nat. Commun.* 3:1034 doi: 10.1038/ncomms2023 (2012).

License: This work is licensed under a Creative Commons Attribution-NonCommercial-Share Alike 3.0 Unported License. To view a copy of this license, visit <http://creativecommons.org/licenses/by-nc-sa/3.0/>