

ARTICLE

Received 20 Jun 2012 | Accepted 17 Jan 2013 | Published 26 Feb 2013

DOI: 10.1038/ncomms2512

OPEN

A new class of plant lipid is essential for protection against phosphorus depletion

Yozo Okazaki¹, Hitomi Otsuki¹, Tomoko Narisawa¹, Makoto Kobayashi¹, Satoru Sawai^{2,†}, Yukiko Kamide¹, Miyako Kusano¹, Toshio Aoki³, Masami Yokota Hirai^{1,4} & Kazuki Saito^{1,2}

Phosphorus supply is a major factor responsible for reduced crop yields. As a result, plants utilize various adaptive mechanisms against phosphorus depletion, including lipid remodeling. Here we report the involvement of a novel plant lipid, glucuronosyldiacylglycerol, against phosphorus depletion. Lipidomic analysis of *Arabidopsis* plants cultured in phosphorus-depleted conditions revealed inducible accumulation of glucuronosyldiacylglycerol. Investigation using a series of sulfolipid sulfoquinovosyldiacylglycerol synthesis-deficient mutants of *Arabidopsis* determined that the biosynthesis of glucuronosyldiacylglycerol shares the pathway of sulfoquinovosyldiacylglycerol synthesis in chloroplasts. Under phosphorus-depleted conditions, the *Arabidopsis sqd2* mutant, which does not accumulate either sulfoquinovosyldiacylglycerol or glucuronosyldiacylglycerol, was the most severely damaged of three sulfoquinovosyldiacylglycerol-deficient mutants. As glucuronosyldiacylglycerol is still present in the other two mutants, this result indicates that glucuronosyldiacylglycerol has a role in the protection of plants against phosphorus limitation stress. Glucuronosyldiacylglycerol was also found in rice, and its concentration increased significantly following phosphorus limitation, suggesting a shared physiological significance of this novel lipid against phosphorus depletion in plants.

¹Metabolomic Function Research Group, RIKEN Plant Science Center, Yokohama 230-0045, Japan. ²Graduate School of Pharmaceutical Sciences, Chiba University, Chiba 260-8675, Japan. ³Department of Applied Biological Sciences, Nihon- University, Fujisawa 252-0880, Japan. ⁴Japan Science and Technology Agency, Core Research for Evolutional Science and Technology, Kawaguchi 332-0012, Japan. †Present address: RIKEN Plant Science Center, Yokohama, Japan. Correspondence and requests for materials should be addressed to K.S. (email: ksaito@psc.riken.jp).

Phosphorus (P) is an essential component in cells and has a crucial role in cellular functionality and metabolism. However, because of limited P availability in many soils, plants often suffer from P deficiency¹. Indeed, it is estimated that crop yields for 40% of the world's arable land are reduced because of P unavailability². As current known P reserves are limited and P-based fertilizers are economically produced, efficient use and the minimization of the loss of P resources (that is, recycling of struvite, magnesium ammonium phosphate) are very important to sustainable agriculture^{3–5}. However, plants have developed several adaptation mechanisms to cope with P-limited environments.

P mobilization by membrane lipid remodelling, that is, changing the composition of the membrane lipids rich in phospholipids, is one of the general mechanisms of adapting to P limitation in plants^{6–9}. Replacement of membrane phospholipids with non-phosphorus glycerolipids such as sulfoquinovosyldiacylglycerol (SQDG) and digalactosyldiacylglycerol (DGDG), which promote the remobilization of P, is typical of metabolic signatures associated with lipid remodelling during phosphate deprivation. These glycolipids are usually found in chloroplast membranes, but DGDG is also found in extraplastidic membranes in P-starved plants^{7,8,10}. Thus far, various aspects of lipid remodelling following P limitation have been investigated. The biosynthetic enzymes involved in the enhanced production of these non-phosphorus glycolipids under P limitation, mainly glycosyltransferases and sugar-donor synthetic enzymes, have been intensely characterized^{11–18}. Phospholipid degradation and P recycling have also been analysed through the characterization of several phospholipases and glycerophosphodiesterases^{15,19–22}. In addition, parts of the signal transduction pathway for lipid remodelling have been studied^{14,15,23,24}. However, detailed metabolite analysis regarding lipid remodelling has been limited. Previous studies have thus far focused only on several well-known lipid classes²⁵ because of the difficulties associated with the comprehensive analysis of plant lipids, which consist of a wide variety of hydrophobic chemicals. In this study, we applied untargeted metabolomic analysis, which allows comprehensive chemical metabolic phenotyping of cells, to investigate lipid remodelling in detail and to identify a new key lipid molecule that has an essential role in protection against P-depletion stress.

Results

A new plant lipid inducibly accumulates under P depletion. To expose plants to P-limiting stress, wild-type *Arabidopsis thaliana* plants grown on phosphate (P_i)-sufficient growth medium for 14 days were transferred to either P_i -sufficient or P_i -depleted growth medium. After another 14 days of growth, lipids were extracted from the leaves and subjected to lipidomic analysis by hydrophilic interaction chromatography (HILIC) coupled with ion trap time-of-flight mass spectrometry²⁶. This analysis indicated that P limitation induced a decrease in phospholipids such as phosphatidylglycerol (PG), phosphatidylethanolamine (PE), phosphatidylinositol (PI) and phosphatidylcholine, with a complementary increase in non-phosphorus glycerolipids, including SQDG (~4.5-fold) and DGDG (~2-fold), as previously reported (Fig. 1a)^{7,12}. The total monogalactosyldiacylglycerol (MGDG) level did not significantly change in response to P limitation, as described below (Fig. 1a). In addition to these changes, a new peak (UK1; retention time, 11.4 min) was found to appear under P limitation but not under P sufficiency. To obtain detailed information regarding the overall metabolic changes induced by P limitation, an orthogonal projection to a latent structure-discriminant analysis (OPLS-DA) model was examined using the lipidomic data obtained from the plants subjected to P-limited and P-sufficient conditions²⁷. The OPLS-DA (Supplementary Fig. S1) and OPLS loading S-plot (Fig. 1b) revealed that the discriminative inducible metabolites associated with the P-limiting treatment were mainly UK1, SQDG and DGDG (Table 1), and the significantly decreased metabolites were phospholipids (Table 2). We focused on the new lipid UK1 for further study.

Mass spectral analyses revealed that UK1 is a lipid class consisting of several similar molecular species whose molecular weights differed by 2 Da (Supplementary Fig. S2a,c). The tandem mass spectrum (MS/MS) of UK1 further revealed that the major product ions were attributable to two fatty acids and glucuronosylglycerol (Fig. 2a, and Supplementary Fig. S2b), which suggests that UK1 is a glucuronosylated diacylglycerol (Fig. 2b). MS/MS results in the positive ion mode, exhibiting the neutral loss attributable to glucuronic acid, also supported this conclusion (Supplementary Fig. S2c,d). The presence of 1,2-diacyl-3-O- α -glucuronosylglycerol (GlcADG, **1**) has been reported in several bacteria and fungi and in two species of




Figure 1 | A new lipid inducibly accumulates in *Arabidopsis* under P depletion. (a) Wild-type *Arabidopsis* plants grown under P-sufficient conditions were transferred to either P-sufficient (+P, blue line) or P-depleted (-P, red line) medium. After cultivation for 14 days, crude lipid extracts from the leaves were analysed by HILIC-MS. The total ion current detected by the negative ion mode is shown. An unknown lipid (denoted as UK1, further identified as GlcADG) appeared only under P-limited conditions. (b) S-plot of OPLS-DA based on ANOVA of the cross-validated residuals (CV-ANOVA). The S-plot demonstrates the covariance versus the correlation in conjunction with the variable trend plots and allows easier visualization of the data. The variables that showed maximum change, and therefore strongly contributed to the class separation, are plotted at the top or bottom of the S-shaped plot. The variables that did not significantly vary are plotted in the middle. Each point represents a lipid molecule detected in the negative ion mode. Red stars denote the $[M-H]^-$ ions of the UK1 species, indicating that the change in the levels of a series of UK1 molecules is highly associated with P-limiting treatment. The molecules at the top and bottom are listed in Tables 1 and 2.

Table 1 | Discriminative metabolites predicted by OPLS-DA with increased levels following P deprivation.

<i>m/z</i>	<i>t_R</i>	Annotation	Fold change in peak heights (– P/ + P)	P (corr)
592.180	3.513	*	‡	0.92
197.811	9.713	†	‡	0.97
765.523	11.403	UK1_34:3 ([M–H] [–])	‡	0.99
591.175	3.534	*	26.9	0.94
818.520	9.713	SQDG_34:2 ([M–H + 1] [–])	5.25	0.97
817.518	9.718	SQDG_34:2 ([M–H] [–])	4.61	0.99
837.495	9.678	SQDG_36:6 ([M–H] [–])	3.95	0.99
816.509	9.719	SQDG_34:3 ([M–H + 1] [–])	3.75	0.99
815.504	9.725	SQDG_34:3 ([M–H] [–])	3.07	0.98
959.604	11.024	DGDG_34:3 ([M + HCO ₂] [–])	2.92	0.99
960.608	11.028	DGDG_34:3 ([M + HCO ₂ + 1] [–])	2.82	0.98
961.615	11.024	DGDG_34:2 ([M + HCO ₂] [–])	2.58	0.97
797.549	2.849	MGDG_34:3 ([M + HCO ₂] [–])	2.11	0.97
983.596	10.986	DGDG_36:5 ([M + HCO ₂] [–])	2.04	0.97
621.441	4.369	Sitosteryl glucoside ([M + HCO ₂] [–])	1.70	0.96
821.546	2.81	MGDG_36:5 ([M + HCO ₂] [–])	1.62	0.94
982.590	10.989	DGDG_36:6 ([M + HCO ₂ + 1] [–])	1.52	0.95
820.535	2.825	MGDG_36:6 ([M + HCO ₂ + 1] [–])	1.50	0.94
819.532	2.824	MGDG_36:6 ([M + HCO ₂] [–])	1.47	0.91
981.588	10.981	DGDG_36:6 ([M + HCO ₂] [–])	1.45	0.92

DGDG, digalactosyldiacylglycerol; MGDG, monogalactosyldiacylglycerol; OPLS-DA, orthogonal projection to a latent structure-discriminant analysis; SQDG, sulfoquinovosyldiacylglycerol. *Signals of unidentified lipids whose levels in leaves are relatively high in young leaves, and gradually decrease with the growth of the plant when the plants are grown under nutrition-sufficient conditions. †Fragment ion that has not been assigned, but it is presumably derived from SQDG because it exhibits almost the same chromatographic behaviour. In addition, this ion is not detected in all the SQDG-deficient mutants of *A. thaliana*. ‡The fold changes could not be calculated because these signals are only observed in P-depleted plants.

Table 2 | Discriminative metabolites predicted by OPLS-DA with decreased levels following P deprivation.

<i>m/z</i>	<i>t_R</i>	Annotation	Fold change in peak heights (– P/ + P)	P (corr)
834.533	12.645	PI_34:2 ([M–H + 1] [–])	– *	– 0.97
714.514	12.256	PE_34:2 ([M–H] [–])	0.14	– 0.99
715.519	12.249	PE_34:2 ([M–H + 1] [–])	0.16	– 0.99
712.498	12.256	PE_34:3 ([M–H] [–])	0.25	– 0.99
742.481	10.458	PG_34:4 ([M–H + 1] [–])	0.27	– 0.99
833.526	12.644	PI_34:2 ([M–H] [–])	0.27	– 0.98
741.478	10.456	PG_34:4 ([M–H] [–])	0.28	– 0.98
713.501	12.264	PE_34:3 ([M–H + 1] [–])	0.31	– 0.99
802.565	15.764	PC_34:2 ([M + HCO ₂] [–])	0.33	– 0.99
719.494	10.504	PG_32:1 ([M–H] [–])	0.37	– 0.98
831.511	12.64	PI_34:3 ([M–H] [–])	0.37	– 0.97
747.525	10.483	PG_34:1 ([M–H] [–])	0.40	– 0.92
738.513	12.196	PE_36:4 ([M–H] [–])	0.44	– 0.95
745.508	10.465	PG_34:2 ([M–H] [–])	0.46	– 0.95
743.494	10.451	PG_34:3 ([M–H] [–])	0.46	– 0.99
744.497	10.459	PG_34:3 ([M–H + 1] [–])	0.48	– 0.99
795.532	2.83	MGDG_34:4 ([M + HCO ₂] [–])	0.53	– 0.91
793.513	2.888	MGDG_34:5 ([M + HCO ₂] [–])	0.76	– 0.91
792.506	2.898	MGDG_34:6 ([M–H + 1] [–])	0.82	– 0.92
791.503	2.894	MGDG_34:6 ([M + HCO ₂] [–])	0.83	– 0.94

MGDG, monogalactosyldiacylglycerol; OPLS-DA, orthogonal projection to a latent structure-discriminant analysis; PC, phosphatidylcholine; PE, phosphatidylethanolamine; PG, phosphatidylglycerol; PI, phosphatidylinositol. *This signal is not detected under P-depleted conditions.

unicellular algae^{28–30}, but it has not yet been described in higher plants. In addition, the physiological function and biosynthetic gene have not been identified in these organisms. Thus, we purified UK1 from *A. thaliana* and GlcADG from the proteobacterium *Brevundimonas diminuta*, previously called *Pseudomonas diminuta*³¹, and compared them with respect to several physicochemical properties.

The new plant lipid was identified as GlcADG. The purified UK1 and bacterial GlcADG showed the same chromatographic behaviour (Fig. 2b) and same fragmentation pattern (Fig. 2a, and Supplementary Figs S3,S4), supporting the hypothesis that UK1 was GlcADG. However, further characterization of the sugar moiety was still required to identify UK1. At first, we intended to analyse the NMR spectra of UK1 using bacterial GlcADG as the




Figure 2 | The novel lipid is glucuronosyldiacylglycerol (GlcADG). (a) MS/MS of $[M-H]^-$ ion of the major UK1 molecule (34:3 UK1, m/z 765.517). FA, fatty acid. The asterisk indicates the key fragment ion attributable to the glucuronosylglycerol moiety. (b) Comparison of the chromatographic behaviour of UK1 and GlcADG isolated from *B. diminuta*. The structure of the most major GlcADG molecule from *Arabidopsis* (34:3 GlcADG) is presented. The species labels, namely, total acyl carbons:total acyl double bonds, are shown in parentheses. Fragment analysis in the negative mode allowed annotation of fatty acyl groups and tentative assignment of these groups to the *sn*-1 and *sn*-2 positions⁶⁰.

reference; however, NMR spectra of GlcADG from *B. diminuta* were not available because the bacterial GlcADG was identified via several other methods of chemical derivatization³¹. Thus, we carefully analysed the structure of these two lipids from two different organisms.

First, GlcADG from *B. diminuta* was hydrolysed with hydrochloric acid, and the hydrolysate was analysed using GC-MS. The compound found in the hydrolysate was identified as glucuronic acid, not galacturonic acid, by comparison with the commercially available standard (Supplementary Figs S5,S6), indicating that the glycosyl moiety of isolated bacterial lipid is glucuronic acid. Then, we analysed NMR spectra of the methyl ester of GlcADG from *B. diminuta*. The ¹H and ¹³C NMR data on the methyl ester of bacterial GlcADG is very similar to previously published data on GlcADG from *Caulobacter bacteroides*³² and *Aspergillus fumigatus*³³, respectively, considering the effect of methylation of the carboxyl group of glucuronic acid. In addition, the ¹H and ¹³C NMR spectra of the methyl ester of bacterial UK1 are also very similar to those of the methyl ester of bacterial GlcADG. Thus, the lipids isolated from *A. thaliana* and *B. diminuta* are both GlcADG (1). The coupling constants of the anomeric proton (H-1') of GlcADG from *A. thaliana* and *B. diminuta* are both 3.7 Hz, indicating that the configuration of the glycosidic bond in these lipids is 'alpha'. The coupling pattern of protons in the uronic acid moiety is also consistent with the conclusion that the sugar moiety of this lipid class is glucuronic acid.

Elucidation of the possible biosynthetic pathway of GlcADG.

The biosynthesis of lipids occurs in the discrete subcellular biogenic membranes of different cellular compartments, and the fatty composition of the glycerolipids synthesized in one compartment is quite different from those synthesized in other compartments. For example, endoplasmic reticulum-derived glycerolipids have an 18-carbon fatty acid at *sn*-2 of glycerol, while glycerolipids synthesized *de novo* in plastids have a 16-carbon fatty acid at *sn*-2 (refs 34,35). Some of these lipids are transported from one compartment to another and transformed into different lipid classes by changing their polar headgroup. This transformation is catalysed by several different enzymes that are not evenly distributed in every organelle. These mechanisms for membrane lipid biosynthesis make the fatty acid composition and diacylglycerol (DAG) composition of each lipid class unique.

Thus, the MS and MS/MS spectral information obtained as lipidomic data were further utilized to elucidate the biosynthetic mechanism of GlcADG production, based on the assumption that each glycerolipid class possesses a characteristic fatty acid composition that reflects their synthetic route. Hierarchical clustering (HCL) of the *z*-scored DAG composition of each lipid class demonstrated that the DAG composition of GlcADG and SQDG were nearly identical, indicating that GlcADG belongs to the same group as the SQDG produced during P limitation (Fig. 3a). Analysis of the negative MS/MS data on these lipids further confirmed the high similarities between the fatty acid compositions of GlcADG and SQDG (Supplementary Fig. S3). It should be noted that all plant GlcADG species with acyl groups consisting of 34 carbons have a palmitic acid moiety at the *sn*-2 position (Supplementary Fig. S3). Palmitic acid (16:0 fatty acid) is a minor acyl group both in galactolipids and at *sn*-2 of phospholipids, but it is the major group at *sn*-2 in SQDG (Supplementary Fig. S3)^{34,35}. These results collectively suggest that the biosynthesis of GlcADG may share, at least in part, the biosynthetic pathway of SQDG.

SQDG synthase is required for GlcADG synthesis.

In *A. thaliana*, SQDG is synthesized in the chloroplast by three unique steps (Fig. 4a): synthesis of UDP-glucose (UDP-Glc) by UDP-Glc pyrophosphorylase 3 (UGP3)¹⁷, assembly of UDP-Glc and sulphite into UDP-sulfoquinovose (UDP-SQ) by UDP-SQ synthase (SQD1)^{11,36}, and the subsequent transfer of the SQ moiety from UDP-SQ to DAG by SQDG synthase (SQD2)^{12,37}. These three SQDG biosynthetic enzymes are located in the chloroplast, and the expression of their coding genes is coordinated in green tissues and significantly upregulated by P limitation^{11,12,17,37-40}. Based on the above-mentioned hypothesis that GlcADG synthesis partly shares the pathway of SQDG synthesis, GlcADG accumulation was investigated in a series of mutants deficient in all three SQDG biosynthetic genes of *A. thaliana* under the P-depleted condition. The *sqd1* and *ugp3* mutants produced GlcADG at a slightly higher level than the wild-type control (Fig. 3b and Supplementary Fig. S7); strikingly, however, no accumulation of GlcADG was observed in the independent *sqd2* mutants. These results clearly demonstrate that GlcADG is synthesized by the action of SQD2 (Fig. 4a), presumably using DAG and a sugar donor—most likely UDP-glucuronic acid (UDP-GlcA)—instead of UDP-SQ, which is used




Figure 3 | Compositional changes of lipid species under P starvation. (a) HCL of glycerolipid classes based on the z-scored DAG compositions of the fatty acid moieties. Key: P-sufficient (+P), P-depleted (-P). To test the rigour of the presented dendrogram, we performed a permutation test using jackknife resampling (permutation, 1,000). The colours of the branches indicate the level of branch rearrangement reliability (jackknife proportions) (black: 95–100% support, blue: 80–90% support). (b) Levels of individual lipid classes are expressed as relative abundance (Rel. ab.) against that of the wild-type grown under P-sufficient conditions. GlcADG was not detected under the P-sufficient condition. Each data point represents the mean \pm s.d. value of four experiments. Different letters indicate statistically significant differences ($P < 0.05$, Tukey’s test) among the tested genotypes grown under identical growth conditions. The absence of letters demonstrates no significant difference among all the tested genotypes for a lipid class. The asterisk denotes ‘not detected’.




Figure 4 | Investigation of GlcADG biosynthesis and growth phenotypes of Arabidopsis mutants. (a) Three enzymes (UGP3, SQD1 and SQD2) are involved in the biosynthesis of SQDG in the chloroplasts of *Arabidopsis*. SQD2 is responsible for SQDG synthesis under P-sufficient conditions. Under P-deficient conditions, SQD2 produces not only SQDG but also GlcADG. (b) Wild-type plants and a series of SQDG-deficient mutants of *A. thaliana* grown under P-sufficient conditions were transferred to either P-sufficient or P-depleted medium. After cultivation for 4 weeks, the growth of each genotype was compared. The individuals showing the most typical phenotype of P deprivation are displayed. Only 2 *sqd2* mutants appeared bleached under P-starvation conditions. Scale bar, 1 cm. Images of other individuals are shown in Supplementary Fig. S11.

in the synthesis of SQDG. However, two other enzymes involved in SQDG synthesis, UGP3 and SQD1, are not involved in the synthesis of GlcADG.

GlcADG deficiency leads to significant growth defects. An *Arabidopsis* mutant with disrupted *SQD2* demonstrates impaired growth in the P-limited condition¹². Therefore, we carefully examined the lipid profiles and growth of a series of SQDG-deficient *Arabidopsis* mutants with identical genetic backgrounds. Under the P-sufficient condition, the total levels of PG were slightly higher in the SQDG-deficient mutants than in the wild-type (Fig. 3b), as previously reported^{11,12,17}. Detailed analysis revealed that 34:2–34:4 PG and 34:3 PI accumulated more in the SQDG-deficient mutants than in the wild-type (Supplementary Fig. S8). Upon P limitation, the *ugp3* and *sqd1* mutants accumulated GlcADG at slightly higher levels than the wild-type (Fig. 3b and Supplementary Fig. S9). In addition, the decrease of PI, PG and PE following P limitation was suppressed in the SQDG-deficient mutants, as the total levels of GlcADG and SQDG were lower (Fig. 3b). In particular, 34:2–34:3 PI and 34:2–34:3 and 36:5 PE were significantly elevated in the *sqd2* mutant compared with the other tested genotypes (Supplementary Fig. S9). In contrast to these significant changes in phospholipids, the deficiency of GlcADG and SQDG led to only a subtle effect on galactolipid (MGDG and DGDG) contents (Fig. 3b). The levels of glucosylceramide (GlcCer) and steryl glucoside (StGlc), which serve along with glycerolipids as membrane lipids, were also analysed. The amount of GlcCer and StGlc increased following P limitation, and the amounts of StGlc in P-starved *sqd2* was higher compared with the other tested genotypes (Supplementary Fig. S10ab). There were no clear differences in the total levels of GlcCer among the tested genotypes when P starved, although there were some differences among their GlcCer profiles (Supplementary Fig. S10c). These results indicate that the lipid remodelling mechanisms induced by P limitation are significantly altered by GlcADG deficiency.

We then investigated the growth of the SQDG-deficient mutants under P-depleted conditions. Under P-sufficient conditions, all SQDG-deficient mutants and the wild-type demonstrated similar growth (Supplementary Fig. S11a). In contrast, under P-depleted conditions, the two independent *sqd2* mutants that did not accumulate SQDG or GlcADG were severely damaged compared with the *sqd1* and *ugp3* mutants, in which SQDG is absent but GlcADG is still produced (Fig. 4b and Supplementary Fig. S11b). These results indicate that GlcADG has a major role in mitigating P-depletion stress in plants. In addition, the leaf colour of the P-starved *sqd2* mutants appeared bleached. The total chlorophyll content was reduced in *sqd2* mutants compared with wild-type, *ugp3* or *sqd1* when plants were grown under P depletion (Supplementary Fig. S12). These results indicate that GlcADG deficiency accelerates the breakdown of chlorophylls and the de-regulation of chloroplast functions.

As described above, the disruption of the *SQD2* gene in *Arabidopsis* led to the de-regulation of lipid remodelling under P depletion, and the shoots of *sqd2* mutants showed enhanced senescence compared with other SQDG-deficient mutants. These data suggest that P_i levels might be affected by disruption of *SQD2*. Thus, we analysed the inorganic P_i level in shoots grown under P-controlled conditions. The inorganic P_i levels in shoots were significantly altered as P_i availability changed, whereas there was no significant difference in the P_i content of the shoots of the tested genotypes (Supplementary Fig. S13). Thus, the severe growth defects observed in the shoots of *sqd2* mutants under P-limited conditions are not necessarily related to the free P_i content in shoots, suggesting an association of GlcADG in the

protection of the membrane environment under P-limited growth conditions.

To obtain further insight into the general role of GlcADG in P depletion in plants, the distribution of GlcADG was also investigated in rice (*Oryza sativa* L.). GlcADG was detected in the leaves of rice grown even under the nutrition-sufficient condition (Supplementary Fig. S14), and the levels were significantly elevated (5–7 times) by P deprivation, as observed in *A. thaliana* (Fig. 5). Significant lipid profile changes upon P limitation were also observed in rice leaves, as observed in *A. thaliana* (Supplementary Fig. S15). These results suggest a general physiological significance for this novel lipid under P-limited conditions across plant species.

Discussion

SQDG is an unusual glycerolipid because its polar headgroup consists of SQ, a 6-deoxy-6-sulfonate-glucose (Fig. 4). Much has been speculated about the possible function of this lipid in plants⁴¹. However, the isolation of SQDG-deficient bacterial mutants^{42,43}, a mutant of the unicellular algae *Chlamydomonas*⁴⁴, and a mutant of the plant *A. thaliana*¹² have all provided some clues about its function. It has become apparent from these mutants that there is no essential role for SQDG in photosynthetic bacteria or in eukaryotic cells containing chloroplasts because all mutants are capable of photoautotrophic growth. The effects on photosynthesis are subtle under normal growth conditions. However, if these SQDG-deficient null mutants are starved for P, they cease growth much earlier than the corresponding wild-type^{12,42,43}. Thus, SQDG has been considered to be of conditional importance in these organisms, including plants. In addition, SQDG and PG contents are known to be maintained by reciprocally balancing their levels with changes in P availability, which is considered to contribute to the maintenance of the negative charge in thylakoid membranes^{12,45}. However, among the three SQDG-deficient mutants of *A. thaliana*, the severe growth defect under the P-limited condition was observed only in the *sqd2* mutant (Fig. 4b and Supplementary Fig. S11)¹², and no studies have investigated




Figure 5 | GlcADG inducibly accumulates in rice leaves upon P limitation.

Comparison of GlcADG levels in the leaves of rice grown under either P-sufficient (+P) or P-depleted (-P) conditions. The levels of each lipid molecule are expressed as relative values compared with the level of 34:2 GlcADG in rice grown under P-sufficient growth conditions. Each data point represents the mean value of three experiments \pm s.d. Asterisks indicate a statistically significant difference from the P-sufficient growth condition ($*P < 0.05$, Welch's *t*-test). More detailed data can be found in Supplementary Figs S14, S15.

the differences in the lipid profiles of these SQDG-deficient mutants^{11,12,17}.

Our present results unexpectedly revealed that the P-starved *sqd2* mutant of *A. thaliana* lacks not only SQDG but also a novel anionic glycolipid GlcADG, which results in a severe phenotype in this mutant that is not observed in the other two SQDG-deficient mutants analysed in this study. In addition, under P limitation, the *ugp3* and *sqd1* mutants accumulated slightly higher levels of GlcADG than the wild-type and showed no apparent growth defects. These results collectively indicate the following: (1) GlcADG is a novel molecule importantly involved in the protection against P-depletion stress in plants; (2) the growth defect of P-starved *sqd2* is attributable to the deficiency of the two anionic glycolipids SQDG and GlcADG; and (3) moderate increases in GlcADG can compensate for the SQDG deficiency, as shown in the *sqd1* and *ugp3* mutants. Our results reinforce the importance of plant SQDG synthase and anionic glycolipids synthesized by SQDG synthase during growth under P-limited conditions.

The details of the biosynthesis of GlcADG remain unknown except for the fact that SQDG synthase is required for GlcADG biosynthesis. We propose a possible biosynthetic scheme for this glycolipid in Fig. 4. However, our lipidomic data using a few SQDG-deficient mutants do not deny the possibility of interconversion of SQDG and GlcADG. Other types of experiments, such as flux analysis, will be important in understanding the metabolic flow underlying the accumulation of these two anionic glycolipids. It is also postulated that the inducible synthesis of GlcADG under P starvation entails increased synthesis of an unidentified sugar donor, presumably UDP-GlcA, in chloroplasts. Our current study suggests the possibility of the presence of an uncharacterized biosynthetic pathway or transportation mechanism for UDP-GlcA in chloroplasts. The synthesis of GlcADG by dehydrogenation of monoglucosyldiacylglycerol, found in several plants⁴⁶, is also possible.

In P-starved *sqd2* mutants, the levels of PI and PE were significantly elevated, which indicates that the complete deficiency of anionic glycolipids causes the de-regulation of not only PG but also other phospholipids, such as PI and PE. Thus, levels

of these phospholipids (PI and PE) and GlcADG are also reciprocally maintained in plant cells under P-limited conditions (Fig. 6), as a similar relationship can be found between SQDG and PG, suggesting some shared functions among GlcADG, PI and PE under P-limited conditions. Levels of 34:3 PI and 34:3 PE were also slightly elevated in *sqd1* and *ugp3* mutants compared with wild-type even under P sufficiency (Supplementary Fig. S8). Thus, these phospholipids may also generally contribute to the maintenance of proper membrane properties in plants even under P sufficiency, and their roles may become more important when plants are deficient in two anionic glycolipids under P-depleted conditions.

Currently, contributions to the compensation of the deficiency of anionic glycolipids are likely to be significant in the order of PG > PI > PE, although it remains unknown whether this order is because of the difference in chemical properties or the difference in the organelle where they are mainly synthesized, that is, chloroplastic PG is synthesized in chloroplasts⁴⁷, such as SQDG and presumably GlcADG, whereas PI and PE are mainly synthesized in the endoplasmic reticulum (Fig. 6). We have never determined where GlcADG is synthesized and accumulates, but GlcADG is likely synthesized in chloroplasts because SQDG synthase is localized in the chloroplast inner membrane³⁷.

The idea that GlcADG accumulates in chloroplasts is plausible considering the localization of SQDG synthase, although we cannot exclude the possibility that somehow GlcADG is exported from chloroplasts to extraplastidic membranes such as DGDG^{6,8}. Given that GlcADG accumulates in extraplastidic space, we might be able to determine the reason why the levels of GlcADG, PI and PE are reciprocally regulated in P-starved plants. In addition, we cannot deny the possibility that levels of PI and PE in the chloroplast membrane are elevated in *sqd2* mutants under P deficiency even though PE is usually not abundant in chloroplasts⁴⁸. Further research to determine the localization of GlcADG accumulation will help us to understand this phenomenon. A study on the bacterium *B. diminuta* also investigated the replacement of acidic phospholipids by GlcADG upon P limitation⁴⁹, suggesting a functional convergence of lipid remodelling involving the accumulation of GlcADG across




Figure 6 | Anionic glycolipids and phospholipids are reciprocally balanced under P depletion. An abbreviated biosynthetic scheme for the major plant polar glycerolipids is represented along with the results of the current report. Previous research on several SQDG-deficient mutants suggested that the levels of SQDG and PG are reciprocally regulated (orange line), while our current report suggests that a similar reciprocal regulation also can be found among GlcADG, PI and PE (red lines). The levels of SQDG, PI and PE might also be reciprocally maintained.

kingdoms. No information regarding the genes involved in GlcADG biosynthesis in *B. diminuta* is currently available.

Lipid remodelling in membrane systems under P-limited conditions is considered to contribute to sustaining the level of cellular inorganic P_i by modifying the P allocation in the cell. As described above, lipid remodelling is severely de-regulated in a P-starved *sqd2* mutant, but there is no significant decrease in free P_i in this mutant (Supplementary Fig. S13). These results suggest that GlcADG is not essential to maintain the free P_i level in P-starved plants but that it is likely to contribute to the control of the membrane environment to ensure the proper functioning of organelles, including chloroplasts, in P-limited, unfavourable environments.

In addition, this study suggests that plants have developed an alternative mechanism to maintain a negative membrane charge by synthesizing uronic acid-containing lipids free of sulphur, which is also an essential macronutrient requiring significant energy for its assimilation⁵⁰. Genome sequence information on organisms capable of synthesizing SQDG has revealed that the two genes coding UDP-SQ synthase and SQDG synthase evolved at different times, and while many genomes show readily identifiable UDP-SQ synthase orthologs, SQDG synthases were differently recruited in cyanobacteria/plants versus non-cyanobacteria and archaea⁵¹. The involvement of plant SQDG synthase in different anionic glycolipids might be related to its ancestral function, which might still be retained in plants to allow them adapt to unfavourable surroundings.

Methods

Chemicals. Primuline, (trimethylsilyl)diazomethane (2.0 M in diethyl ether), and 1,2-didecanoyl-sn-glycero-3-phosphocholine were obtained from Sigma-Aldrich, and polypropylene glycol 2000 (PPG-2000) was obtained from Alfa Aesar. Glucuronic acid was purchased from Tokyo Chemical Industry. Unless stated otherwise, all other chemicals were purchased from Wako Pure Chemical Industries.

Nuclear magnetic resonance. ¹H NMR spectra were recorded in CDCl₃ supplemented with a drop of D₂O for H-D exchange. Other NMR spectra (¹³C, COSY, HMQC, HMBC and DEPT) were recorded in absolute CDCl₃. Tetramethylsilane was used for the internal standard.

Plant materials. *A. thaliana*—Seeds of T-DNA insertion lines for *sqd2-1* (SALK_070595) and *sqd2-2* (SALK_139798) were purchased from ABRC at Ohio State University⁵². The T-DNA insertion sites of these lines were confirmed by sequencing PCR fragments (Supplementary Fig. S16). A PCR fragment at the left border of the T-DNA of *sqd2-1* was amplified using *Lba1* and *SALK_070595_Fw*, and the right border of the T-DNA of *sqd2-2* was amplified using *SALK_070595_Fw* and *RBa1* (Supplementary Table S1). The *sqd1*, *ugp3-1* and *ugp3-2* mutants were isolated in our previous study based on the results of genome PCR¹⁷ and used again in this research. Seeds of the *sqd2* mutant in the *Ws-0* background (provisionally named *sqd2-3* in this study) and its wild-type were kind gifts from Professor Benning¹². Surface-sterilized seeds were sown on agar-solidified Murashige and Skoog medium⁵³ containing 1% (w/v) sucrose at 22 °C under 16 h light/8 h dark cycles. Photon flux density was 50 μmol m⁻² sec⁻¹. The seedlings were kept on agar for 14 days before transfer to plates with controlled phosphate (P_i) concentrations. For P-limitation experiments, sterile *Arabidopsis* medium⁵⁴ was used but at half concentration and containing 0.8% (w/v) agar, 20 mM MES (pH 6.0) and different concentrations of KH₂PO₄ as indicated⁷. For lipid analysis, leaves were harvested 6 h after the onset of the light phase, frozen in liquid nitrogen and stored at -80 °C until lipid extraction.

Rice—Dehulled seeds of *O. sativa* L. cv. Nipponbare were sterilized and germinated on agar-solidified Murashige and Skoog medium at 28 °C under a 16 h light/8 h dark cycle. To expose plants to P-limiting stress, 5-day-old seedlings were transplanted and hydroponically cultured in the P-controlled medium used for the stress treatment of *A. thaliana*. The culture medium was changed every week.

Lipidomics by HILIC-MS. Plant lipids were extracted following the method of Blich and Dyer⁵⁵ with slight modifications²⁶, and directly subjected to the extract to LC-MS analysis. The lipidomic data were recorded on an LC-MS-IT-TOF mass spectrometer using electrospray ionization, combined with an LC-20AD HPLC system (Shimadzu)²⁶. Parameters for MS were as follows: scan range, m/z 150–1,600; loop time, 0.6 s; CDL and heat block temperature, 160 °C; nebulizer gas (N₂),

1.51 min⁻¹. More details are found in our previous report²⁶ and metadata in 'Supplementary Data Set.xls'.

Peak picking and alignment were performed using the Profiling Solution software (version 1.0.76.0; Shimadzu). The profiling solution parameters were as follows: ion m/z tolerance, 20 mDa; ion RT tolerance, 0.1 min; ion intensity threshold, 2e4; detect isomer valley, 20%; allow some ion without isotope peak, ON²⁶. The data matrix containing ~480 ions consisting of negatively detected ions was exported from the profiling solution software and normalized relative to the intensity of the [M + HCO₂]⁻ ion of the internal standard (1,2-didecanoyl-glycero-3-phosphocholine). Annotation was based on the theoretical m/z values of each lipid species and the retention time of identified compounds with the same polar head²⁶.

Statistics for the analysis of lipidomic data. Normalized LC-MS data were pareto-scaled and subjected to OPLS-DA using SIMCA-P (version 11.0.0.0; Umetrics). Tukey's test and Welch's *t*-test were performed using their respective functions (TukeyHSD and *t*-test) in the statistical software R version 2.14.0 (<http://www.r-project.org/>). HCL and jackknife testing were performed on Multi-Experiment Viewer (MeV) version 4.8 (<http://www.tm4.org/mev/>). For HCL, Pearson's correlations were used to analyse the distance metric. Average linkage clustering was selected for the linkage method. The data set used for HCL is available as 'Supplementary Data Set.xls'. Metadata also can be found in the same file.

Preparation of GlcADG from *A. thaliana* and *B. diminuta*. *A. thaliana* plants (Col-0 accession, 340 g fresh weight) grown under P deficiency as described above were extracted using the same protocol to prepare samples for lipidomic analysis. The resulting lipid extracts (4.1 g) was loaded onto a silica gel column (silica gel 60 N, spherical, neutral; 25-cm long, 7-cm i.d.; Kanto Chemical) equilibrated with chloroform. The column was washed with chloroform (3 l), chloroform-acetone (1:1, v/v, 3.6 l), acetone (1 l), acetone-methanol (1:1, v/v, 1 l) and methanol (1 l). The acetone-methanol fractions containing UK1 were subjected to preparative thin-layer chromatography on Silica Gel 60 plates containing no fluorescent indicator (Merck) using chloroform-methanol-H₂O (65:25:4, v/v/v). After spraying a primuline solution, lipids were detected under UV light. The fluorescent spots containing UK1 were scraped off and further subjected to preparative thin-layer chromatography with a chloroform-methanol-acetic acid solution (75:25:3, v/v/v) to obtain UK1 (1, 9.6 mg). MS/MS of all detectable UK1 molecules, are found in Fig. 2 and Supplementary Figs S2,S3.

B. diminuta ATCC 11568 (Strain JCM 2788) was provided by RIKEN BRC (Wako, Japan). The strain was rehydrated according to the provider's instructions and cultivated with continuous rotation at 180 r.p.m. in lysogeny broth containing PPG-2000 (50 p.p.m.) as antifoam. After a 48-h incubation at 30 °C, the cells were collected by centrifugation. The dried cells (4.43 g) were extracted with mixtures of chloroform, methanol and diethyl ether to yield crude lipid extracts (320 mg) as previously reported^{56,57}. The extracts were separated to purify GlcADG (17.8 mg) using the same protocol described above for the purification of UK1. MS and MS/MS of the bacterial GlcADG are found in Supplementary Fig. S4.

Preparation of the methyl ester of GlcADG. GlcADG from *B. diminuta* and UK1 from *A. thaliana* were methylated with (trimethylsilyl)diazomethane as previously described⁵⁸. The ¹H and ¹³C NMR spectra and key HMBCs of bacterial GlcADG are shown in Supplementary Figs S17–S19.

Methyl ester of GlcADG from *B. diminuta*. ¹H NMR (500 MHz, CDCl₃) δ 0.84–0.93 (6 H, *m*, -CH₂CH₃), 1.19–1.45 (ca 36 H, *m*, -CH₂-), 1.54–1.66 (4 H, *m*, -OCOCH₂CH₂-), 1.93–2.10 (5 H, *m*, -CH=CHCH₂-), 2.25–2.39 (4 H, *m*, -OCOCH₂R), 3.57 (1 H, *dd*, *J* = 3.7, 9.5 Hz, H-2'), 3.67 (1 H, *dd*, *J* = 5.7, 10.9 Hz, H-3a), 3.72 (1 H, triplet-like *dd*, *J* = 8.9, 9.5 Hz, H-4'), 3.78 (1 H, triplet-like *dd*, *J* = 8.9, 9.5 Hz, H-3'), 3.83 (3 H, *s*, CH₃O-6'), 3.90 (1 H, *dd*, *J* = 4.6, 10.9 Hz, H-3b), 4.15 (1 H, *dd*, *J* = 5.7, 11.9 Hz, H-1a), 4.16 (1 H, *d*, *J* = 9.5 Hz, H-5'), 4.41 (1 H, *dd*, *J* = 4.2, 11.9 Hz, H-1b), 4.94 (1 H, *d*, *J* = 3.7 Hz, H-1'), 5.26 (1 H, *m*, H-2), 5.33–5.36 (2 H, *m*, -CH=CH-); ¹³C NMR (125 MHz, CDCl₃) δ 14.1 (-CH₂CH₃), 22.66–22.70 (-CH₂CH₂CH₃), 24.9 (-OCOCH₂CH₂CH₂-), 27.2 (-CH=CHCH₂CH₂-), 28.7–31.9 (-CH₂CH₂CH₂-), 34.1–34.2 (-OCOCH₂CH₂-), 52.9 (CH₃O-6'), 61.9 (C-1), 67.1 (C-3), 69.6 (C-2), 70.6 (C-5'), 71.5 (C-2'), C-4'), 73.6 (C-3'), 99.1 (C-1'), 129.8 (-CH=CH-), 130.0 (-CH=CH-), 170.5 (C6'), 173.2 (-OCOR of fatty acid), 173.5 (-OCOR of fatty acid).

Methyl ester of UK1 from *A. thaliana*. ¹H NMR (500 MHz, CDCl₃) δ 0.85–0.93 (ca 4 H, *m*, -CH₂CH₃), 0.95–1.02 (ca 3 H, -CH=CHCH₂CH₃), 1.2–1.4 (ca 40 H, *m*, -CH₂-), 1.5–1.7 (ca 13 H, *m*, -OCOCH₂CH₂-), 2.01–2.13 (ca 6 H, *m*, -CH=CHCH₂-), 2.27–2.37 (ca 4 H, *m*, -OCOCH₂R), 2.74–2.84 (ca 4 H, *m*, -CH=CHCH₂CH=CH-), 3.58 (1 H, *dd*, *J* = 3.7, 9.5 Hz, H-2'), 3.68 (1 H, *dd*, *J* = 5.7, 10.9 Hz, H-3a), 3.72 (1 H, triplet-like *dd*, *J* = 8.9, 9.5 Hz, H-4'), 3.79 (1 H, triplet-like *dd*, *J* = 8.9, 9.5 Hz, H-3'), 3.84 (3 H, *s*, CH₃O-6'), 3.91 (1 H, *dd*, *J* = 4.6, 10.9 Hz, H-3b), 4.14 (1 H, *dd*, *J* = 5.7, 12.0 Hz, H-1a), 4.17 (1 H, *d*, *J* = 9.5 Hz, H-5'), 4.42 (1 H, *dd*, *J* = 4.3, 12.0 Hz, H-1b), 4.95 (1 H, *d*, *J* = 3.7 Hz, H-1'), 5.23–5.29 (ca 1 H,

m, H-2), 5.29–5.44 (ca 8 H, *m*, -CH = CH-); ^{13}C NMR (125 MHz, CDCl_3) δ 14.06–14.3 (-CH₂CH₃), 20.5 (-CH = CHCH₂CH₃), 22.6–22.7 (-CH₂CH₂CH₃), 24.8 (-OCOCH₂CH₂CH₂-), 25.5–25.6 (-CH = CHCH₂CH = CH-), 27.2 (-CH = CHCH₂CH₂-), 29.1–31.9 (-CH₂CH₂CH₂-), 34.1–34.2 (-OCOCH₂CH₂-), 52.8 (CH₃O-6'), 61.9 (C-1), 67.1 (C-3), 69.7 (C-2), 70.6 (C-5'), 71.5 (C-2', C-4'), 73.6 (C-3'), 99.1 (C-1'), 127.1–132.0 (olefinic carbons of fatty acids), 170.4 (C6'), 173.1 (-OCOR of fatty acid), 173.5 (-OCOR of fatty acid).

Quantification of inorganic P_i in plants. Leaves (30 mg) were frozen in liquid nitrogen and homogenized using a Mixer Mill MM300 (Retsch, Haan, Germany) for 2 min at 15 Hz with a zirconium bead and then extracted with 50% (v/v) aqueous methanol (300 μl) followed by centrifugation at 12,000 *g*. The milling process was carried out under cryogenic conditions (the grinding jars were cooled with liquid nitrogen before each milling process to ensure a low milling temperature and keep the plants frozen). The measurement of P_i content in the supernatants was accomplished by forming a phosphomolybdate complex and in turn reducing it to a molybdenum blue complex⁵⁹.

References

- Raghothama, K. G. Phosphate transport and signaling. *Curr. Opin. Plant Biol.* **3**, 182–187 (2000).
- Vance, C. P. Symbiotic nitrogen fixation and phosphorus acquisition. *Plant nutrition in a world of declining renewable resources. Plant Physiol.* **127**, 390–397 (2001).
- Cordell, D., Rosemarin, A., Schroder, J. J. & Smit, A. L. Towards global phosphorus security: A systems framework for phosphorus recovery and reuse options. *Chemosphere* **84**, 747–758 (2011).
- Gilbert, N. Environment: the disappearing nutrient. *Nature* **461**, 716–718 (2009).
- Van Kauwenbergh, S. J. *World phosphate rock reserves and resources* (International Fertilizer Development Center, 2010).
- Andersson, M. X., Stridh, M. H., Larsson, K. E., Liljenberg, C. & Sandelius, A. S. Phosphate-deficient oat replaces a major portion of the plasma membrane phospholipids with the galactolipid digalactosyldiacylglycerol. *FEBS Lett.* **537**, 128–132 (2003).
- Härtel, H., Dörmann, P. & Benning, C. DGD1-independent biosynthesis of extraplastidic galactolipids after phosphate deprivation in *Arabidopsis*. *Proc. Natl Acad. Sci. USA* **97**, 10649–10654 (2000).
- Jouhet, J. *et al.* Phosphate deprivation induces transfer of DGDG galactolipid from chloroplast to mitochondria. *J. Cell Biol.* **167**, 863–874 (2004).
- Benning, C. & Ohta, H. Three enzyme systems for galactoglycerolipid biosynthesis are coordinately regulated in plants. *J. Biol. Chem.* **280**, 2397–2400 (2005).
- Andersson, M. X., Larsson, K. E., Tjellström, H., Liljenberg, C. & Sandelius, A. S. Phosphate-limited oat. The plasma membrane and the tonoplast as major targets for phospholipid-to-glycolipid replacement and stimulation of phospholipases in the plasma membrane. *J. Biol. Chem.* **280**, 27578–27586 (2005).
- Essigmann, B., Guler, S., Narang, R. A., Linke, D. & Benning, C. Phosphate availability affects the thylakoid lipid composition and the expression of *SQD1*, a gene required for sulfolipid biosynthesis in *Arabidopsis thaliana*. *Proc. Natl Acad. Sci. USA* **95**, 1950–1955 (1998).
- Yu, B., Xu, C. & Benning, C. *Arabidopsis* disrupted in *SQD2* encoding sulfolipid synthase is impaired in phosphate-limited growth. *Proc. Natl Acad. Sci. USA* **99**, 5732–5737 (2002).
- Kobayashi, K. *et al.* Type-B monogalactosyldiacylglycerol synthases are involved in phosphate starvation-induced lipid remodeling, and are crucial for low-phosphate adaptation. *Plant J.* **57**, 322–331 (2009).
- Kobayashi, K., Masuda, T., Takamiya, K. & Ohta, H. Membrane lipid alteration during phosphate starvation is regulated by phosphate signaling and auxin/cytokinin cross-talk. *Plant J.* **47**, 238–248 (2006).
- Gaude, N., Nakamura, Y., Scheible, W. R., Ohta, H. & Dörmann, P. Phospholipase C5 (NPC5) is involved in galactolipid accumulation during phosphate limitation in leaves of *Arabidopsis*. *Plant J.* **56**, 28–39 (2008).
- Awai, K. *et al.* Two types of MGDG synthase genes, found widely in both 16:3 and 18:3 plants, differentially mediate galactolipid syntheses in photosynthetic and nonphotosynthetic tissues in *Arabidopsis thaliana*. *Proc. Natl Acad. Sci. USA* **98**, 10960–10965 (2001).
- Okazaki, Y. *et al.* A chloroplastic UDP-glucose pyrophosphorylase from *Arabidopsis* is the committed enzyme for the first step of sulfolipid biosynthesis. *Plant Cell* **21**, 892–909 (2009).
- Kelly, A. A. & Dörmann, P. DGD2, an arabidopsis gene encoding a UDP-galactose-dependent digalactosyldiacylglycerol synthase is expressed during growth under phosphate-limiting conditions. *J. Biol. Chem.* **277**, 1166–1173 (2002).
- Cruz-Ramirez, A., Oropeza-Aburto, A., Razo-Hernandez, F., Ramirez-Chavez, E. & Herrera-Estrella, L. Phospholipase DZ2 plays an important role in extraplastidic galactolipid biosynthesis and phosphate recycling in *Arabidopsis* roots. *Proc. Natl Acad. Sci. USA* **103**, 6765–6770 (2006).
- Nakamura, Y. *et al.* A novel phosphatidylcholine-hydrolyzing phospholipase C induced by phosphate starvation in *Arabidopsis*. *J. Biol. Chem.* **280**, 7469–7476 (2005).
- Nakamura, Y. *et al.* *Arabidopsis* lipins mediate eukaryotic pathway of lipid metabolism and cope critically with phosphate starvation. *Proc. Natl Acad. Sci. USA* **106**, 20978–20983 (2009).
- Cheng, Y. *et al.* Characterization of the *Arabidopsis* glycerophosphodiester phosphodiesterase (GDPD) family reveals a role of the plastid-localized AtGDPD1 in maintaining cellular phosphate homeostasis under phosphate starvation. *Plant J.* **66**, 781–795 (2011).
- Bari, R., Datt Pant, B., Stitt, M. & Scheible, W. R. PHO2, microRNA399, and PHR1 define a phosphate-signaling pathway in plants. *Plant Physiol.* **141**, 988–999 (2006).
- Narise, T. *et al.* Involvement of auxin signaling mediated by IAA14 and ARF7/19 in membrane lipid remodeling during phosphate starvation. *Plant Mol. Biol.* **72**, 533–544 (2010).
- Li, M., Welti, R. & Wang, X. Quantitative profiling of *Arabidopsis* polar glycerolipids in response to phosphorus starvation. Roles of phospholipases D ζ 1 and D ζ 2 in phosphatidylcholine hydrolysis and digalactosyldiacylglycerol accumulation in phosphorus-starved plants. *Plant Physiol.* **142**, 750–761 (2006).
- Okazaki, Y., Kamide, Y., Hirai, M. Y. & Saito, K. Plant lipidomics based on hydrophilic interaction chromatography coupled to ion trap time-of-flight mass spectrometry. *Metabolomics* doi:10.1007/s11306-011-0318-z (2011).
- Wiklund, S. *et al.* Visualization of GC/TOF-MS-based metabolomics data for identification of biochemically interesting compounds using OPLS class models. *Anal. Chem.* **80**, 115–122 (2008).
- Hölzl, G. & Dörmann, P. Structure and function of glycolipids in plants and bacteria. *Prog. Lipid Res.* **46**, 225–243 (2007).
- Eichenberger, W. & Gribo, C. Diacylglycerol- α -D-glucuronide from *Ochromonas danica* (Chrysothymaceae). *J. Plant Physiol.* **144**, 272–276 (1994).
- Eichenberger, W. & Gribo, C. Lipids of *Pavlova lutheri*: cellular site and metabolic role of DGCC. *Phytochemistry* **45**, 1561–1567 (1997).
- Wilkinson, S. G. Lipids of *Pseudomonas diminuta*. *Biochim. Biophys. Acta* **187**, 492–500 (1969).
- Batrakov, S. G., Nikitin, D. I., Sheichenko, V. I. & Ruzhitsky, A. O. Unusual lipid composition of the gram-negative, freshwater, stalked bacterium *Caulobacter bacteroides* NP-105. *Biochim. Biophys. Acta* **1347**, 127–139 (1997).
- Fontaine, T. *et al.* Characterization of glucuronic acid containing glycolipid in *Aspergillus fumigatus* mycelium. *Carbohydr. Res.* **344**, 1960–1967 (2009).
- Browse, J., Warwick, N., Somerville, C. R. & Slack, C. R. Fluxes through the prokaryotic and eukaryotic pathways of lipid synthesis in the '16:3' plant *Arabidopsis thaliana*. *Biochem. J.* **235**, 25–31 (1986).
- Heinz, E. & Roughan, P. G. Similarities and differences in lipid metabolism of chloroplasts isolated from 18:3 and 16:3 plants. *Plant Physiol.* **72**, 273–279 (1983).
- Sanda, S., Leustek, T., Theisen, M. J., Garavito, R. M. & Benning, C. Recombinant *Arabidopsis* SQD1 converts UDP-glucose and sulfite to the sulfolipid head group precursor UDP-sulfoquinovose *in vitro*. *J. Biol. Chem.* **276**, 3941–3946 (2001).
- Tietje, C. & Heinz, E. Uridine-diphospho-sulfoquinovose: diacylglycerol sulfoquinovosyltransferase activity is concentrated in the inner membrane of chloroplast envelopes. *Planta* **206**, 72–78 (1998).
- Seifert, U. & Heinz, E. Enzymatic characteristics of Udp-sulfoquinovose - diacylglycerol sulfoquinovosyltransferase from chloroplast envelopes. *Bot. Acta* **105**, 197–205 (1992).
- Heinz, E. *et al.* Synthesis of different nucleoside 5'-diphospho-sulfoquinovoses and their use for studies on sulfolipid biosynthesis in chloroplasts. *Eur. J. Biochem.* **184**, 445–453 (1989).
- Shimajima, M. Biosynthesis and functions of the plant sulfolipid. *Prog. Lipid Res.* **50**, 234–239 (2011).
- Benning, C. Biosynthesis and Function of the Sulfolipid Sulfoquinovosyl Diacylglycerol. *Annu. Rev. Plant Physiol. Plant Mol. Biol.* **49**, 53–75 (1998).
- Benning, C., Beatty, J. T., Prince, R. C. & Somerville, C. R. The sulfolipid sulfoquinovosyldiacylglycerol is not required for photosynthetic electron transport in *Rhodospirillum rubrum* but enhances growth under phosphate limitation. *Proc. Natl Acad. Sci. USA* **90**, 1561–1565 (1993).
- Güler, S., Seeliger, A., Härtel, H., Renger, G. & Benning, C. A null mutant of *Synechococcus* sp. PCC7942 deficient in the sulfolipid sulfoquinovosyl diacylglycerol. *J. Biol. Chem.* **271**, 7501–7507 (1996).
- Sato, N., Sonoike, K., Tsuzuki, M. & Kawaguchi, A. Impaired photosystem II in a mutant of *Chlamydomonas reinhardtii* defective in sulfoquinovosyl diacylglycerol. *Eur. J. Biochem.* **234**, 16–23 (1995).
- Yu, B. & Benning, C. Anionic lipids are required for chloroplast structure and function in *Arabidopsis*. *Plant J.* **36**, 762–770 (2003).
- Jamieson, G. R. & Reid, E. H. Sugar components of galactosyl diglycerides from green plants. *Phytochemistry* **15**, 135–136 (1976).

47. Xu, C. *et al.* The *pgp1* mutant locus of *Arabidopsis* encodes a phosphatidylglycerolphosphate synthase with impaired activity. *Plant Physiol.* **129**, 594–604 (2002).
48. Block, M. A., Douce, R., Joyard, J. & Rolland, N. Chloroplast envelope membranes: a dynamic interface between plastids and the cytosol. *Photosynth. Res.* **92**, 225–244 (2007).
49. Minnikin, D. E., Abdolrahimzadeh, H. & Baddiley, J. Replacement of acidic phosphates by acidic glycolipids in *Pseudomonas diminuta*. *Nature* **249**, 268–269 (1974).
50. Takahashi, H., Kopriva, S., Giordano, M., Saito, K. & Hell, R. Sulfur assimilation in photosynthetic organisms: molecular functions and regulations of transporters and assimilatory enzymes. *Annu. Rev. Plant Biol.* **62**, 157–184 (2011).
51. Benning, C., Gravito, R. M. & Shimojima, M. in *Sulfur Metabolism in Phototrophic Organisms (Advances in Photosynthesis and Respiration)* (eds Hell, R., Dahl, C., Knaff, D. B. & Leustek, T.) Ch. 10, 185–200 (Springer, 2008).
52. Alonso, J. M. *et al.* Genome-wide insertional mutagenesis of *Arabidopsis thaliana*. *Science* **301**, 653–657 (2003).
53. Murashige, T. & Skoog, F. A revised medium for rapid growth and bio assays with tobacco tissue cultures. *Physiol. Plant* **15**, 473–497 (1962).
54. Estelle, M. A. & Somerville, C. Auxin-resistant mutants of *Arabidopsis thaliana* with an altered morphology. *Mol. Gen. Genet.* **206**, 200–206 (1987).
55. Bligh, E. G. & Dyer, W. J. A rapid method of total lipid extraction and purification. *Can. J. Biochem. Physiol.* **37**, 911–917 (1959).
56. Wilkinson, S. G. Glycosyl diglycerides from *Pseudomonas rubescens*. *Biochim. Biophys. Acta* **164**, 148–156 (1968).
57. Wilkinson, S. G. Studies on the cell walls of *Pseudomonas* species resistant to ethylenediaminetetra-acetic acid. *J. Gen. Microbiol.* **54**, 195–213 (1968).
58. Hashimoto, N., Aoyama, T. & Shioiri, T. New methods and reagents in organic synthesis. 14. A simple efficient preparation of methyl esters with trimethylsilyldiazomethane (TMSCHN₂) and its application to gas chromatographic analysis of fatty acids. *Chem. Pharm. Bull.* **29**, 1475–1478 (1981).
59. Ames, B. N. Assay of inorganic phosphate, total phosphate and phosphatases. *Methods Enzymol.* **8**, 115–118 (1966).
60. Murphy, R. C. in *Handbook of Lipid Research* (ed Snyder, F.) vol. 7 Chapter 7, 223–226 (Plenum Press, 1993).

Acknowledgements

We are grateful to Professor Christoph Benning (Michigan State University) for kindly donating the *sqd2* mutant seeds (provisionally named as *sqd2-3*) and the wild-type *A. thaliana* seeds (Ws-0 accession). We also thank our colleague Dr Ryo Nakabayashi (RIKEN Plant Science Centre) for his advice regarding the NMR experiments. The stock of *B. diminuta* was provided by the Japan Collection of Microorganisms, RIKEN BRC (Wako, Japan), which is participating in the National BioResource Project, MEXT, Japan. This work was supported in part by grant aids from the Strategic International Research Cooperative Programme of Japan Science Technology (Metabolomics for a Low Carbon Society, JST-NSF) and a Grant-in-Aid for Scientific Research from the Ministry of Education, Culture, Sports, Science and Technology of Japan. This research was also supported by the Japan Advanced Plant Science Network.

Author contributions

Y.O. performed the metabolite profiling of plants. H.O., T.N., Y.K. and M.Y.H. performed the molecular biology experiments. Makoto K. and Miyako K. performed the GC-MS analysis. S.S. and T.A. performed the NMR experiments. Y.O. and K.S. designed the experiments and wrote the manuscript.

Additional information

Supplementary Information accompanies this paper at <http://www.nature.com/naturecommunications>

Competing financial interests: The authors declare no competing financial interests.

Reprints and permission information is available online at <http://npg.nature.com/reprintsandpermissions/>

How to cite this article: Okazaki, Y. *et al.* A new class of plant lipid is essential for protection against phosphorus depletion. *Nat. Commun.* 4:1510 doi: 10.1038/ncomms2512 (2013).



This work is licensed under a Creative Commons Attribution-NonCommercial-ShareAlike 3.0 Unported License. To view a copy of this license, visit <http://creativecommons.org/licenses/by-nc-sa/3.0/>