ARTICLE

Received 14 Jul 2013 | Accepted 8 Oct 2013 | Published 14 Nov 2013

DOI: 10.1038/ncomms3726

Caspase-mediated activation of *Caenorhabditis elegans* CED-8 promotes apoptosis and phosphatidylserine externalization

Yu-Zen Chen^{1,2,*}, James Mapes^{1,*}, Eui-Seung Lee¹, Riley Robert Skeen-Gaar¹ & Ding Xue¹

During apoptosis, phosphatidylserine (PS), normally restricted to the inner leaflet of the plasma membrane, is exposed on the surface of apoptotic cells and serves as an 'eat-me' signal to trigger phagocytosis. It is poorly understood how PS exposure is activated in apoptotic cells. Here we report that CED-8, a *Caenorhabditis elegans* protein implicated in controlling the kinetics of apoptosis and a homologue of the XK family proteins, is a substrate of the CED-3 caspase. Cleavage of CED-8 by CED-3 activates its proapoptotic function and generates a carboxyl-terminal cleavage product, acCED-8, that promotes PS externalization in apoptotic cells and can induce ectopic PS exposure in living cells. Consistent with its role in promoting PS externalization in apoptotic cells, *ced-8* is important for cell corpse engulfment in *C. elegans*. Our finding identifies a crucial link between caspase activation and PS externalization, which triggers phagocytosis of apoptotic cells.

¹Department of Molecular, Cellular, and Developmental Biology, University of Colorado, Boulder, Colorado 80309, USA. ²Division of Biochemistry and Cellular Molecular Biology, Graduate Institute of Biomedical Sciences, Chang Gung University, Taoyuan, Taiwan 333, China. *These authors contributed equally to this work. Correspondence and requests for materials should be addressed to D.X. (email: Ding.Xue@Colorado.EDU).

Phospholipids are major components of the plasma membrane and are asymmetrically distributed in the inner and outer leaflets of the plasma membrane, with phosphatidylcholine and sphingomyelin predominantly in the outer leaflet and phosphatidylserine (PS) and phosphatidylethanolamine in the inner leaflet of the plasma membrane¹. During apoptosis, the asymmetrical distribution of phospholipids is disrupted, leading to the flipping of PS to the outer leaflet of the plasma membrane^{2,3}. The externalized PS then serves as an 'eat-me' signal to trigger removal of apoptotic cells⁴.

Caspases are a family of cysteine proteases that are activated during apoptosis to initiate multiple cell suicidal or disassembly events, including nuclear membrane breakdown, chromatin fragmentation and mitochondria elimination⁵. They promote cell death execution by cleaving and activating proapoptotic caspase substrates or inactivating anti-apoptotic caspase substrates⁶. How PS is externalized during apoptosis and whether caspases have a direct role in activating PS externalization are poorly understood and have been a long-standing issue in the field of apoptosis^{4,7}. In the nematode *Caenorhabditis elegans*, the caspase-encoding gene *ced-3* is required for the death of almost all cells programmed to die⁸, but how the CED-3 caspase kills the cell through cleavage of its downstream protein substrates remains largely unknown. DRP-1, a dynamin-related GTPase, and DCR-1, a ribonuclease important for small RNA biogenesis, have been identified as two important CED-3 substrates that are activated by CED-3 cleavage to promote mitochondrial elimination and to initiate chromosome fragmentation during apoptosis, respectively^{9,10}. As in mammals, PS externalization occurs during *C. elegans* apoptosis^{11–13} and is one of the earliest cell death events observed in worms¹⁴. In animals deficient in *ced-3*, few cells undergo apoptosis and PS is not externalized in cells that are programmed to die^{11,14}. These observations suggest that CED-3 is required to activate the PS externalization process in apoptotic cells.

The *ced-8* gene was first identified as a gene important for controlling the kinetics of apoptosis in *C. elegans*¹⁵. In *ced-8* loss-of-function (*lf*) mutants, the numbers of cell corpses in early embryonic stages are markedly reduced and more apoptotic cells appear in late embryonic stages and in the L1 larval stage, suggesting a delay of early embryonic cell deaths to later embryonic stages¹⁵. *ced-8* encodes a protein with sequence similarity to the human XK membrane transporter¹⁵. Loss of human XK has been associated with the McLeod syndrome, a hereditary disease characterized by abnormalities in the neuromuscular and hematopoietic systems, including acanthocytic red blood cells, loss of erythrocyte Kx antigen and reduction of Kell antigens^{16–18}. The mechanism by which *ced-8* affects apoptosis is unknown, but genetic analysis suggests that it functions downstream of or in parallel to the *C. elegans* cell death inhibitor *ced-9* (ref. 15).

In this study, we present evidence that CED-8 is a substrate of the CED-3 caspase and acts downstream of CED-3 to promote cell killing. Cleavage of CED-8 by CED-3 generates an activated cleavage product, acCED-8, that is necessary and sufficient to mediate the proapoptotic functions of CED-8. Moreover, we find that CED-8 is important for PS externalization in apoptotic cells and phagocytosis and that acCED-8 can induce ectopic PS exposure in living cells. Our study thus identifies an important caspase mediator that activates PS externalization in response to caspase activation.

Results

CED-8 is a substrate of the CED-3 caspase. In wild-type *C. elegans* embryos, most cell deaths occur during early

embryogenesis (comma, 1.5-fold and 2-fold embryonic stages) and less cell deaths are seen during late embryogenesis (3-fold and 4-fold embryonic stages; Fig. 1a). In a strong *lf ced-8(n1891)* mutant, the numbers of cell corpses in early embryonic stages are greatly reduced and significantly more apoptotic cells are found in late embryonic stages and in the L1 larval stage¹⁵, suggesting a delay of embryonic cell deaths (Fig. 1a). To understand how *ced-8* acts to effect apoptosis, we first determined whether *ced-8* acts downstream or upstream of the CED-3 caspase. We performed epistatic analysis of *ced-8* and *ced-3* using an integrated transgene (*smIs111*) that expresses an activated form of CED-3 (acCED-3) under the control of the promoter of the *C. elegans* cell death initiator *egl-1* (ref. 19). When *smIs111* was placed in the *ced-1(e1735); ced-3(n2433)* mutant background, in which almost all naturally occurring cell deaths in *C. elegans* are blocked by a strong *lf ced-3(n2433)* mutation (*ced-1(e1735)* blocks cell corpse engulfment and sensitizes quantification of apoptotic cells²⁰), acCED-3 still induced ectopic cell death and an average of 8 and 11 persistent cell corpses in 2-fold- and 4-fold-stage embryos, respectively (Fig. 1b). The *ced-8(n1891)* mutation significantly reduced the number of ectopic cell deaths induced by acCED-3 in *ced-1(e1735) smIs111; ced-3(n2433)* embryos (Fig. 1b), indicating that *ced-8* probably acts downstream of *ced-3* to promote apoptosis.

As effectors of apoptosis that act downstream of caspases are often substrates of caspases^{6,9,10,21}, we tested whether CED-8 is a substrate of the CED-3 protease. A glutathione S-transferase CED-8 fusion (GST-CED-8) was synthesized and labelled with ³⁵S-methionine in rabbit reticulocyte lysate and incubated with purified CED-3 protease²². ³⁵S-Met-labelled GST-CED-8 was cleaved by CED-3, yielding multiple cleavage products of ~51, 45, 34 and 28 kD (Fig. 1c, lanes 1 and 2). On the basis of the sizes of the cleavage products and the known consensus CED-3 cleavage sites, D/EXXD, in which X can be any amino acid²², we inferred the locations of two principle CED-3 cleavage sites: between Asp 21 and Ala 22 and between Asp 163 and Gly 164. The amino-acid sequences amino terminal to these two potential cleavage sites (P₄ to P₁; EQED and ENTD) are consistent with the consensus CED-3 cleavage sites. When Asp 21 of CED-8 was replaced by Ala and the resulting mutant protein, GST-CED-8(D21A) was examined in the CED-3 cleavage assay, the 51- and 28-kD cleavage products expected from CED-3 cleavage at CED-8 Asp 21 were lost (Fig. 1c, lanes 3 and 4), confirming that CED-3 cleaves CED-8 at Asp 21. Similarly, substitution of CED-8 Asp 163 with Ala blocked the production of the 45- and 34-kD cleavage products expected from CED-3 cleavage at CED-8 Asp 163 (Fig. 1c, lanes 5 and 6). Moreover, substitutions of both Asp 21 and Asp 163 with Ala prevented cleavage of CED-8 by CED-3 (Fig. 1c, lanes 7 and 8), indicating that these are two major *in vitro* CED-3 cleavage sites. Analysis of the CED-8 protein sequence using the TOPCONS program (consensus prediction of membrane protein topology) yields uniformed predictions that residues 1–34 are inside the cell, whereas residues 145–218 are extracellular and correspond to the site of the extracellular disulphide linkage of XK to Kell^{16,23}. Therefore, Asp 163 is unlikely accessible to CED-3 *in vivo*.

We examined whether CED-8 is cleaved by CED-3 in cells undergoing apoptosis using a yellow fluorescent protein (YFP) and cyan fluorescent protein (CFP) dual reporter assay. We generated a CFP fusion to the N terminus of wild-type CED-8 (CFP::CED-8) and a YFP fusion to the N terminus of CED-8(D21E) [YFP::CED-8(D21E)], which is resistant to CED-3 cleavage because D21E blocks CED-3 cleavage of CED-8 as well as D21A (Supplementary Fig. S1). Both fusion proteins were expressed in *C. elegans* under the control of the native *ced-8*

Figure 1 | *ced-8* acts downstream of *ced-3* and encodes a substrate of CED-3. (a) Appearance of apoptotic cell corpses in different stages of wild-type (N2) and *ced-8(n1891)* embryos. **(b)** *ced-8* acts downstream of *ced-3*. Cell corpses were scored in 2-fold and 4-fold *ced-1(e1735) smls111; ced-3(n2433)* and *ced-1(e1735) smls111; ced-3(n2433); ced-8(n1891)* embryos. **(c)** Cleavage of CED-8 by CED-3 *in vitro*. Top: ³⁵S-Met-labelled GST-CED-8 proteins (wild-type or mutants) were incubated with or without 5 ng purified CED-3 protease at 30 °C for 2 h and then resolved by 15% SDS-polyacrylamide gel electrophoresis. Bottom: a schematic diagram of GST-CED-8 and cleavage products generated by CED-3 (indicated by arrows). **(d)** Demonstration of *in vivo* cleavage of CED-8 by CED-3. CFP, YFP, CFP/YFP merged and differential contrast interference images of a comma-stage *ced-1(e1735)* embryo expressing CFP::CED-8 and YFP::CED-8(D21E) are shown. CFP (pseudo-colour in red) is observed in the cytoplasm of apoptotic cells (arrowheads). YFP (pseudo-colour in green) remains localized to the plasma membrane. Scale bar, 5 μm. **(e,f)** Rescue of the delay-of-cell-death defect of the *ced-8(n1891)* mutant by various *ced-8* transgenes. Cell corpses were scored in the following animals: **(e)** N2, *ced-8(n1891)*, *ced-8(n1891); ex[P_{ced-8}CED-8::GFP]#1* and *ced-8(n1891); ex[P_{ced-8}CED-8(D163A)::GFP]#1*. **(f)** N2, *ced-8(n1891)*, *ced-8(n1891); ex[P_{ced-8}CED-8(D21A)::GFP]#1* and *ced-8(n1891); ex[P_{ced-8}acCED-8::GFP]#1*. Comma- and 4-fold-stage transgenic embryos were examined. **(a,b,e,f)** The y-axis represents average number of cell corpses scored and error bars represent s.e.m. (*n* = 15). The significance of differences between results was determined by two-tailed Student's *t*-tests, **P* < 0.05; ***P* < 0.001.

promoter. In living cells, both CFP::CED-8 and YFP::CED-8(D21E) displayed identical plasma membrane localization indistinguishable from that of CED-8::GFP (green fluorescent protein) reported previously (Fig. 1d and Supplementary Fig. S2a,b)¹⁵. In cells undergoing apoptosis (arrowheads, Fig. 1d), CFP was observed instead in the cytoplasm, suggesting that following CED-3 cleavage CFP disassociated from CED-8 localized on the plasma membrane and diffused into the

cytoplasm. In comparison, YFP remained associated with CED-8(D21E) and localized to the plasma membrane in these apoptotic cells (Fig. 1d), indicating that CED-3 cleaves CED-8 at Asp 21 *in vivo*.

CED-8 cleavage by CED-3 is critical for apoptosis. We next examined whether the two *in vitro* CED-3 cleavage sites in CED-8

are important for its proapoptotic function *in vivo*. We tested whether transgenes expressing wild type or mutant forms of CED-8::GFP fusions under the control of the *ced-8* gene promoter can rescue the delay-of-cell-death defect of the *ced-8(n1891)* mutant at two signature embryonic stages, the early comma embryonic stage and the late 4-fold embryonic stage (Fig. 1a). Compared with wild-type embryos, which had an average of 9 cell corpses in the comma stage and few cell corpse in the 4-fold stage, *ced-8(n1891)* embryos had 1–2 cell corpses in the comma stage and an average of 15 cell corpses in the 4-fold embryonic stage (Fig. 1a)¹⁵. Transgenes expressing wild-type CED-8 [*P_{ced-8}*CED-8::GFP] or mutant CED-8(D163A) [*P_{ced-8}*CED-8(D163A)::GFP] fully rescued the delay-of-cell-death defect of the *ced-8(n1891)* mutant and restored wild-type cell corpse patterns to the *ced-8(n1891)* embryos (Fig. 1e), indicating that Asp 163 is not important for CED-8's proapoptotic function. By contrast, *P_{ced-8}*CED-8(D21A)::GFP did not rescue the delay-of-cell-death defect of the *ced-8(n1891)* mutant (Fig. 1f), although the GFP intensity on the plasma membrane of *P_{ced-8}*CED-8(D21A)::GFP transgenic embryos was similar to that of *P_{ced-8}*CED-8::GFP or *P_{ced-8}*CED-8(D163A)::GFP embryos (Supplementary Fig. S2). This result indicates that CED-3 cleavage at the Asp 21 site is critical for CED-8's proapoptotic function *in vivo*. Finally, expression of the carboxy-terminal cleavage fragment of CED-8 by CED-3 [*P_{ced-8}*CED-8(22–458)::GFP] also fully rescued the *ced-8(n1891)* mutant (Fig. 1f), indicating that CED-8(22–458), which we named acCED-8, is sufficient to mediate the proapoptotic function of CED-8 and represents the CED-3-activated form of CED-8.

CED-8 is crucial for PS externalization in apoptotic cells. To probe how *ced-8* regulates apoptosis, we examined whether *ced-8* affects the PS externalization process, a hallmark of apoptosis^{1,2,7}, and one of the earliest detectable cell death events in *C. elegans*¹⁴. We used a genetically encoded PS sensor, secreted Annexin V::GFP fusion (ANV::GFP)¹⁴, to monitor the PS externalization process. In early comma-stage wild-type *C. elegans* embryos carrying an integrated transgene *smIs76*, which expresses ANV::GFP under the control of the *C. elegans* heat-shock promoters¹⁴, an average of nine cell corpses were observed (Fig. 1a) and almost all of them were strongly labelled by ANV::GFP (Fig. 2a,d). In comma-stage *smIs76; ced-8(n1891)* embryos, only a few cell corpses were observed and ~60% of them were labelled by ANV::GFP (Figs 1a and 2b–d). However, in those apoptotic cells that were labelled, ANV::GFP labelling often was weak or barely detectable (Fig. 2c,e), with an average fluorescent intensity ~11% of that seen in apoptotic cells in wild-type embryos (Fig. 2e). These results indicate that loss of *ced-8* blocks or greatly reduces PS externalization in cells that already died at this stage. As the death of most cells programmed to die at the comma stage was deferred to later embryonic stages, we examined whether apoptotic cells observed in late-stage *ced-8(n1891)* embryos were labelled by ANV::GFP. In 4-fold *ced-8(n1891)* embryos, there are an average of 15 cell corpses compared with an average of 0.5 cell corpse in wild-type 4-fold embryos (Fig. 1a). However, only 5% of these apoptotic cells in *smIs76; ced-8(n1891)* 4-fold embryos were labelled by ANV::GFP (Fig. 2d), indicating that PS was not externalized in cells that died in late-stage *ced-8(n1891)* embryos either. We obtained similar results using another genetically encoded PS sensor, secreted GFP::Lactadherin fusion (GFP::Lact)¹⁴. In *smIs434* animals, which carry an integrated transgene expressing GFP::Lact under the control of the heat-shock promoters¹⁴, *ced-8(n1891)* not only reduced the percentage of apoptotic cells labelled by GFP::Lact (Supplementary Fig. S3a) but also markedly decreased the

intensity of GFP::Lact labelling on apoptotic cells (Supplementary Fig. S3b), providing further evidence that loss of *ced-8* greatly reduces PS externalization in apoptotic cells.

We then performed time-lapse microscopy to follow the PS externalization process in cells undergoing apoptosis. As reported previously¹⁴, in wild-type *C. elegans* embryos, ANV::GFP was detected on the surface of the dying cell before it adopted the typical button-like cell corpse morphology (Supplementary Fig. S4a,b). The intensity of ANV::GFP increased as the dying cell shrank and displayed the dense button-like morphology (Supplementary Fig. S4c), and reached its maximum around the time when it was engulfed by one of its neighbours (Supplementary Fig. S4d), which were also weakly labelled by ANV::GFP¹⁴. In the *ced-8(n1891)* embryo, ANV::GFP labelling of the apoptotic cell was much weaker (Supplementary Fig. S4h–n) and was first detected after the dying cell already adopted the button-like morphology (Supplementary Fig. S4h,i), the appearance of which was delayed compared with that in wild-type embryos (20 versus 10 min in the N2 embryo; Supplementary Fig. S4c,i). The intensity of ANV::GFP on the apoptotic cell did not increase subsequently and the apoptotic cell persisted for a significantly longer time period (120 versus 30 min in the N2 embryo; Supplementary Fig. S4c–g, i–n), which suggests a potential defect in cell corpse engulfment (see below). The observations from the time-lapse microscopy provide further support that *ced-8* has a critical role in PS externalization in apoptotic cells.

It is possible that in *ced-8(n1891)* animals, the cells programmed to die in early embryonic stages already initiated the suicide programme and exposed PS, but had not yet adopted the button-like morphology. We thus quantified the total number of ANV::GFP rings in comma-stage embryos regardless of their morphology and found that wild-type embryos had an average of 12 ANV::GFP rings, most of which displayed apoptotic cell morphology (Fig. 2f). By contrast, *ced-8(n1891)* comma-stage embryos had an average of 1.5 ANV::GFP rings (Fig. 2f), confirming that most cells programmed to die at the comma stage did not expose PS at this stage. We further sensitized quantification of PS exposure in apoptotic cells by inactivating the *ced-7* gene, which encodes an ATP-binding cassette transporter important for cell corpse engulfment²⁴ and promotes time-dependent efflux of exposed PS from the surface of apoptotic cells to the extracellular space through PS-containing small vesicles^{14,25}. Loss of *ced-7* leads to retention of exposed PS on the surface of aged or unengulfed apoptotic cells and thus sensitizes scoring of apoptotic cells with surface-exposed PS. In comma-stage *ced-7(n2094)* embryos, almost all apoptotic cells (97%) were labelled by ANV::GFP, whereas in 4-fold *ced-7(n2094)* embryos, 52% of unengulfed cell corpses still had surface-exposed PS (Fig. 2d). In comparison, only 6% of unengulfed cell corpses in *ced-1(e1735)* 4-fold embryos have surface-exposed PS because of efflux and loss of exposed PS from apoptotic cells¹⁴. In *ced-7(n2094); ced-8(n1891)* embryos, 63% of cell corpses in the comma stage were labelled by ANV::GFP and only 11% of unengulfed cell corpses were labelled by ANV::GFP in late 4-fold embryos (Fig. 2d), which had the same number of unengulfed apoptotic cells as that in *ced-7(n2094)* 4-fold embryos (Fig. 4c). These observations confirm that loss of *ced-8* also significantly reduces PS exposure in unengulfed apoptotic cells. Taken together, these results indicate that *ced-8* is crucial for PS externalization in apoptotic cells.

CED-8 is activated by CED-3 to promote PS externalization.

We next examined whether CED-3 cleavage of CED-8 at the Asp 21 site is critical for CED-8 to promote PS externalization in

Figure 2 | PS externalization in apoptotic cells is blocked or greatly reduced in the *ced-8* mutant. (a–c) ANV::GFP and differential contrast interference images of comma-stage embryos from *smls76* and *smls76; ced-8(n1891)* animals are shown. Arrowheads indicate apoptotic cells labelled by ANV::GFP. In *smls76; ced-8(n1891)* embryos, apoptotic cells were either not labelled (indicated by an arrow, **b**) or were very weakly labelled (**c**) by ANV::GFP. Insets show enhanced GFP images of apoptotic cells. Scale bars, 5 μ m. **(d)** Percentages of apoptotic cells labelled by ANV::GFP in the indicated strains and stages are shown. **(e)** Relative intensity of ANV::GFP that labelled apoptotic cells in comma-stage embryos was quantified in the indicated strains (see Methods). Results are shown using the histogram ($n = 30$), with mean $\pm$ s.e.m. indicated. Apoptotic cells that were not labelled by ANV::GFP are presented with zero intensity. **(f)** The number of ANV::GFP rings in the indicated strains was scored in the comma- and 4-fold-stage embryos regardless of their cell morphology. **(g)** Rescue of the PS externalization defect of the *ced-8(n1891)* mutant by various *ced-8* transgenes. Quantification of the percentage of apoptotic cells labelled by ANV::GFP and the relative intensity of ANV::GFP in transgenic embryos was performed as in **d** and **e**. **(d,f,g)** Results are presented as mean $\pm$ s.e.m. ($n = 15$ each). The significance of differences between results was determined by two-tailed Student's *t*-tests, * $P < 0.05$; ** $P < 0.001$. Heat-shock treatment was performed at 33°C for 30 min and embryos were examined after 2 h recovery at 20 °C. Exposure time for each fluorescent image was identical (12.5 ms).

apoptotic cells. Transgenes expressing wild-type or altered forms of CED-8 under the control of the *ced-8* gene promoter were introduced into *smls76; ced-8(n1891)* animals. The percentage of apoptotic cells labelled by ANV::GFP and the intensity of ANV::GFP labelling in comma-stage transgenic embryos were examined (Fig. 2g). Expression of wild-type CED-8 or acCED-8 in *smls76; ced-8(n1891)* embryos restored ANV::GFP labelling of apoptotic cells to the levels seen in *smls76* embryos (Fig. 2g), indicating full rescue of the PS externalization defect. By contrast,

ANV::GFP labelling of apoptotic cells in *smls76; ced-8(n1891)* embryos expressing CED-8(D21E) was similar to that seen in *smls76; ced-8(n1891)* embryos (Fig. 2g), indicating no rescue of the PS externalization defect. These results indicate that CED-3 cleavage at the Asp 21 site is also critical for activating CED-8's function in promoting PS externalization in apoptotic cells.

Given that acCED-8 is necessary and sufficient to mediate the PS externalization activity of CED-8 in apoptotic cells, we examined whether acCED-8 can induce PS exposure in non-

apoptotic cells. We induced global expression of acCED-8 under the control of the *C. elegans* heat-shock promoters (P_{hsp} acCED-8) and then examined PS exposure using GFP::Lact, which is more sensitive in detecting PS exposure in living cells than ANV::GFP (Supplementary Fig. S5)^{14,26}. In *smIs434* animals expressing GFP::Lact, GFP::Lact only labelled apoptotic cells (arrowheads, Fig. 3a). Expression of acCED-8 in *smIs434* embryos resulted in GFP::Lact labelling of all cells, including both living cells and apoptotic cells (Fig. 3c), suggesting that acCED-8 can induce ectopic PS exposure on the surface of all cells. In comparison, expression of full-length CED-8 (P_{hsp} CED-8) in *smIs434* embryos did not alter the staining pattern of GFP::Lact or cause widespread, ectopic PS exposure on the surface of all cells (Fig. 3b). In *ced-3(n2433); smIs434* animals, where almost all apoptosis is blocked, expression of acCED-8, but not full-length CED-8, still caused all cells to externalize PS (Fig. 3d,e), indicating that acCED-8 is capable of activating PS externalization independent of the CED-3 caspase and apoptosis. These results suggest that CED-8, once activated by CED-3, promotes PS externalization in cells through the generation of acCED-8.

Figure 3 | Expression of acCED-8 induces ectopic PS exposure in all cells.

(a–e) GFP::Lact and differential contrast interference images of comma-stage embryos from the following strains are shown: *smIs434* (a), *smIs434; ex[P_{hsp}CED-8]* (b), *smIs434; ex[P_{hsp}acCED-8]* (c), *ced-3(n2433); smIs434; ex[P_{hsp}CED-8]* (d) and *ced-3(n2433); smIs434; ex[P_{hsp}acCED-8]* (e).

Arrowheads indicate apoptotic cells. Heat-shock treatment was performed at 33°C for 45 min and embryos were examined after 2 h recovery at 20 °C. Exposure time for each fluorescent image was identical (100 ms). Scale bars, 5 μm.

CED-8 is important for cell corpse engulfment. As surface-exposed PS is an important ‘eat-me’ signal for phagocytosis^{2,4,27} and because cell corpses in *ced-8(n1891)* embryos appear to persist significantly longer than those in wild-type embryos (Supplementary Fig. S4), we investigated whether *ced-8* affects cell corpse engulfment in *C. elegans*, in addition to its previously reported role in regulating the kinetics of cell death¹⁵. We first used four-dimensional microscopy to measure the duration of embryonic cell corpses in *ced-8(n1891)* animals²⁸. In wild-type embryos, on average cell corpses persisted for 34 min. In *ced-8(n1891)* embryos, cell corpses persisted for 96 min, which is almost three times longer than that of wild-type animals (Fig. 4a). These results indicate that the cell corpse engulfment process is significantly compromised in the *ced-8(n1891)* mutant.

In *C. elegans*, two partially redundant pathways mediate cell corpse removal, with *ced-1*, *ced-6*, *ced-7* and *ttr-52* genes functioning in one pathway and *ced-2*, *ced-5* and *ced-12* genes acting in the other^{29–31}. To interrogate the functioning pathway of *ced-8*, we constructed and analysed double mutants between the *ced-8(n1891)* mutation and strong *lf* alleles of the above engulfment genes. *ced-8(n1891)* markedly enhanced the cell corpse engulfment defect of the *ced-2*, *ced-5* or *ced-12* mutants at the late 4-fold embryonic stage and L1 larval stage (Fig. 4b), but only mildly enhanced the corpse engulfment defect of the *ced-1*, *ced-6*, *ced-7* or *ttr-52* mutants at the L1 larval stage (Fig. 4c). These results are consistent with *ced-8* acting to promote PS externalization, which serves as an ‘eat-me’ signal for both cell corpse engulfment pathways to promote phagocytosis of apoptotic cells (Fig. 5)^{26,28,31}.

Discussion

As a multipass plasma membrane protein, how CED-8 affects programmed cell death has been a major unresolved issue in *C. elegans* apoptosis. In this study, we demonstrate that CED-8 acts downstream of the CED-3 caspase to promote apoptosis and is a substrate of CED-3. Cleavage of CED-8 by CED-3 is required to activate CED-8’s proapoptotic functions and generates a C-terminal cleavage product, acCED-8, that is sufficient to mediate the proapoptotic functions of CED-8. Importantly, *ced-8* is also critical for PS externalization in apoptotic cells, which is an unexpected new role for CED-8. Cleavage of CED-8 by CED-3 is also required to activate its PS externalization activity. Moreover, ectopic expression of acCED-8, but not full-length CED-8, can induce PS externalization in all living cells in the absence of the CED-3 caspase or apoptosis. Our study thus identifies a crucial link between caspase activation and PS externalization^{4,27}.

Consistent with its role in promoting PS externalization in apoptotic cells, we find that *ced-8* is important for cell corpse engulfment. Loss of *ced-8* causes persistence of embryonic cell corpses, mildly enhances the cell corpse engulfment defects of *ced-1*, *ced-6*, *ced-7* and *ttr-52* mutants and strongly enhances the cell corpse engulfment defects of *ced-2*, *ced-5* and *ced-12* mutants. As PS could be an ‘eat-me’ signal for both cell corpse engulfment pathways in *C. elegans*^{26,28,31}, our observations are consistent with *ced-8* acting in both engulfment pathways to promote clearance of apoptotic cells (Fig. 5). However, given the significantly stronger effect of *ced-8(n1891)* in enhancing the engulfment defect of the *ced-2* pathway and marginal effect in enhancing the engulfment defect of the *ced-1* pathway, *ced-8* may mainly act through the *ced-1*, *ced-6*, *ced-7* and *ttr-52* engulfment pathway to promote phagocytosis. As *ced-1* encodes a phagocyte receptor that recognizes and promotes engulfment of apoptotic cells with surface-exposed PS through its binding partner TTR-52, a PS-binding extracellular bridging protein^{31,32}, our

Figure 4 | *ced-8* is important for cell corpse engulfment in *C. elegans*. (a) Time-lapse microscopy analysis of the durations of cell corpses. The persistence of 25 cell corpses from N2 embryos ($n=3$) and 24 cell corpses from *ced-8(n1891)* embryos ($n=8$) was monitored. The numbers in parentheses indicate the average persistence for cell corpses $\pm$ s.e.m. The y-axis indicates the number of cell corpses with a specific duration range (shown by the x-axis). The durations of four cell divisions in the MS cell lineage from the MS cell to the MS.aaaa cell³⁸ were also followed to ensure that the embryos assayed had similar rates of development. The average duration of four cell divisions for N2 embryos is 89 ± 6 and 91 ± 3 min for *ced-8(n1891)* embryos. (b,c) The *ced-8(n1891)* mutation enhances the corpse clearance defect of engulfment mutants. Cell corpses were scored in 4-fold embryos and in L1 larvae from the indicated strains. The y-axis represents the mean of cell corpses scored in the whole embryos or in the head region of L1 larvae ($n=15$). Error bar represents s.e.m. The significance of differences between results was determined by two-tailed Student's *t*-tests, * $P<0.05$; ** $P<0.001$.

finding delineates a new apoptotic signalling cascade, starting from CED-3 caspase activation during apoptosis, followed by CED-3-mediated activation of CED-8, acCED-8-mediated activation of PS exposure on the surface of apoptotic cells, and subsequent recognition and removal of PS-exposed apoptotic cells by the TTR-52 and CED-1 phagocytosis pathway, which is probably aided by the *ced-2*, *ced-5* and *ced-12* phagocytosis pathway, also capable of recognizing and engulfing apoptotic cells with surface-exposed PS (Fig. 5).

A new study reports a similar finding that a mammalian XK family protein, Xkr8, and its *C. elegans* homologue CED-8 have a conserved role in promoting PS exposure in apoptotic cells³³. The PS-externalizing activity of Xkr8 also appears to be activated by caspase cleavage. Interestingly, CED-8 is cleaved and activated by the CED-3 caspase at the N terminus, whereas Xkr8 is cleaved

and potentially activated by mammalian caspases at its C terminus (Supplementary Fig. S6). Although the activated form of Xkr8 has not been identified³³, the N-terminal cleavage product of Xkr8 shares an overall sequence similarity to acCED-8, including all putative transmembrane domains. Moreover, the 21-residue CED-8 N-terminal sequence removed by CED-3 cleavage seems to be unique to *C. elegans* and is not found in Xkr8 or Xkr8 homologues in other organisms. It remains to be determined whether CED-8 and Xkr8 are regulated by similar factors and activated through a conserved mechanism, and whether activated CED-8 and activated Xkr8 promote PS externalization in apoptotic cells through a similar mechanism.

CED-8 shares sequence similarity to the XK transporters. However, the substrates that XK family proteins might transport remain unknown^{18,33}. Given our new findings, it is conceivable

Figure 5 | CED-8 working model. CED-8 is cleaved and activated by CED-3 during apoptosis to generate acCED-8, which promotes PS externalization and normal progression of apoptosis through undefined mechanisms (?). Surface-exposed PS then promotes phagocytosis of apoptotic cells through TTR-52, a PS-binding extracellular bridging molecule, and its binding partner, the CED-1 phagocyte receptor. Surface-exposed PS may also promote phagocytosis of apoptotic cells through the CED-2/CED-5/CED-12 engulfment pathway.

that acCED-8 may directly transport PS across the plasma membrane to effect PS exposure. However, because CED-8 also affects the kinetics of apoptosis, it is more likely that acCED-8 is a general caspase mediator that interacts with several downstream effectors to promote cell death execution (Fig. 5). In this scenario, acCED-8 may regulate the activity of a lipid-transporting enzyme, leading to PS externalization in apoptotic cells. Moreover, because loss of *ced-8* does not completely abolish PS externalization in *C. elegans* apoptotic cells, additional mechanisms or regulators must be responsible for the residual PS externalization activity in the *ced-8* mutant during apoptosis.

Methods

Strains and culture conditions. *C. elegans* strains were maintained using standard procedures³⁴. The following alleles were used in the genetic analyses: LGI, *ced-1(e1735)*, *ced-12(n3261)*, *smIs111*; LGIII, *ced-6(n2095)*, *ced-7(n2094)*, *ttr-52(tm2078)*; LGIV, *ced-2(n1994)*, *ced-3(n2433)*, *ced-5(n1812)*; LGV, *smIs76*; and LGX, *ced-8(n1891)*, *smIs434*. *smIs76* and *smIs434* are integrated transgenes generated by ultraviolet irradiation of animals carrying extrachromosomal arrays containing *P_{hsp}ANV::GFP* and *P_{hsp}GFP::Lact*, respectively¹⁴. They were backcrossed four times with wild-type animals before being used for genetic analyses.

Transgenic animals. Transgenic animals were generated using standard protocols³⁵. *P_{ced-8}CED-8::GFP*, *P_{ced-8}CED-8(D21A)::GFP*, *P_{ced-8}CED-8(D163A)::GFP* or *P_{ced-8}acCED-8::GFP* was injected into *unc-76(e911)*; *ced-8(n1891)* animals at 5 ng μl^{-1} along with an *unc-76* rescuing plasmid (at 50 ng μl^{-1}) as a co-injection marker. *P_{ced-8}CFP::CED-8* and *P_{ced-8}YFP::CED-8(D21E)* were co-injected into *ced-1(e1375)*; *unc-76(e911)* animals at 25 ng μl^{-1} each along with the *unc-76* rescuing plasmid (at 50 ng μl^{-1}). *P_{hsp}CED-8* or *P_{hsp}acCED-8* was injected into *smIs434* or *ced-3(n2433)*; *smIs434* animals at 25 ng μl^{-1} along with the *P_{sur}-mCherry* plasmid (25 ng μl^{-1}) as a co-injection marker, which directs mCherry expression in all somatic cells in most developmental stages³⁶.

Quantification of cell corpses. The number of cell corpses in living embryos or in the head region of L1 larvae was scored using Nomarski optics³⁷.

Quantification of ANV::GFP or GFP::Lact intensity on corpses. To score the number of cell corpses labelled by ANV::GFP or GFP::Lact, apoptotic cells were first identified using differential contrast interference optics and then examined for the presence of ANV::GFP or GFP::Lact on their surface. To measure the ANV::GFP or GFP::Lact fluorescence intensity on cell corpses, fluorescence micrographs were captured using a Zeiss Axioplan 2 microscope (Zeiss) equipped with a charge-coupled device camera and the Slidebook 5.0 software (Intelligent Imaging Innovations). GFP intensity was determined by measuring the fluorescence signals from three random areas on the plasma membrane of each apoptotic cell (1a) and three random areas on normal cells (1b). The average value of 1a minus the average value of 1b was then used as the value for relative ANV::GFP or GFP::Lact intensity.

Quantification of ANV::GFP rings. *C. elegans* embryos were mounted on slides using the egg salt buffer (118 mM NaCl, 48 mM KCl, 2 mM CaCl_2 , 2 mM MgCl_2 , 5 mM HEPES pH 7.5) and the number of ANV::GFP rings was then scored regardless of the cell morphology.

Heat-shock treatment. For time-lapse microscopy, *smIs76* or *smIs76*; *ced-8(n1891)* embryos were placed at 33°C for 30 min, followed by recovery

at room temperature for 1 h. For analysis of ANV::GFP or GFP::Lact labelling of comma-stage embryos, embryos were placed at 33°C for 45 min, followed by recovery at room temperature 2 h. For analysis of ANV::GFP or GFP::Lact labelling of 4-fold-stage embryos, embryos were placed at 33°C for 45 min, followed by recovery at room temperature for 3 h.

Microscopy. For fluorescent microscopy, *C. elegans* embryos were mounted on agar pads and observed using $\times 100$ optics on Axioplan 2 microscopes (Zeiss). Fluorescence and differential contrast interference images were captured using a charge-coupled device camera (PCO SensiCam) coupled with the Slidebook 5.0 software.

To perform time-lapse microscopy, *C. elegans* embryos were mounted on slides with agar pad in egg salt buffer and cover slips were sealed with Beeswax and Vaseline (1:1). Images in a 45-micron z-series (3 micron per section) were captured every 2 or 5 min for 180–360 min. Neutral density filter was used to allow only 12% light transmission. Four-dimensional microscopy analysis of cell corpse durations in the N2 and *ced-8(n1891)* embryos were performed as described previously²⁸.

Molecular biology. *ced-8*-coding region was amplified using PCR from the pBluescript KS(+)–*ced-8* vector¹⁵ and subcloned into the pET-41b vector (Novagen) through its *SpeI* and *EcoRV* sites. Deletions of CED-8 or single amino-acid substitutions in CED-8 were generated by site-directed mutagenesis of the resulting pET-41b–CED-8 vector. The *P_{ced-8}CED-8::GFP* vector was made by fusing a 1.2-kb (*SmaI/NheI*) *ced-8* promoter to a *ced-8* cDNA (*NheI/AgeI*) in frame with GFP from pPD95.77. The *P_{ced-8}CFP::CED-8* and *P_{ced-8}YFP::CED-8(D21E)* vectors were made by fusing the 1.2 kb *ced-8* promoter to CFP::CED-8 and YFP::CED-8(D21E), respectively. *P_{hsp}acCED-8* was made by PCR amplification of *ced-8* cDNA encoding amino acids 22–458 and subcloning the PCR fragment into the pPD49.78 and pPD49.83 vectors through *NheI* and *EcoR V* sites.

CED-3 protease assays. CED-3 protease assays were carried out using the standard protocol²². Briefly, GST–CED-8, GST–CED-8(D21A), GST–CED-8(D21E), GST–CED-8(D163A) and GST–CED-8(D21A, D163A) were synthesized and labelled with ³⁵S-methionine using TNT Transcription/Translation coupled system (Promega) and incubated with 5 ng of purified CED-3 at 30°C for 2 h. The reactions were then resolved by 15% SDS–PAGE.

Statistical analysis. *P*-values represent analysis from two-tailed Student's *t*-tests. **P* < 0.05; ***P* < 0.001.

References

- Daleke, D. L. Regulation of transbilayer plasma membrane phospholipid asymmetry. *J. Lipid Res.* **44**, 233–242 (2003).
- Fadok, V. A. *et al.* Exposure of phosphatidylserine on the surface of apoptotic lymphocytes triggers specific recognition and removal by macrophages. *J. Immunol.* **148**, 2207–2216 (1992).
- Fadok, V. A. *et al.* Different populations of macrophages use either the vitronectin receptor or the phosphatidylserine receptor to recognize and remove apoptotic cells. *J. Immunol.* **149**, 4029–4035 (1992).
- Savill, J. & Fadok, V. Corpse clearance defines the meaning of cell death. *Nature* **407**, 784–788 (2000).
- Steller, H. Mechanisms and genes of cellular suicide. *Science* **267**, 1445–1449 (1995).
- Boyce, M., Degterev, A. & Yuan, J. Caspases: an ancient cellular sword of Damocles. *Cell Death Differ.* **11**, 29–37 (2004).
- Fadok, B. & Xue, D. The ins and outs of phospholipid asymmetry in the plasma membrane: roles in health and disease. *Crit. Rev. Biochem. Mol. Biol.* **44**, 264–277 (2009).
- Ellis, H. M. & Horvitz, H. R. Genetic control of programmed cell death in the nematode *C. elegans*. *Cell* **44**, 817–829 (1986).
- Breckenridge, D. G. *et al.* *Caenorhabditis elegans drp-1* and *fis-2* regulate distinct cell-death execution pathways downstream of *ced-3* and independent of *ced-9*. *Mol. Cell* **31**, 586–597 (2008).
- Nakagawa, A., Shi, Y., Kage-Nakadai, E., Mitani, S. & Xue, D. Caspase-dependent conversion of Dicer ribonuclease into a death-promoting deoxyribonuclease. *Science* **328**, 327–334 (2010).
- Wang, X. *et al.* *C. elegans* mitochondrial factor WAH-1 promotes phosphatidylserine externalization in apoptotic cells through phospholipid scramblase SCRM-1. *Nat. Cell Biol.* **9**, 541–549 (2007).
- Zullig, S. *et al.* Aminophospholipid translocase TAT-1 promotes phosphatidylserine exposure during *C. elegans* apoptosis. *Curr. Biol.* **17**, 994–999 (2007).
- Venegas, V. & Zhou, Z. Two alternative mechanisms that regulate the presentation of apoptotic cell engulfment signal in *Caenorhabditis elegans*. *Mol. Biol. Cell* **18**, 3180–3192 (2007).
- Mapes, J. *et al.* CED-1, CED-7, and TTR-52 regulate surface phosphatidylserine expression on apoptotic and phagocytic cells. *Curr. Biol.* **22**, 1267–1275 (2012).

15. Stanfield, G. M. & Horvitz, H. R. The *ced-8* gene controls the timing of programmed cell deaths in *C. elegans*. *Mol. Cell* **5**, 423–433 (2000).
16. Khamlichi, S. *et al.* Purification and partial characterization of the erythrocyte Kx protein deficient in McLeod patients. *Eur. J. Biochem.* **228**, 931–934 (1995).
17. Ho, M. *et al.* Isolation of the gene for McLeod syndrome that encodes a novel membrane transport protein. *Cell* **77**, 869–880 (1994).
18. Danek, A. *et al.* Neuroacanthocytosis: new developments in a neglected group of dementing disorders. *J. Neurol. Sci.* **229–230**, 171–186 (2005).
19. Kokel, D., Li, Y., Qin, J. & Xue, D. The nongenotoxic carcinogens naphthalene and para-dichlorobenzene suppress apoptosis in *Caenorhabditis elegans*. *Nat. Chem. Biol.* **2**, 338–345 (2006).
20. Hedgecock, E. M., Sulston, J. E. & Thomson, J. N. Mutations affecting programmed cell deaths in the nematode *Caenorhabditis elegans*. *Science* **220**, 1277–1279 (1983).
21. Nicholson, D. W. & Thornberry, N. A. Caspases: killer proteases. *Trends Biochem. Sci.* **22**, 299–306 (1997).
22. Xue, D., Shaham, S. & Horvitz, H. R. The *Caenorhabditis elegans* cell-death protein CED-3 is a cysteine protease with substrate specificities similar to those of the human CPP32 protease. *Genes Dev.* **10**, 1073–1083 (1996).
23. Russo, D., Redman, C. & Lee, S. Association of XK and Kell blood group proteins. *J. Biol. Chem.* **273**, 13950–13956 (1998).
24. Wu, Y. C. & Horvitz, H. R. The *C. elegans* cell corpse engulfment gene *ced-7* encodes a protein similar to ABC transporters. *Cell* **93**, 951–960 (1998).
25. Zhang, Y., Wang, H., Kage-Nakadai, E., Mitani, S. & Wang, X. *C. elegans* secreted lipid-binding protein NRF-5 mediates PS appearance on phagocytes for cell corpse engulfment. *Curr. Biol.* **22**, 1276–1284 (2012).
26. Darland-Ransom, M. *et al.* Role of *C. elegans* TAT-1 protein in maintaining plasma membrane phosphatidylserine asymmetry. *Science* **320**, 528–531 (2008).
27. Fadok, V. A., Bratton, D. L., Frasch, S. C., Warner, M. L. & Henson, P. M. The role of phosphatidylserine in recognition of apoptotic cells by phagocytes. *Cell. Death Differ.* **5**, 551–562 (1998).
28. Wang, X. *et al.* Cell corpse engulfment mediated by *C. elegans* phosphatidylserine receptor through CED-5 and CED-12. *Science* **302**, 1563–1566 (2003).
29. Ellis, R. E., Jacobson, D. M. & Horvitz, H. R. Genes required for the engulfment of cell corpses during programmed cell death in *Caenorhabditis elegans*. *Genetics* **129**, 79–94 (1991).
30. Reddien, P. W. & Horvitz, H. R. The engulfment process of programmed cell death in *Caenorhabditis elegans*. *Annu. Rev. Cell Dev. Biol.* **20**, 193–221 (2004).
31. Wang, X. *et al.* *Caenorhabditis elegans* transthyretin-like protein TTR-52 mediates recognition of apoptotic cells by the CED-1 phagocyte receptor. *Nat. Cell Biol.* **12**, 655–664 (2010).
32. Zhou, Z., Hartwig, E. & Horvitz, H. R. CED-1 is a transmembrane receptor that mediates cell corpse engulfment in *C. elegans*. *Cell* **104**, 43–56 (2001).
33. Suzuki, J., Denning, D. P., Imanishi, E., Horvitz, H. R. & Nagata, S. Xk-related protein 8 and CED-8 promote phosphatidylserine exposure in apoptotic cells. *Science* **341**, 403–406 (2013).
34. Brenner, S. The genetics of *Caenorhabditis elegans*. *Genetics* **77**, 71–94 (1974).
35. Mello, C. C., Krame, J. M., Stinchcomb, D. & Ambros, V. Efficient gene transfer in *C. elegans*: Extrachromosomal maintenance and integration of transforming sequences. *EMBO J.* **10**, 3959 (1992).
36. Gu, T., Orita, S. & Han, M. *Caenorhabditis elegans* SUR-5, a novel but conserved protein, negatively regulates LET-60 Ras activity during vulval induction. *Mol. Cell Biol.* **18**, 4556–4564 (1998).
37. Parrish, J. *et al.* Mitochondrial endonuclease G is important for apoptosis in *C. elegans*. *Nature* **412**, 90–94 (2001).
38. Sulston, J. E., Schierenberg, E., White, J. G. & Thomson, J. N. The embryonic cell lineage of the nematode *Caenorhabditis elegans*. *Dev. Biol.* **100**, 64–119 (1983).

Acknowledgements

We thank members of the Xue lab for helpful discussions. This work was supported by US National Institutes of Health Grants (R01 GM59083, R01 GM79097 and R01 GM088241) to D.X.

Author contributions

Y.Z.C., J.M., E.S.L. and R.R.S. performed the experiments; Y.Z.C., J.M. and D.X. conceived and designed the experiments, analysed and interpreted data, and wrote the manuscript.

Additional information

Supplementary Information accompanies this paper at <http://www.nature.com/naturecommunications>

Competing financial interests: The authors declare no competing financial interests.

Reprints and permission information is available online at <http://npg.nature.com/reprintsandpermissions/>

How to cite this article: Chen, Y.-Z. *et al.* Caspase-mediated activation of *Caenorhabditis elegans* CED-8 promotes apoptosis and phosphatidylserine externalization. *Nat. Commun.* 4:2726 doi: 10.1038/ncomms3726 (2013).