

ARTICLE

Received 11 Sep 2013 | Accepted 1 Nov 2013 | Published 2 Dec 2013

DOI: 10.1038/ncomms3856

Discovery of chlamydial peptidoglycan reveals bacteria with murein sacculi but without FtsZ

Martin Pilhofer^{1,2,*}, Karin Aistleitner^{3,*}, Jacob Biboy⁴, Joe Gray⁵, Erkin Kuru⁶, Edward Hall⁶, Yves V. Brun⁶, Michael S. VanNieuwenhze⁶, Waldemar Vollmer⁴, Matthias Horn³ & Grant J. Jensen^{1,2}

Chlamydiae are important pathogens and symbionts with unique cell biological features. They lack the cell-division protein FtsZ, and the existence of peptidoglycan (PG) in their cell wall has been highly controversial. FtsZ and PG together function in orchestrating cell division and maintaining cell shape in almost all other bacteria. Using electron cryotomography, mass spectrometry and fluorescent labelling dyes, here we show that some environmental chlamydiae have cell wall sacculi consisting of a novel PG type. Treatment with fosfomycin (a PG synthesis inhibitor) leads to lower infection rates and aberrant cell shapes, suggesting that PG synthesis is crucial for the chlamydial life cycle. Our findings demonstrate for the first time the presence of PG in a member of the *Chlamydiae*. They also present a unique example of a bacterium with a PG sacculus but without FtsZ, challenging the current hypothesis that it is the absence of a cell wall that renders FtsZ non-essential.

¹Division of Biology, California Institute of Technology, Pasadena, California 91125, USA. ²Howard Hughes Medical Institute, Pasadena, California 91125, USA. ³Division of Microbial Ecology, University of Vienna, Vienna, A-1090, Austria. ⁴Institute for Cell and Molecular Biosciences, The Centre for Bacterial Cell Biology, Newcastle University, Newcastle upon Tyne, NE2 4AX, UK. ⁵Institute for Cell and Molecular Biosciences, Pinnacle Laboratory, Newcastle University, Newcastle upon Tyne, NE2 4AX, UK. ⁶Indiana University, Bloomington, Indiana 47405, USA. *These authors contributed equally to this work. Correspondence and requests for materials should be addressed to M.H. (email: horn@microbial-ecology.net) or to G.J.J. (email: jensen@caltech.edu).

Chlamydiae are members of the *Planctomycetes–Verrucomicrobia–Chlamydiae* (PVC) bacterial superphylum¹. Like most other bacteria, some PVC bacteria are already known to possess peptidoglycan (PG), that is, chains of alternating N-acetylglucosamine and N-acetylmuramic acid sugars crosslinked by short peptides. PVC bacteria also display striking eukaryote-like and archaea-like cell biological features, which have suggested intriguing hypotheses about their role in cellular evolution^{2,3}.

In *Verrucomicrobia* and almost all other bacteria, septal PG synthesis is orchestrated by the FtsZ cytoskeleton^{4,5}. In contrast, *Planctomycetes* lack both PG and FtsZ². This fits to the notion that PG-loss renders FtsZ dispensable, like in mycoplasmas⁶ or L-form bacilli⁷. Although from genome sequences it is clear that chlamydiae do not possess FtsZ, the presence or absence of chlamydial PG has been highly controversial^{8–16}. One early study reported the colorimetric detection of muramic acid in chlamydiae¹⁴, but more reliable chromatographic methods subsequently failed to confirm this result^{11,15}. All attempts to purify chlamydial sacculi have failed^{10,12} and no periplasmic density layers have been detected between the inner and outer membranes of chlamydiae by electron microscopy (EM) (including for instance^{17–19}). The apparent absence of PG in chlamydiae is surprising, however, as despite their highly reduced genomes, a nearly complete pathway for the synthesis of PG is present in the genomes of all chlamydiae¹³. In addition, several of the chlamydial PG biosynthetic enzymes have been characterized and shown to be functional *in vitro* and in complementation assays^{20–25}.

Here we look for the evidence of PG cell walls in two diverse and deeply rooting chlamydiae²⁶, *Protochlamydia amoebophila* and *Simkania negevensis*. Through electron cryotomography (ECT), biochemical purification, enzymatic digestion, mass spectrometry (MS), fluorescence microscopy and antibiotic treatment, we show that *P. amoebophila* are indeed surrounded by sacculi containing a new type of PG. In contrast, no evidence of PG is found in *S. negevensis*. These results prove that some chlamydiae do in fact synthesize PG sacculi, explaining the presence of PG-synthetic genes, but raising new questions about the identity and purpose of the modifications and the mechanisms of cell division in the absence of FtsZ.

Results

ECT of the chlamydial cell envelope. Two diverse and deeply rooting members of the chlamydial phylum, *S. negevensis* and *P. amoebophila*, were imaged by ECT in a near-native state. Bacteria were purified from amoeba cultures, plunge-frozen and 25 and 20 tomograms were collected of intact cells (Fig. 1). Density profiles through the cell envelopes of the two species were quite different. Although four layers were resolved in *Simkania* envelopes (Fig. 1b,c), five layers were resolved in *Protochlamydia* (Fig. 1e,f). Because the individual leaflets of lipid bilayers can be resolved in some cryotomograms, especially when the images are taken close to focus, in the case of *Simkania*, it is unclear whether the four layers represent the two leaflets of the outer and inner membranes ('O' and 'A' being the two leaflets of the outer membrane, and layers 'B' and 'I' being the two leaflets of the inner membrane), or whether one or more of these layers are non-membranous. The facts that layers O and A have fairly similar contrast and are consistently spaced even through the undulations are consistent with them being two leaflets of a single (outer) bilayer membrane. Their separation (~5 nm), however, is much larger than typical phospholipid bilayer membranes, whose two density peaks (from the phospholipid head groups) are only 3.7–4 nm apart²⁷. Similarly, *Simkania* layers B and I may be the

two leaflets of a single (inner) membrane, as they have similar contrast and a consistent spacing, but again they appear too far apart. In contrast to the *Simkania* envelope, the profile of *Protochlamydia* surprisingly resembled those of other Gram-negative bacteria with two membranes and a PG cell wall^{28–30}. Between the *Protochlamydia* outer and inner membranes (labelled 'O' and 'I', respectively) there appeared to be three additional layers (labelled 'C'–'E'). The similar-looking three layers in *Treponema pallidum* (from the outside in) were identified as proteinaceous (lipoproteins), PG and again proteinaceous²⁹. By analogy this suggests that layer C is composed of lipoproteins (perhaps connecting the outer membrane to the cell wall) and other outer-membrane-associated proteins, layer D is a cell wall and layer E is composed of lipoproteins and inner-membrane-associated proteins (perhaps including the penicillin-binding proteins responsible for cell wall synthesis, the lipoprotein OmcA and the cysteine-rich protein OmcB). Although other interpretations remain possible (cysteine-rich disulphide-crosslinked envelope proteins have been suggested to be the functional equivalent of PG in chlamydiae³¹), the most important and clear observation was that *Protochlamydia* exhibit a distinct periplasmic layer (D).

Purification and imaging of sacculi. In order to explore whether any of the observed periplasmic layers consisted of PG, we attempted to purify sacculi by boiling chlamydial cells (obtained from asynchronously infected amoeba cultures) in 4% sodium dodecyl sulphate. Strikingly, in three independent experiments, we observed sacculus-like structures in preparations from *Protochlamydia* (Fig. 2a–d), but not from *Simkania* (two experiments). *Protochlamydia* sacculi diameters (679 nm ± 34 s.d. *n* = 10) and morphologies matched the size and shape of intact cells. *Protochlamydia* sacculi had one or two 5–7 nm thick layers (arrowheads in Fig. 2e), plus mesh-like (up to 30 nm long) high-density aggregates attached to the outside (arrows in Fig. 2d).

Digestion and biochemical analyses of sacculi. To check for the presence of PG in the purified sacculi, we digested the samples with cellosyl, a glycan strand-cleaving PG muramidase. Cellosyl released soluble material from insoluble sacculi, which was reduced with sodium borohydride and analysed by high-pressure liquid chromatography (HPLC) using conditions for separating muropeptides³². The chromatogram (Fig. 2f) showed three main peaks in the monomeric region (20–50 min) and many peaks after 60 min that are poorly separated at higher retention time (>75 min) forming a 'hump', which is typical for highly crosslinked and/or incompletely digested PG material³³. The retention times and overall pattern of cellosyl digestion products were different, however, from those of muropeptide mixtures obtained from other Gram-positive and Gram-negative bacteria.

To characterize this material, the three main cellosyl products in the monomeric region and one well-separated main product at the beginning of the 'hump' region were analysed by MS. The determined neutral masses of the earlier three products were higher than what would be expected for monomeric muropeptides, but the masses of products 1 and 2 and of products 2 and 3 both differed by 71 Da, a typical feature of monomeric muropeptides with a tri-, tetra- and pentapeptide, respectively, due to the presence of none, one or two D-alanine residues (Fig. 2f). In MS/MS analysis, the three products were fragmented in a similar way, showing that they are related (Supplementary Fig. S1, Table 1). For all three peaks, we observed mass differences to the parent ion corresponding to the loss of GlcNAc, GlcNAcMurNAc(r) (r, indicates reduction to N-acetylmuramitol),




Figure 1 | Chlamydial cell envelopes are multi-layered. *Simkania* (a–c) and *Protochlamydia* (d–f) cells were purified from asynchronously infected amoeba cultures, plunge-frozen and imaged by ECT. Shown are tomographic slices through reticulate bodies (a,d and b,e enlarged) and corresponding density profiles (c,f) of the cell envelopes. Profiles are enlarged, aligned and cropped relative to the outer membrane. Distances between peaks (in nm) are indicated. In contrast to the *Simkania* profile, the *Protochlamydia* profile resembles those of other bacteria with PG cell walls (see text for a full discussion of each profile and layer). Scale bar, 100 nm in a,d and 20 nm in b,c,e,f.




Figure 2 | *Protochlamydia* synthesize purifiable sacculi that contain PG. Cryoprojections (a) and tomographic slices (b–e) through sacculi-like structures purified from *Protochlamydia* cells. We were unable to obtain similar structures from *Simkania*. Sacculi had one or two layers (arrowheads) plus short high-density filaments (arrows) on the outside (c enlarged in d,e). Sacculi were digested, reduced and separated by HPLC (f). MS analysis of peaks 1–3 (neutral masses indicated in Da) indicated the presence of modified PG (see also Table 1 and Supplementary Fig. S1). Scale bars, 100 nm.

GlcNAcMurNAc(r)-L-alanine and GlcNAcMurNAc(r)-L-alanine-D-glutamate, confirming that products 1–3 are all muropeptides. The neutral masses of the *Protochlamydia* products 1, 2 and 3 were all 314.12 Da larger than the masses of the reduced monomeric muropeptides (with tri-, tetra- or pentapeptide) from Gram-negative bacteria³⁴, however, suggesting the presence of a common modification in the *Protochlamydia* muropeptides. The

neutral mass of product 4 was consistent with a peptide crosslinked dimer of product 2. Additional mass differences that occurred in all fragmentation spectra indicated the presence of the same and as yet unknown modifications with 129 and 203 Da, respectively, explaining the higher mass of *Protochlamydia* muropeptides 1–3 compared to the monomeric muropeptides of *Escherichia coli*.

Table 1 | Fragmentation masses from fractions of reduced *P. amoebophila* muropeptides.

Fraction number or muro peptide	Fragment	Mass of fragments, H ⁺ form (m/z)		Proposed structure
		Determined	Calculated	
1	A	1,167.3447	1,167.4908	[M + H] ⁺ - H ₂ O
	B	1,056.3231	1,056.5014	[M + H] ⁺ - 129.2
	C	982.3848	982.4220	[M + H] ⁺ - GlcNAc
	D	853.3392	853.4220	[M + H] ⁺ - 129.0 - GlcNAc
	E	705.3032	705.3059	[M + H] ⁺ - GlcNAcMurNAc(r)
	F	634.3148	634.2688	[M + H] ⁺ - GlcNAcMurNAc(r)Ala
	G	505.2668	505.2262	[M + H] ⁺ - GlcNAcMurNAc(r)AlaGlu
2	A	1,238.3288	1,238.5255	[M + H] ⁺ - H ₂ O
	B	1,127.4075	1,127.5361	[M + H] ⁺ - 129.1
	C	1,053.3859	1,053.4567	[M + H] ⁺ - GlcNAc
	D	924.3934	924.4567	[M + H] ⁺ - 129.0 - GlcNAc
	E	776.3376	776.3406	[M + H] ⁺ - GlcNAcMurNAc(r)
	F	705.3202	705.3035	[M + H] ⁺ - GlcNAcMurNAc(r)Ala
	G	576.3141	576.2609	[M + H] ⁺ - GlcNAcMurNAc(r)AlaGlu
3	A	1,309.4104	1,309.5684	[M + H] ⁺ - H ₂ O
	B	1,198.3875	1,198.5790	[M + H] ⁺ - 129.2
	C	1,124.3796	1,124.4996	[M + H] ⁺ - GlcNAc
	D	995.3871	995.4996	[M + H] ⁺ - 129.0 - GlcNAc
	E	847.3485	847.3835	[M + H] ⁺ - GlcNAcMurNAc(r)
	F	776.3356	776.3464	[M + H] ⁺ - GlcNAcMurNAc(r)Ala
	G	647.3268	647.3038	[M + H] ⁺ - GlcNAcMurNAc(r)AlaGlu
Tetra from <i>C. jejuni</i>	A	946.4128	946.3869	[M + Na] ⁺ - H ₂ O
	B	n.d.	835.4202	[M + Na] ⁺ - 129
	C	761.3908	761.3181	[M + Na] ⁺ - GlcNAc
	D	n.d.	632.3181	[M + Na] ⁺ - 129 - GlcNAc
	E	484.3317	484.2019	[M + Na] ⁺ - GlcNAcMurNAc(r)
	F	413.2619	413.1648	[M + Na] ⁺ - GlcNAcMurNAc(r)Ala
	G	284.3200	284.1222	[M + Na] ⁺ - GlcNAcMurNAc(r)AlaGlu

n.d., not detected. Fraction numbers correspond to peak numbers in Fig. 2f. Fragments correspond to fragments in Supplementary Fig. S1. The parental ion ([M + H]⁺ or [M + Na]⁺) of each muropeptide fragments to ions lacking either water, an unknown modification with ~129 Da, a GlcNAc residue, ~129 Da and GlcNAc, a GlcNAcMurNAc(r) disaccharide (r, indicates reduction to muramitol), GlcNAcMurNAc(r)Ala or GlcNAcMurNAc(r)AlaGlu (see column 'Proposed structure'). Tetra, reduced disaccharide tetrapeptide muropeptide fraction from *Campylobacter jejuni*⁵⁶. The Na⁺ containing ion with m/z 964.4202 was fragmented.

To confirm that PG was a major component of *Protochlamydia* sacculi and check for the presence of disulphide-crosslinked protein components, purified sacculi were subjected to lysozyme and dithiothreitol treatment, respectively, and imaged with negative stain EM. Only the incubation with lysozyme degraded the sacculi (Supplementary Fig. S2).

Fluorescence imaging of D-alanine incorporation in vivo. To further confirm the presence of PG in *Protochlamydia* sacculi, we tested whether fluorescently labelled amino acids (fluorescent D-amino acids (FDAA); labelled D-alanine in our experiments)³⁵ would be incorporated into chlamydial cells *in vivo*. Incubation of amoeba cultures continuously infected with *Protochlamydia* (including reticulate bodies, elementary bodies and transitional stages) with FDAA (HADA (hydroxy coumarin-carbonyl-amino-D-alanine) and BADA (4,4-difluoro-5,7-dimethyl-4-bora-3a,4a-diaza-s-indacene-3-propionic acid-3-amino-D-alanine)) resulted in multiple strong and chlamydial cell-sized signals inside amoeba cells (Fig. 3a–d, Supplementary Fig. S3). In many cases, the FDAA labelling in infected amoebae overlapped with staining of the chlamydial cells by 4',6-diamidino-2-phenylindole (DAPI) or chlamydiae-specific fluorescence *in situ* hybridization (FISH). Not all cells stained by FISH/DAPI showed a corresponding FDAA signal, at least in part because chlamydial cells were in different developmental stages, including non-replicating elementary bodies. No signals were detected when uninfected amoebae were incubated with FDAA (Supplementary Fig. S3D, E), or when infected amoebae were incubated with dimethylsulphoxide only. Interestingly, purified *Protochlamydia* elementary bodies (which cannot undergo cell division and are therefore probably

not actively synthesizing new PG) also did not show labelling upon incubation with FDAA (Supplementary Fig. S3B,C), indicating that PG synthesis takes place during *Protochlamydia* replication inside the host. Amoeba cultures infected with *Simkania*, on the other hand, showed either no signals (using BADA) or signals similar to the background level (using HADA) upon labelling with FDAA (Supplementary Fig. S4A,B), consistent with the absence of purifiable sacculi. Purified *Simkania* cells were not labelled by either dye (Supplementary Fig. S4C,D).

Protochlamydia sensitivity to cell-wall-targeting antibiotics.

Owing to the high conservation of PG throughout the bacterial domain of life, many antibacterial drugs target PG synthesis. The so-called chlamydial anomaly¹⁶ is that despite the fact that PG has not been detected in pathogenic chlamydiae, these organisms are sensitive to cell-wall-targeting β-lactam antibiotics. Penicillin, for instance, leads to the formation of enlarged aberrant cells^{36,37} and blocks the conversion between developmental stages³⁸. Environmental chlamydiae, in contrast, are resistant to β-lactams^{39,40} –possibly due to putative β-lactamases encoded in their genomes. To explore the role of the PG sacculus in the *Protochlamydia* life cycle, we used an alternative PG synthesis-targeting antibiotic (fosfomycin) to treat infected amoeba cells. The addition of 500 μg ml⁻¹ fosfomycin to *Protochlamydia*-infected amoeba cultures led to a significant decrease in infection rate (20.2% ± 8 infected amoebae for fosfomycin-treated cultures versus 95.8% ± 2.2 infected amoebae for untreated cultures; *P* < 0.0001, unpaired *t*-test). *Protochlamydia* cells within treated cultures were also up to eight-times larger than normal




Figure 3 | *Protochlamydia* incorporate D-alanine *in vivo* and are sensitive to fosfomycin. *Protochlamydia*-infected amoebae (a–d), but not purified *Protochlamydia* cells (Supplementary Fig. S2) or uninfected amoeba cells (Supplementary Fig. S2), stained positively for new PG synthesis with fluorescently labelled D-alanine dyes HADA (a) and BADA (Supplementary Fig. S2), confirming the synthesis of PG in actively growing *Protochlamydia*. Specific FISH staining of chlamydial cells (b, overlay with a shown in c) and the eukaryotic host (d, shows overlay with c) are shown. The treatment of *Protochlamydia*-infected amoeba cultures (amoeba cell outlines in white) with cell-wall-synthesis-targeting fosfomycin (h, control shown in g) resulted in a dramatic decrease in infection rate and aberrant or larger chlamydial cell shapes (shown are immunofluorescent stainings of chlamydial outer membrane proteins), suggesting a crucial role of PG in the *Protochlamydia* life cycle. Fosfomycin did not affect *Simkania* (f, control shown in e). Scale bars, 5 μm in d, 10 μm in e–h.

(diameters of up to 6 μm) (Fig. 3g,h). Lower fosfomycin concentrations (25 $\mu\text{g ml}^{-1}$ and 100 $\mu\text{g ml}^{-1}$) induced the formation of fewer aberrant forms and did not affect the infection rate (not shown). Fosfomycin treatment of *Simkania*-infected amoeba cultures led to only a slight decrease in infection rate (59.9% \pm 5.6 infected amoebae for fosfomycin-treated cultures versus 67% \pm 1.7 infected amoebae for untreated cultures) and no differences in cell size were detected (Fig. 3e,f).

Discussion

We conclude that *Protochlamydia* synthesize sacculi containing PG that can be hydrolysed by cellosyl, contains monomeric and crosslinked muropeptides, and carries yet unknown modifications at virtually every subunit. No evidence of PG in *Simkania* was found. Fluorescence imaging of D-Ala incorporation *in vivo* and monitoring of cell wall antibiotic sensitivity further suggested that the *Protochlamydia* PG sacculus has an important role in cell cycle and shape. This challenges previous speculations that chlamydiae synthesize a small ring of PG only during cell division⁴¹. Because this might still be true for *Simkania* and pathogenic chlamydiae, however, our data prompts a reconsideration of whether these organisms lack PG entirely (and the effects of β -lactams are pleiotropic) or if they synthesize novel PG structures that are not purified by standard sacculus preparation protocols.

The presence of sacculi in *Protochlamydia* but not in *Simkania* matches the less complete set of synthetic genes in the latter: *Simkania*, as well as pathogenic chlamydiae, lack an undecaprenyl-diphosphate phosphatase (UppP) and alanine/glutamine racemases (Alr, MurI) (Supplementary Table S1)^{13,42,43}. Interestingly, transglycosylases have not been found in any chlamydial genomes (Supplementary Table S1)¹³ or in the genomes of a few other PG-possessing bacteria^{44,45}, so some other enzyme(s) must be capable of synthesizing glycan strands.

The presence of PG sacculi in *Protochlamydia* and in the *Chlamydiae*'s sister phylum *Verrucomicrobia*⁴⁶, together with the fact that the more basal chlamydial lineages have more complete PG synthesis pathways make it likely that the last common chlamydial ancestor synthesized a PG sacculus. The detection of a PG-containing sacculus in *Protochlamydia* challenges the view that FtsZ is essential in PG-possessing bacteria^{4,6,7}, however, because to our knowledge, *Protochlamydia* is the first example of a bacterium with a PG cell wall, but without FtsZ. Studying cell division and septal development in this organism could help clarify the role of FtsZ and the evolutionary transition to PG- and FtsZ-independency.

Methods

Cultivation of organisms. *Acanthamoeba castellanii* Neff infected with *P. amoebophila* UWE25, or *A. castellanii* UWC1 infected with *S. negevensis*, were cultivated in TSY medium (30 g l^{-1} trypticase soy broth, 10 g l^{-1} yeast extract, pH 7.3) at 20 °C. Amoebal growth was monitored by light microscopy and medium was exchanged every 3–6 days. The presence and identity of the chlamydial symbionts was verified by isolation of DNA from cultures followed by amplification and sequencing of the 16S rRNA genes. In addition, FISH using specific probes combined with DAPI staining of infected cultures was performed using specific probes for the respective symbiont⁴⁷. Amoebae infected with chlamydiae were allowed to attach on slides and were fixed with 4% formaldehyde at 20 °C. Cells were hybridized for 1.5 h at 46 °C at a formamide concentration of 25% with the *Protochlamydia*-specific probe E25-454 (5'-GGA TGT TAG CCA GCT CAT-3') or the *Simkania*-specific probe Simneg-183 (5'-CAG GCT ACC CCA GCT CTC-3') and the probe EUB338 (ref. 48). Subsequently, cells were stained with DAPI (0.5 $\mu\text{g ml}^{-1}$ in PBS) for 5 min, and slides were analysed using an epifluorescence microscope.

Purification of chlamydiae. Infected *A. castellanii* cultures were harvested by centrifugation (7,197 \times g, 10 min), washed in Page's Amoebic Saline (PAS)⁴⁹, centrifuged and resuspended in PAS. Amoeba cells were disrupted by vortexing with an equal volume of glass beads for 3 min. Glass beads and cell debris were removed by centrifugation (5 min, 300 \times g). The supernatant was filtered through a 1.2 μm filter and centrifuged at maximum speed for 10 min, 18,000 \times g. The obtained pellet was resuspended in PAS.

Plunge-freezing. For plunge-freezing, copper/rhodium EM grids (R2/2 or R2/1, Quantifoil) were glow-discharged for 1 min. A 20 × -concentrated bovine serum albumin-treated solution of 10 nm colloidal gold (Sigma) was added to purified chlamydiae or sacculi (1:4 v/v) immediately before plunge freezing. A 4-μl droplet of the mixture was applied to the EM grid, then automatically blotted and plunge-frozen into a liquid ethane-propane mixture⁵⁰ using a Vitrobot (FEI Company)⁵¹.

Electron cryotomography. Images were collected using a Polara 300 kV FEG transmission electron microscope (FEI Company) equipped with an energy filter (slit width 20 eV; Gatan) on a lens-coupled 4 k × 4 k UltraCam CCD (Gatan). Pixels on the CCD represented 0.95 nm (22,500 ×) or 0.63 nm (34,000 ×) at the specimen level. Typically, tilt series were recorded from -60° to +60° with an increment of 1° at 10 μm under-focus. The cumulative dose of a tilt-series was 180–220 e⁻/Å². UCSF Tomo⁵² was used for automatic acquisition of tilt-series and 2D projection images. Three-dimensional reconstructions were calculated using the IMOD software package⁵³ or Raptor⁵⁴. Tomograms were visualized using 3dMOD⁵³. Density profiles were generated using ImageJ.

Fluorescent labelling of PG. Newly synthesized PG was labelled using FDAA³⁵. *A. castellanii* cells continuously infected with *Protochlamydia* or *Simkania* were harvested and resuspended in a mixture of TSY and PAS (1:1). Cells were incubated with 1.5 mM HADA³⁵ or BADA for 6 h with gentle shaking. Cells were pelleted, washed three times and fixed with 4% formaldehyde followed by FISH using the chlamydia-specific probe Chls-0523 or DAPI-staining. As a control, uninfected amoebae and purified chlamydiae were treated in the same way.

Antibiotic treatment of infected amoeba cultures. *A. castellanii* were seeded into the wells of a multi-well dish and infected with purified *Simkania* and *Protochlamydia* elementary bodies⁵⁵. After centrifugation at 600 × g for 10 min, the medium was exchanged for TSY supplemented with fosfomycin (0, 25, 100 or 500 μg ml⁻¹, respectively). Medium was exchanged once at 48 h post infection. Cells were fixed with methanol at 96 h post infection followed by immunofluorescence analysis using either anti-PomS antibodies⁵⁵ or anti-*Simkania* antibodies raised against purified chlamydiae. The number of infected amoebae was counted for each treatment.

Preparation and composition analysis of sacculi. Chlamydial cells were purified from 1.81 (*Protochlamydia*) and 31 (*Simkania*) of infected amoeba culture and, depending on the amount of harvested cells, resuspended in 2.6–5.2 ml 4% SDS (w/v). After shipping (overnight, room temperature), the suspensions were dripped into 4% SDS (preheated to 90 °C, 6 ml final volume) and stirred for 2.5 h at 90 °C. Sacculi were pelleted (30 min, 135,000 × g) in a TLA-100.3 rotor (Beckman Coulter), washed 4 × in 3 ml water, resuspended in 150 μl water and supplemented with 0.02% (w/v) sodium azide. Muropeptides were released from the PG by an overnight incubation with the muramidase Cellosyl (Hoechst, Frankfurt am Main, Germany) on a thermal shaker at 37 °C and 800 rpm. The sample was boiled for 10 min and centrifuged for 10 min at 13,500 × g. The muropeptides present in the supernatant were reduced with sodium borohydride and separated on a 250 × 4.6 mm 3-μm ProntoSil 120-3-C18 AQ reversed phase column (Bischoff, Germany) at 55 °C using a 135 min gradient from 50 mM sodium phosphate pH 4.31 to 75 mM sodium phosphate pH 4.95, 15% methanol and a flow rate of 0.5 ml min⁻¹ (ref. 32). Muropeptides were detected at 205 nm. Fractions were collected, concentrated in a SpeedVac, acidified by 1% trifluoroacetic acid and analysed by offline electrospray MS on a Finnigan LTQ-FT mass spectrometer (ThermoElectron, Bremen, Germany) in a positive ion mode using mass scans over the mass range from $m/z = 300$ to $m/z = 1,900$ at a typical spray voltage of 1.1–1.5 kV³⁴. Parent ion scans were acquired with an FT MS resolution setting of 100,000 (at $m/z = 400$) with a typical mass accuracy of 3 ppm. MS/MS scans were performed in the ion trap, which has a typical mass accuracy for the fragment ion of ± 0.3 Da. MS spectra were deconvoluted to generate uncharged masses using the QualBrowser program (ThermoElectron, Bremen, Germany)³⁴.

Negative-stain EM of treated sacculi preparations. Purified sacculi were incubated at 37 °C for 12 h in 20 μl total volume with either lysozyme (10 mg ml⁻¹), dithiothreitol (5 mM) or phosphate buffered saline. Samples were applied to a Formvar-coated, carbon-coated, glow-discharged copper EM grid (Electron Microscopy Sciences). Samples were aspirated, stained with 1.5% uranylacetate and imaged on an FEI Tecnai T12 electron microscope.

References

- Wagner, M. & Horn, M. The *Planctomycetes*, *Verrucomicrobia*, *Chlamydiae* and sister phyla comprise a superphylum with biotechnological and medical relevance. *Curr. Opin. Biotechnol.* **17**, 241–249 (2006).
- Devos, D. P. & Reynaud, E. G. Evolution. Intermediate steps. *Science* **330**, 1187–1188 (2010).
- Reynaud, E. G. & Devos, D. P. Transitional forms between the three domains of life and evolutionary implications. *Proc. Biol. Sci.* **278**, 3321–3328 (2011).
- Erickson, H. P., Anderson, D. E. & Osawa, M. FtsZ in bacterial cytokinesis: cytoskeleton and force generator all in one. *Microbiol. Mol. Biol. Rev.* **74**, 504–528 (2010).
- Pilhofer, M. *et al.* Characterization and evolution of cell division and cell wall synthesis genes in the bacterial phyla *Verrucomicrobia*, *Lentisphaerae*, *Chlamydiae*, and *Planctomycetes* and phylogenetic comparison with rRNA genes. *J. Bacteriol.* **190**, 3192–3202 (2008).
- Lluch-Senar, M., Querol, E. & Pinol, J. Cell division in a minimal bacterium in the absence of ftsZ. *Mol. Microbiol.* **78**, 278–289 (2010).
- Leaver, M., Dominguez-Cuevas, P., Coxhead, J. M., Daniel, R. A. & Errington, J. Life without a wall or division machine in *Bacillus subtilis*. *Nature* **457**, 849–853 (2009).
- Manire, G. P. & Tamura, A. Preparation and chemical composition of the cell walls of mature infectious dense forms of meningopneumonitis organisms. *J. Bacteriol.* **94**, 1178–1183 (1967).
- Tamura, A. & Manire, G. P. Preparation and chemical composition of the cell membranes of developmental reticulate forms of meningopneumonitis organisms. *J. Bacteriol.* **94**, 1184–1188 (1967).
- Caldwell, H. D., Kromhout, J. & Schachter, J. Purification and partial characterization of the major outer membrane protein of *Chlamydia trachomatis*. *Infect. Immun.* **31**, 1161–1176 (1981).
- Fox, A. *et al.* Muramic acid is not detectable in *Chlamydia psittaci* or *Chlamydia trachomatis* by gas chromatography-mass spectrometry. *Infect. Immun.* **58**, 835–837 (1990).
- Barbour, A. G., Amano, K., Hackstadt, T., Perry, L. & Caldwell, H. D. *Chlamydia trachomatis* has penicillin-binding proteins but not detectable muramic acid. *J. Bacteriol.* **151**, 420–428 (1982).
- McCoy, A. J. & Maurelli, A. T. Building the invisible wall: updating the chlamydial peptidoglycan anomaly. *Trends Microbiol.* **14**, 70–77 (2006).
- Perkins, H. R. & Allison, A. C. Cell-wall constituents of Rickettsiae and Psittacosis-Lymphogranuloma organisms. *J. Gen. Microbiol.* **30**, 469–480 (1963).
- Garrett, A. J., Harrison, M. J. & Manire, G. P. A search for the bacterial mucopeptide component, muramic acid, in *Chlamydia*. *J. Gen. Microbiol.* **80**, 315–318 (1974).
- Moulder, J. W. Why is *Chlamydia* sensitive to penicillin in the absence of peptidoglycan? *Infect. Agents. Dis.* **2**, 87–99 (1993).
- Matsumoto, A. Recent progress of electron microscopy in microbiology and its development in future: from a study of the obligate intracellular parasites, chlamydia organisms. *J. Electron. Microsc.* **28**, 57–64 (1979).
- Tamura, A., Matsumoto, A., Manire, G. P. & Higashi, N. Electron microscopic observations on the structure of the envelopes of mature elementary bodies and developmental reticulate forms of *Chlamydia psittaci*. *J. Bacteriol.* **105**, 355–360 (1971).
- Huang, Z. *et al.* Cryo-electron tomography of *Chlamydia trachomatis* gives a clue to the mechanism of outer membrane changes. *J. Electron. Microsc.* **59**, 237–241 (2010).
- McCoy, A. J. & Maurelli, A. T. Characterization of *Chlamydia* MurC-Ddl, a fusion protein exhibiting D-alanyl-D-alanine ligase activity involved in peptidoglycan synthesis and D-cycloserine sensitivity. *Mol. Microbiol.* **57**, 41–52 (2005).
- McCoy, A. J., Sandlin, R. C. & Maurelli, A. T. In vitro and in vivo functional activity of *Chlamydia* MurA, a UDP-N-acetylglucosamine enolpyruvyl transferase involved in peptidoglycan synthesis and fosfomycin resistance. *J. Bacteriol.* **185**, 1218–1228 (2003).
- Hesse, L. *et al.* Functional and biochemical analysis of *Chlamydia trachomatis* MurC, an enzyme displaying UDP-N-acetylmuramate:amino acid ligase activity. *J. Bacteriol.* **185**, 6507–6512 (2003).
- Patin, D., Bostock, J., Blanot, D., Mengin-Lecreulx, D. & Chopra, I. Functional and biochemical analysis of the *Chlamydia trachomatis* ligase MurE. *J. Bacteriol.* **191**, 7430–7435 (2009).
- Patin, D., Bostock, J., Chopra, I., Mengin-Lecreulx, D. & Blanot, D. Biochemical characterisation of the chlamydial MurF ligase, and possible sequence of the chlamydial peptidoglycan pentapeptide stem. *Arch. Microbiol.* **194**, 505–512 (2012).
- Henrichfreise, B. *et al.* Functional conservation of the lipid II biosynthesis pathway in the cell wall-less bacteria *Chlamydia* and *Wolbachia*: why is lipid II needed? *Mol. Microbiol.* **73**, 913–923 (2009).
- Horn, M. Chlamydiae as symbionts in eukaryotes. *Annu. Rev. Microbiol.* **62**, 113–131 (2008).
- Leforestier, A., Lemerrier, N. & Livolant, F. Contribution of cryoelectron microscopy of vitreous sections to the understanding of biological membrane structure. *Proc. Natl Acad. Sci. USA* **109**, 8959–8964 (2012).
- Tocheva, E. I., Li, Z. & Jensen, G. J. Electron cryotomography. *Cold Spring Harb. Perspect. Biol.* **2**, a003442 (2010).
- Liu, J. *et al.* Cellular architecture of *Treponema pallidum*: novel flagellum, periplasmic cone, and cell envelope as revealed by cryo electron tomography. *J. Mol. Biol.* **403**, 546–561 (2010).

30. Tocheva, E. I. *et al.* Peptidoglycan remodeling and conversion of an inner membrane into an outer membrane during sporulation. *Cell* **146**, 799–812 (2011).
31. Hatch, T. Disulfide cross-linked envelope proteins: the functional equivalent of peptidoglycan in chlamydiae? *J. Bacteriol.* **178**, 1–5 (1996).
32. Glauner, B. Separation and quantification of mucopeptides with high-performance liquid chromatography. *Anal. Biochem.* **172**, 451–464 (1988).
33. de Jonge, B. L. & Tomasz, A. Abnormal peptidoglycan produced in a methicillin-resistant strain of *Staphylococcus aureus* grown in the presence of methicillin: functional role for penicillin-binding protein 2A in cell wall synthesis. *Antimicrob. Agents. Chemother.* **37**, 342–346 (1993).
34. Bui, N. K. *et al.* The peptidoglycan sacculus of *Myxococcus xanthus* has unusual structural features and is degraded during glycerol-induced myxospore development. *J. Bacteriol.* **191**, 494–505 (2009).
35. Kuru, E. *et al.* In Situ probing of newly synthesized peptidoglycan in live bacteria with fluorescent D-amino acids. *Angew. Chem.* **51**, 12519–12523 (2012).
36. Matsumoto, A. & Manire, G. P. Electron microscopic observations on the effects of penicillin on the morphology of *Chlamydia psittaci*. *J. Bacteriol.* **101**, 278–285 (1970).
37. Skilton, R. J. *et al.* Penicillin induced persistence in *Chlamydia trachomatis*: high quality time lapse video analysis of the developmental cycle. *PLoS One* **4**, e7723 (2009).
38. Beatty, W. L., Morrison, R. P. & Byrne, G. I. Persistent chlamydiae: from cell culture to a paradigm for chlamydial pathogenesis. *Microbiol. Rev.* **58**, 686–699 (1994).
39. Maurin, M., Bryskier, A. & Raoult, D. Antibiotic susceptibilities of *Parachlamydia acanthamoeba* in amoebae. *Antimicrob. Agents Chemother.* **46**, 3065–3067 (2002).
40. Kahane, S., Gonen, R., Sayada, C., Elion, J. & Friedman, M. G. Description and partial characterization of a new chlamydia-like microorganism. *FEMS. Microbiol. Lett.* **109**, 329–334 (1993).
41. Brown, W. J. & Rockey, D. D. Identification of an antigen localized to an apparent septum within dividing chlamydiae. *Infect. Immun.* **68**, 708–715 (2000).
42. Horn, M. *et al.* Illuminating the evolutionary history of chlamydiae. *Science* **304**, 728–730 (2004).
43. Collingro, A. *et al.* Unity in variety—the Pan-genome of the *Chlamydiae*. *Mol. Biol. Evol.* **28**, 3253–3270 (2011).
44. McPherson, D. C. & Popham, D. L. Peptidoglycan synthesis in the absence of Class A penicillin-binding proteins in *Bacillus subtilis*. *J. Bacteriol.* **185**, 1423–1431 (2003).
45. Arbeloa, A. *et al.* Role of class A penicillin-binding proteins in PBP5-mediated β -lactam resistance in *Enterococcus faecalis*. *J. Bacteriol.* **186**, 1221–1228 (2004).
46. Hedlund, B. P., Gosink, J. J. & Staley, J. T. Phylogeny of *Prostheco bacter*, the fusiform caulobacters: members of a recently discovered division of the Bacteria. *Int. J. Syst. Bacteriol.* **46**, 960–966 (1996).
47. Schmitz-Esser, S. *et al.* Diversity of bacterial endosymbionts of environmental *Acanthamoeba* isolates. *Appl. Environ. Microbiol.* **74**, 5822–5831 (2008).
48. Amann, R. I. *et al.* Combination of 16S rRNA-targeted oligonucleotide probes with flow cytometry for analyzing mixed microbial populations. *Appl. Environ. Microbiol.* **56**, 1919–1925 (1990).
49. Page, F. C. *An Illustrated Key to Freshwater and Soil Amoebae* (Freshwater Biological Association, 1976).
50. Tivol, W. F., Briegel, A. & Jensen, G. J. An improved cryogen for plunge freezing. *Microsc. Microanal.* **14**, 375–379 (2008).
51. Iancu, C. V. *et al.* Electron cryotomography sample preparation using the Vitrobot. *Nat. Protocols* **1**, 2813–2819 (2006).
52. Zheng, S. Q. *et al.* UCSF tomography: an integrated software suite for real-time electron microscopic tomographic data collection, alignment, and reconstruction. *J. Struct. Biol.* **157**, 138–147 (2007).
53. Kremer, J. R., Mastronarde, D. N. & McIntosh, J. R. Computer visualization of three-dimensional image data using IMOD. *J. Struct. Biol.* **1996/01/01**, 71–76 (1996).
54. Amat, F. *et al.* Markov random field based automatic image alignment for electron tomography. *J. Struct. Biol.* **161**, 260–275 (2008).
55. Aistleitner, K. *et al.* Identification and characterization of a novel porin family highlights a major difference in the outer membrane of chlamydial symbionts and pathogens. *PLoS One* **8**, e55010 (2013).
56. Frirdich, E. *et al.* Peptidoglycan-modifying enzyme Pgp1 is required for helical cell shape and pathogenicity traits in *Campylobacter jejuni*. *PLoS Pathog.* **8**, e1002602 (2012).

Acknowledgements

This work was funded by the Austrian Science Fund FWF (Y277-B03 to M.H.), the European Research Council (ERC StG 'EvoChlamy' to M.H.), NIH grant GM094800B (to G.J.J.), the Caltech Center for Environmental Microbiology Interactions (to G.J.J., M.P.), a gift from the Gordon and Betty Moore Foundation to Caltech, the BBSRC (BB/I020012/1 to W.V.), NIH grant AI059327 (to M.S.V.) and NIH grant GM051986 (to Y.V.B.). We thank Elitza Tocheva for discussions on the preparation of sacculi.

Author contributions

M.P. initiated the study. M.P. and K.A. performed all experiments except HPLC/MS analyses of sacculi, which were done by J.B., J.G. and W.V. E.K., E.H., Y.V.B. and M.S.V. provided the FDAA dyes and advice on the FDAA labelling experiments. M.P., K.A., W.V., M.H. and G.J.J. designed the experiments and wrote the manuscript.

Additional information

Supplementary Information accompanies this paper at <http://www.nature.com/naturecommunications>

Competing financial interests: The authors declare no competing financial interests.

Reprints and permission information is available online at <http://npg.nature.com/reprintsandpermissions/>

How to cite this article: Pilhofer, M. *et al.* Discovery of chlamydial peptidoglycan reveals bacteria with murein sacculi but without PtsZ. *Nat. Commun.* **4**:2856 doi: 10.1038/ncomms3856 (2013).