

ARTICLE

Received 24 Jan 2014 | Accepted 16 May 2014 | Published 18 Jun 2014

DOI: 10.1038/ncomms5140

TOPLESS mediates brassinosteroid-induced transcriptional repression through interaction with BZR1

Eunkyoo Oh^{1,*}, Jia-Ying Zhu^{1,*}, Hojin Ryu^{2,3}, Ildoo Hwang² & Zhi-Yong Wang¹

Brassinosteroid (BR) regulates plant development by activating the transcription factor brassinazole resistant 1 (BZR1), which activates and represses different target genes to switch cellular programmes. The mechanisms that determine BZR1's transcriptional activities remain largely unknown. Here we show that BZR1 represses target genes by recruiting the Groucho/TUP1-like transcriptional corepressor TOPLESS (TPL). Specific deletion or mutation of an evolutionarily conserved ERF-associated amphiphilic repression (EAR) motif at the carboxy terminus abolishes BZR1's abilities to regulate gene expression and cell elongation, but these defects are rescued by TPL fusion to the EAR motif-mutated BZR1. The EAR motif in BZR1 mediates recruitment of TPL to BZR1-repressed promoters. A triple *tpl* mutant (*tpl;tpr1;tpr4*) shows reduced BR sensitivity and suppresses the gain-of-function *bzr1-1D* mutant phenotype. BR repression of gene expression also requires histone deacetylases that interact with TPL. Our study demonstrates key roles of the EAR motif and TPL in BR regulation of gene expression and plant growth.

¹Department of Plant Biology, Carnegie Institution for Science, 260 Panama Street, Stanford, California 94305, USA. ²Department of Life Sciences, POSTECH Biotech Center, Pohang University of Science and Technology, Pohang 790-784, Korea. ³Microbial Plant Activation Lab, Agricultural Microbiology Division, National Academy of Agricultural Science, RDA, Suwon 441-707, South Korea. * These authors contributed equally to this work. Correspondence and requests for materials should be addressed to Z.-Y.W. (email: zywang24@stanford.edu).

Steroids are important hormones for regulation of gene expression and development in both animals and plants. Unlike animal steroid hormones, which directly bind to nuclear receptor transcription factors, the plant steroid hormone brassinosteroid (BR) binds to a transmembrane receptor kinase, brassinosteroid insensitive1 (BRI1)¹, which has a leucine-rich repeat extracellular domain similar to the Toll receptors in metazoans². BRI1 signalling activates the members of the brassinazole resistant1 (BZR1) family of transcription factors through a phosphorylation-mediated signal transduction pathway^{3,4}. BZR1 activates and represses different target genes to programme genome expression and cell growth. The mechanisms that specify BZR1's function in transcription activation and repression are not understood.

BZR1 is a plant-specific transcription factor with a DNA-binding domain that recognizes BR response element (CGTG(T/C)G)⁵. Both subcellular localization and DNA-binding activity of BZR1 are tightly regulated by phosphorylation. In the absence of BR, a glycogen synthase kinase 3 (GSK3)-like kinase named BIN2 interacts with BZR1 via a docking motif at the C terminus of BZR1 and phosphorylates BZR1 (refs 6–8). The phosphorylated BZR1 is unable to bind DNA and also is retained in the cytosol due to binding to the 14-3-3 proteins^{9,10}. BR binding to BRI1 triggers a signalling cascade that leads to inactivation of BIN2 (ref. 11), resulting in dephosphorylation of BZR1 catalysed by protein phosphatase 2A (PP2A)¹². Dephosphorylated BZR1 enters the nucleus and binds to target promoters to mediate BR-responsive gene expression³. BZR2 (also named BES1), the close homologue of BZR1, is regulated through similar mechanisms^{6,12–14}.

Similar to many plant transcription factors, BZR1 functions as a transcriptional repressor for some promoters but an activator for others. Previous genome-wide identification of direct binding sites of BZR1 coupled with transcriptome profiling showed that BZR1 binds to promoters of both BR-repressed and -induced genes^{5,15}. Apparently, BZR1 transcriptional activity is partly determined by the context of promoter *cis*-elements and *trans*-factors. Indeed the genome-wide data showed that BR response element is enriched in the promoters of BR-repressed gene, whereas the E-box motif (CANNTG), a binding site for bHLH factors, is more enriched in the promoters of BR-activated genes¹⁵. BZR1 and its homologous BZR2/BES1 interact with the bHLH factor phytochrome interacting factor4 (PIF4) and BZR1 shares with PIF4 over 50% of their target genes in the genome. BZR2/BES1 also interacts with BES1-interacting Myc-like1 (BIM1). The BZR1-PIF4 and BZR2-BIM1 heterodimers bind to the E-box motif in the promoters of target genes and activate gene expression^{13,16}. Several other transcription factors and histone-modifying enzymes have been reported to interact with BZR2/BES1 and participate in BZR2/BES1 activation of gene expression^{17–19}. In addition, recent studies have suggested that myeloblastosis transcription factor like2 (MYBL2) and homeodomain-leucine zipper protein1 (HAT1) interact with BZR2/BES1 and repress several BR target genes^{20,21}. However, it remains unclear how BZR1 represses gene expression, and specifically how its gene repression and activation activities are mediated by interactions with general transcriptional regulators.

In this study, a genetic screen for the intragenic suppressors of the gain-of-function mutant *bzr1-1D* reveals an important function of an evolutionarily conserved ERF-associated amphiphilic repression (EAR) motif in BZR1 regulation of target gene expression. We further show that BZR1 EAR motif mediates interaction with the Groucho/TUP1-like corepressor TOPLESS (TPL), selectively recruiting TPL to the promoters of BZR1-repressed target genes. BZR1 repression of gene expression requires both TPL and also the activity of histone deacetylase

(HDA) that is known to interact with TPL. Our study unveils the molecular mechanism of BR repression of gene expression through BZR1–TPL interaction.

Results

EAR motif is crucial for BZR1 functions. The gain-of-function mutant BZR1 protein, *bzr1-1D*, is constitutively dephosphorylated and active even in the BR deficiency condition due to a strong interaction with PP2A¹². Thus, the hypocotyl elongation of *bzr1-1D* seedling is resistant to BR biosynthesis inhibitors, brassinazole (BRZ) and propiconazole (PPZ)¹². We identified suppressors of the *bzr1-1D* mutant through an EMS mutagenesis screen. One of the mutants, *bzs285*, carried a mutation that creates a stop codon after amino acid 284, resulting in deletion of 52 amino acids at the C terminus of *bzr1-1D* protein (Fig. 1a,b). The truncated region includes a BIN2 docking motif and an LxLxL type of EAR motif. As the BIN2 docking motif mediates inactivation of BZR1 through BIN2-mediated phosphorylation⁸, the suppression of the *bzr1-1D* phenotype by the *bzs285* mutation was most likely to be due to the loss of the EAR motif. The EAR motif is highly conserved in the BZR1 family of transcription factors across land plants from bryophytes to angiosperms (Supplementary Fig. 1a). In addition, all other BZR1-like proteins in *Arabidopsis* have the EAR motif at their C terminus (Supplementary Fig. 1b), implicating an indispensable role of the EAR motif in BZR1 functions.

To functionally characterize the EAR motif of BZR1, we generated transgenic plants overexpressing the EAR motif-deleted BZR1 (*BZR1ΔEAR-OX*) or *bzr1-1D* (*bzr1-1DΔEAR-OX*). Transgenic plants overexpressing wild-type BZR1 (*BZR1-OX*) were almost indistinguishable from wild-type plants (Fig. 1c) due to posttranslational regulation of BZR1 activity by BIN2, whereas overexpression of *bzr1-1D* (*bzr1-1D-OX*) caused BR-activation phenotypes similar to the *bzr1-1D* mutant (Fig. 1c), including leaf bending, organ fusion, and insensitivity to BR inhibitors²². In contrast, both *BZR1ΔEAR-OX* and *bzr1-1DΔEAR-OX* plants displayed severe dwarfism (Fig. 1c), similar to the BR-insensitive or BR-deficient mutant plants, suggesting that the EAR-deleted BZR1 proteins have dominant negative effects on BR responses. Consistent with *bzr1-1D* suppression by *bzs285*, *bzr1-1DΔEAR-OX* plants were sensitive to BR biosynthetic inhibitor PPZ (Fig. 1d) and showed no organ-fusion phenotypes in contrast to the fusion between cauline leaves and axillary branches often observed in the *bzr1-1D* and *bzr1-1D-OX* plants²² (Fig. 1e). The *bzr1-1DΔEAR-OX* seedlings grown in the dark also showed shorter hypocotyls (Fig. 1d). These results confirm an essential role of the EAR motif in BZR1 function.

To determine whether the dwarfism is caused by a defect in BR signalling or BR biosynthesis, we examined BR response of the *bzr1-1DΔEAR-OX*. Although the hypocotyls of *bzr1-1D* were more elongated by BR treatment than those of wild type, hypocotyls of *bzr1-1DΔEAR-OX* were insensitive to BR (Fig. 1f), indicating that BR signalling is impaired in the *bzr1-1DΔEAR-OX* plants. BR inhibits primary root growth mainly through reducing cell division in the root meristem²³. Root growth of *bzr1-1D* was more sensitive to BR inhibition than wild type, whereas roots of the *bzr1-1DΔEAR-OX* were insensitive to BR (Fig. 1g). Similar to *bzr1-1DΔEAR-OXs*, *BZR1ΔEAR-OXs* also showed BR insensitivity in hypocotyl elongation (Fig. 1h). The severe dwarfism and BR insensitivity of *BZR1ΔEAR-OX* and *bzr1-1DΔEAR-OX* are most likely to be due to the dominant negative effects of the EAR motif-deleted BZR1 over endogenous BZR1 and BZR1-like transcription factors. Together, these results demonstrate that the EAR motif is essential for BZR1 functions in various BR responses.




Figure 1 | EAR motif is essential for BZR1 function in BR responses. (a) Glutamine (Q) at position 285 in BZR1 is converted to stop codon in the *bzs285* mutant. BIN2-DM: BIN2 docking motif; EAR: EAR motif. (b) Intragenic mutant (*bzs285*) suppresses the constitutive activation of BR response of *bzs1-1D*. Seedlings were grown in the dark on the medium containing 2 μ M PPZ. Scale bar, 5 mm. (c) Both *BZR1*Δ*EAR*-OXs and *bzs1-1D*Δ*EAR*-OXs show severe dwarfism. Plants were grown in the long-day condition (16 h of light and 8 h of dark) for 4 weeks. Two independent lines (1 and 2) for the *BZR1*Δ*EAR*-OXs and *bzs1-1D*Δ*EAR*-OXs are shown. (d) The hypocotyl elongation of *bzs1-1D*Δ*EAR*-OX is inhibited by PPZ treatment. Seedlings were grown in the dark for 5 days on the medium containing either the mock or 2 μ M PPZ. Error bars indicate the s.d. ($n = 10$ plants). Scale bar, 5 mm. (e) Axillary branches and cauline leaves are not fused in *bzs1-1D*Δ*EAR*-OX. (f,g) *bzs1-1D*Δ*EAR*-OXs are insensitive to BR in the promotion of hypocotyl elongation (f) and the inhibition of primary root growth (g). Seedlings were grown under the white light for 5 days on the medium containing either mock or 1,000 nM brassinolide (BL). Scale bar, 5 mm. (h) *BZR1*Δ*EAR*-OXs are insensitive to BR in the promotion of hypocotyl elongation. Seedlings were grown under the white light for 5 days on the medium containing either mock or 1,000 nM brassinolide (BL). Error bars in (f-h) indicate the s.d. ($n = 10$ plants).

EAR motif mediates BZR1 regulation of gene expression. We next analysed the effects of the EAR motif deletion on BZR1 regulation of target gene expression. BZR1 is known to directly repress BR biosynthesis genes, including *DWF4* and *CPD*, to provide the feedback regulation of BR level⁵. The expression levels of *DWF4* and *CPD* were dramatically reduced in *bzs1-1D* compared with wild type, but was only marginally reduced in the *bzs1-1D*Δ*EAR*-OX plants (Fig. 2a), indicating that the transcriptional repression activity is impaired in *bzs1-1D*Δ*EAR*. BZR1 also activates expression of many of its target genes¹⁵.

Although the expression levels of *SAURs*, *IAA19* and *PRE5* were highly increased by *bzs1-1D*, the expression of *SAUR19*, *IAA19* and *PRE5* was not activated, and *SAUR15* expression was only slightly activated by *bzs1-1D*Δ*EAR* (Fig. 2b), suggesting that the EAR motif is also required for the BZR1-mediated transcriptional activation of target genes.

To more directly examine the functions of the EAR motif in BZR1 regulation of gene expression, we generated transgenic plants expressing *bzs1-1D* or *bzs1-1D*Δ*EAR* fused to the glucocorticoid receptor (GR), *bzs1-1D*-GR or *bzs1-1D*Δ*EAR*-GR,




Figure 2 | EAR motif mediates BZR1 regulation of gene expression. (a,b) The deletion of EAR motif abolishes BZR1 regulation of gene expression. Seedlings were grown on the medium containing 2 μ M PPZ in the dark for 5 days before harvesting for RNA extraction. Gene expression levels are normalized to that of *PP2A* and are shown relative to the expression levels in wild type (Col-0). (c,d) The EAR motif-deleted *bzr1-1D* neither represses *CPD* expression (c) nor activates *SAUR15* expression (d). Four-week-old leaves of transgenic plants expressing *bzr1-1D-GR* or *bzr1-1DΔEAR-GR* were treated with mock or 40 μ M of dexamethasone (DEX) for 4 h. Ratios of expression levels in DEX-treated to mock-treated are shown. Each bar represents independent transgenic plant. (e,f) The EAR motif-deleted *bzr1-1D* cannot activate *PRE5* promoter activity (e) and cannot repress *CPD* promoter activity (f) in a transient gene expression assay. The 2 kb of *PRE5* promoter or 1 kb of *CPD* promoter fused to firefly luciferase reporter gene was co-transfected with 35S::*bzr1-1D* or 35S::*bzr1-1DΔEAR* into *Arabidopsis* mesophyll protoplast. The firefly luciferase activities were normalized by *Renilla* luciferase as an internal control and are shown relative to that of protoplasts transfected with 35S::GFP (GFP). ** $P < 0.01$ and * $P < 0.05$ by Student's *t*-test. (g) The deletion of EAR motif does not affect BZR1 DNA binding. Transgenic plants overexpressing *BZR1* and *BZR1ΔEAR* were treated with 100 nM BL and cross-linked for the ChIP assays. In the ChIP assays, the enrichment of DNA was calculated as the ratio between transgenic plants and wild-type control (Col-0), normalized to that of the *PP2A* coding region as an internal control. Error bars in (a,b,e,f,g) indicate the s.d. ($n = 3$).

which allows inducible nuclear localization of the fusion proteins. Transient DEX treatment (4 h) of the *bzr1-1D-GR* plants dramatically repressed *CPD* expression and activated *SAUR15* expression, confirming the inducible nuclear localization and function of *bzr1-1D-GR* *in planta* (Fig. 2c,d). In contrast, the DEX treatment of the *bzr1-1DΔEAR-GR* plants did not repress *CPD* or activate *SAUR15* (Fig. 2c,d), indicating that the EAR motif is necessary for both transcriptional repression and activation activities of BZR1. In fact, the *CPD* expression levels were slightly increased by DEX treatment in *bzr1-1DΔEAR-GR* plants (Fig. 2c), most likely to be due to the competition by the transcriptionally inactive *bzr1-1DΔEAR* protein with the endogenous BZR1 for target promoter occupancy. Consistent with the results of the DEX treatment assays, in transient gene expression assays using *Arabidopsis* mesophyll protoplasts, *bzr1-1D* increased *PRE5* promoter activity and decreased *DWF4* promoter activity, whereas *bzr1-1DΔEAR* could neither activate *PRE5* nor repress *DWF4* (Fig. 2e,f). The deletion of the EAR motif did not affect the BZR1 binding to the promoters of target genes (Fig. 2g). These results indicate that the deletion of the EAR motif abolishes both gene repression and activation functions of BZR1.

To examine the possibility that the deletion of the EAR motif affects the phosphorylation of BZR1, we checked the phosphorylation status of BZR1 and *BZR1ΔEAR* before and after BR treatment. Wild-type BZR1 protein was mostly phosphorylated

when seedlings were grown on the medium containing PPZ and dephosphorylated after BR treatment, but *BZR1ΔEAR* was highly dephosphorylated even without BR treatment (Supplementary Fig. 2a). Such high level of dephosphorylated *BZR1ΔEAR* is likely to be responsible for the strong phenotypes of *BZR1ΔEAR-OX* plants in contrast to the lack of phenotype of *BZR1-OX* plants (Fig. 1). As the EAR motif is located near the BIN2-docking motif⁸, it is likely to be that the deletion of the EAR motif affects BZR1–BIN2 interaction. However, *BZR1ΔEAR* still interacted with BIN2 in a yeast two-hybrid assay (Supplementary Fig. 2b). Moreover, the endogenous BZR1 was also partially dephosphorylated before BR treatment in the *BZR1ΔEAR-OX* plants (Supplementary Fig. 2a), suggesting that BR signalling upstream of BZR1 is activated in the *BZR1ΔEAR-OX* plants, most likely to be due to the loss of the feedback inhibition of BR synthesis and upstream BR signalling. Indeed, quantitative reverse transcription–PCR (qRT–PCR) analysis showed that the expression level of *BIN2* is decreased in *bzr1-1DΔEAR-OX* compared with that in wild type (Supplementary Fig. 2c), which probably contributes to the decreased BZR1 phosphorylation.

SRDX fusion restores the functions of EAR motif-deleted BZR1. An EAR motif-based artificial transcriptional repression domain (LDDLLELRGFA), named SRDX, confers a dominant transcriptional repression activity to the linked transcription

factor and has been widely used in functional studies of transcription factors in plants²⁴. To determine whether SRDX provides the same function as the EAR motif in BZR1, we generated transgenic plants overexpressing *bzr1-1DΔEAR* fused with SRDX at the C terminus (*bzr1-1DΔEAR-SRDX-OX*) (Fig. 3a). On the medium containing PPZ, the hypocotyls of wild type and *bzr1-1DΔEAR-OX* were much shorter than that of *bzr1-1D*, indicating the inability of *bzr1-1DΔEAR* to promote hypocotyl elongation (Fig. 3b). In contrast, *bzr1-1DΔEAR-SRDX-OX* had much longer hypocotyl than wild type (Fig. 3b). In addition, the SRDX fusion restored the BR sensitivity of *bzr1-1DΔEAR-OX* in the inhibition of primary root growth (Fig. 3c), indicating that the functions of EAR motif can be replaced by SRDX.

Consistent with the established SRDX function, the SRDX fusion restored the transcriptional repression activity of *bzr1-1DΔEAR* as shown by downregulation of *DWF4* expression in the *bzr1-1DΔEAR-SRDX-OX* plants (Fig. 3d). However, to our

surprise, *bzr1-1DΔEAR-SRDX* also activated expression of *SUAR15* and *IAA19* (Fig. 3e), which are direct targets of BZR1, suggesting that SRDX also mediates the transcriptional activation function of BZR1.

BZR1 interacts with TPL through the EAR motif. In several transcription factors, the EAR motif has been known to mediate transcriptional repression through interaction with TPL and the homologous TPL-related (TPR) proteins^{25–28}. Our recent tandem affinity purification of BZR1 complex followed by mass spectrometry analysis suggested that TPL associates with BZR1 *in vivo*²⁹. We therefore tested whether BZR1 directly interacts with TPL and TPRs. Indeed, in the yeast two-hybrid assay, BZR1 interacted with TPL and also other TPRs (Fig. 4a,c). Deletion of the EAR motif or mutation of the conserved Leu residues in the EAR motif abolished the BZR1–TPL interaction (Fig. 4a,b), indicating that BZR1 interacts with TPL through the EAR motif.




Figure 3 | SRDX fusion restores the functions of EAR motif-deleted BZR1. (a) Box diagram of BZR1, BZR1ΔEAR and BZR1ΔEAR-SRDX. (b,c) SRDX fusion restores both BZR1 promotion of hypocotyl elongation and inhibition of primary root growth. Seedlings were grown in the dark for 5 days on the medium containing 2 μM PPZ (b) or grown under light for 5 days on the medium containing either mock or 1,000 nM BL (c). The error bars indicate the s.d. ($n=10$ plants). Scale bar, 5 mm. (d,e) qRT-PCR analyses of BZR1-regulated genes. RNA was extracted from seedlings grown in the dark for 5 days on the medium containing 2 μM PPZ. Gene expression levels are normalized to that of *PP2A* and are shown relative to the expression levels in wild type. The error bars indicate the s.d. ($n=3$).




Figure 4 | BZR1 interacts with TOPLESS through the EAR motif. (a) A yeast two-hybrid assay shows the interaction between BZR1 and TPL. Yeast clones were grown on the synthetic dropout (+ His) or synthetic dropout without histidine (-His) plus 50 mM 3AT medium. (b) Mutation in the EAR motif abolishes the BZR1-TPL interaction. Core Leu residues of the EAR motif were substituted to Ala. Yeast clones were grown on the synthetic dropout (+ His) or synthetic dropout without histidine (- His) plus 1 mM 3AT medium. (c) BZR1 interacts with TPL and other TPRs. Yeast clones were grown on the synthetic dropout (+ His) or synthetic dropout without histidine (- His) plus 5 mM 3AT medium. TPL-N, N-terminal domain of TPL (1-344); TPR1-N, 1-343; TPR2-N, 1-335; TPR3-N, 1-335; and TPR4-N, 1-344. (d) BZR1 interacts with TPL through EAR motif *in vivo*. *Arabidopsis* mesophyll protoplasts were transfected with TPL-Myc together with *bzr1-1D-GFP* or *bzr1-1DΔEAR-GFP*, and the extracted proteins were immunoprecipitated by anti-GFP antibody. Gel blots were probed with anti-Myc or anti-GFP antibody. (e) TPL binds to *DWF4* promoter through BZR1. GST-TPL-N (amino acids 1-344) was incubated with a biotin-labelled *DWF4* promoter DNA (400 bps) together with MBP or MBP-BZR1 and pulled down with streptavidin agarose beads. Gel blots were probed with anti-GST or anti-MBP antibody. The full scans of the gel blots (d,e) are shown in Supplementary Fig. 4. (f) The TPL DNA binding on the promoters of *CPD* and *DWF4* are enhanced by BR treatment. TPL DNA-binding was determined by ChIP assay using *TPLp::TPL-HA* transgenic plants. One-week seedlings grown on the medium containing 2 μM PPZ were treated with mock (M) or 100 nM BL for 4 h before crosslinking. Error bars indicate the s.d. ($n=3$) and ** $P<0.01$ by Student's *t*-test.

To confirm the BZR1-TPL interaction *in vivo*, we performed co-immunoprecipitation (co-IP) assays using *Arabidopsis* mesophyll protoplasts expressing *bzr1-1D-GFP* or *bzr1-1DΔEAR-GFP* and TPL-Myc. The TPL-Myc protein was co-immunoprecipitated by

anti-YFP antibody specifically when *bzr1-1D-GFP* was co-expressed (Fig. 4d), confirming the *in vivo* BZR1-TPL interaction. The deletion of EAR motif greatly reduced the TPL-BZR1 interaction *in vivo* (Fig. 4d), which is consistent with

the result of the yeast two-hybrid assay (Fig. 4a). Thus, both yeast two-hybrid and co-IP assays demonstrate that BZR1 interacts with TPL through the EAR motif.

TPL is not a DNA-binding protein, but it is usually recruited to specific promoter DNA through interaction with DNA-binding transcription factors. To test whether BZR1 recruits TPL to the promoters of BZR1 target genes, we first carried out DNA pull-down assays with a biotin-labelled *DWF4* promoter. As expected, TPL was pulled down by the *DWF4* promoter DNA only in the presence of BZR1 but not when BZR1 was absent (Fig. 4e). Next, we carried out chromatin immunoprecipitation (ChIP) assays using the transgenic plants expressing TPL-HA driven by the native *TPL* promoter, to test the TPL binding to the promoters of BZR1 targets *in vivo*. Consistent with the results of the DNA pull-down assays, TPL is associated with the promoters of BZR1-repressed genes (*DWF4* and *CPD*) *in vivo* and the TPL binding to these promoters was enhanced by BR treatment (Fig. 4f), which is known to induce BZR1 DNA binding⁵. TPL also bound to the

promoters of BR-activated genes (*SAUR15* and *IAA19*), but unlike the binding to *DWF4* and *CPD* promoters the TPL binding to *SAUR15* and *IAA19* promoters was not enhanced by BR treatment (Fig. 4f). TPL is likely to be also recruited to these promoters through other BR-independent transcription factors, such as the AUX/IAA proteins, which also interact with TPL and mediate auxin-responsive expression of *SAUR15* and *IAA19* (refs 27,30). Together, these results indicate that through its EAR motif, BZR1 recruits TPL to the promoters of BR-repressed genes.

TPL is required for BR responses. To further verify the requirement of TPL/TPRs activities in BR responses, we tested the BZR1-repressed genes in the *tpl; tpr1; tpr4* triple mutant lacking *TPL* and two *TPR* genes²⁸. As expected, *bzr1-1D* caused a much milder repression of *CPD* and *DWF4* in *tpl; tpr1; tpr4* than in wild-type background (Fig. 5a), supporting that the full TPL/TPRs activities are required for BZR1 repression of these target




Figure 5 | TOPLESS is required for BR responses. (a) The BZR1 transcriptional repression activity is compromised in *tpl;tpr1;tpr4*. Total RNA were extracted from etiolated seedlings grown on the medium containing 2 μ M PPZ. The relative gene expression levels were determined by qRT-PCR. Numbers indicate the ratios of expression levels (*bzr1-1D*/Col-0 or *bzr1-1D;tpl;tpr1;tpr4* / *tpl;tpr1;tpr4*). Error bars indicate the s.d. ($n = 3$). (b) The *tpl;tpr1;tpr4* triple mutant is less sensitive to BR. Seedlings were grown for 5 days on the medium containing either mock or 1,000 nM BL. Numbers indicate ratio of hypocotyl lengths (BL-treated/mock-treated). Error bars indicate the s.d. ($n = 10$ plants) and $**P < 0.01$ by Student's *t*-test. Scale bar, 5 mm. (c) The *tpl;tpr1;tpr4* triple mutation suppresses the long hypocotyl phenotype of *bzr1-1D*. Seedlings were grown on the medium containing 2 μ M PPZ for 6 days in the dark. Error bars indicate the s.d. ($n = 10$ plants). n.s., not significant and $**P < 0.01$ by Student's *t*-test. Scale bar, 5 mm. (d) Scatter plot of fresh weight measurements of 3-week-old plants grown on the medium containing 2 μ M PPZ. This experiment was repeated with similar results. $**P < 0.01$ by Student's *t*-test ($n = 10$ plants). (e) TPL fusion at the C terminus restores the transcriptional repression activity of *bzr1-1DmEAR*. Relative expression fold changes of *CPD* and *DWF4* were determined by qRT-PCR in *Arabidopsis* plants transformed with the indicated constructs, compared with wild type (Col-0). Error bars indicate the s.e. ($n = 4$). (f) TPL fusion restores the activity of *bzr1-1DmEAR* in promoting hypocotyl elongation. Seedlings were grown in the dark on the medium containing 1 μ M BRZ for 5 days. Error bars indicate the s.e. ($n = 10$).

genes. In contrast, *bzr1-1D* activation of *IAA19* and *EXP8* was not significantly affected by the *tpl; tpr1; tpr4* mutation, and *SAUR15* activation by *bzr1-1D* was only slightly decreased in *tpl; tpr1; tpr4* (Supplementary Fig. 3). These results indicate that TPL/TPRs play a major role in BZR1 repression of target genes but not in BZR1 activation of gene expression. Consistent with the defect in the BZR1 repression of gene expression, the *tpl; tpr1; tpr4* seedlings were significantly less sensitive to BR treatment (Fig. 5b) in hypocotyl elongation compared with wild type. In addition, the *tpl; tpr1; tpr4* triple mutation partially suppressed the PPZ-insensitive phenotypes of the *bzr1-1D* mutation both in the dark and under light (Fig. 5c,d). These results demonstrate that the normal level of TPL/TPR activity is required for BZR1-mediated gene repression and hypocotyl cell elongation.

The BR-related phenotypes of the *tpl; tpr1; tpr4* triple mutant are weaker and likely to be also less specific than typical BR-insensitive mutants, because of genetic redundancy and broad functions of TPL/TPRs as general transcription repressors. To further confirm the essential and sufficient role of the BZR1–TPL interaction in the BR repression of gene expression, we generated transgenic plants overexpressing *bzr1-1DmEAR* (mutation in the EAR motif) and *bzr1-1DmEAR-TPL* fusion protein. The mutation of the core Leu residues in the EAR motif (mEAR) abolished the BZR1 interaction with TPL (Fig. 4b) and the BZR1 transcriptional repression activities on *DWF4* and *CPD* (Fig. 5e). TPL fusion to the C terminus of *bzr1-1DmEAR* restored the BZR1 transcriptional repression activities (Fig. 5e) as well as the hypocotyl promoting activity (Fig. 5f), indicating that the TPL recruitment through the EAR motif is necessary and sufficient for these BZR1 functions.

TPL/TPRs modulate light sensitivity. The BR repression of gene expression has been shown to mediate the BR inhibition of light-induced seedling morphological changes, photomorphogenesis^{31,32}. BR-activated BZR1 directly binds to the promoters of *GATA4* and *BZS1*, which are positive regulators of photomorphogenesis, and represses their expression levels^{31,32}. As TPL/TPRs are involved in the BZR1 repression of gene expression, we tested whether TPL/TPRs are also required for BZR1 to repress *GATA4* and *BZS1*. First, we showed, using ChIP assays, that TPL was recruited to the promoters of *GATA4* and

BZS1 in a BR-dependent manner (Fig. 6a). Consistently, BZR1 repression of *GATA4* and *BZS1* expression was diminished in the *tpl; tpr1; tpr4* mutant compared with wild type (Fig. 6b). Next, we tested whether loss of TPL/TPRs affects light response of seedlings. Figure 6c shows that *tpl; tpr1; tpr4* is hypersensitive to the red light inhibition of hypocotyl elongation, which is consistent with the high expression levels of *GATA4* and *BZS1*. Together, these results suggest that BZR1–TPL/TPRs interactions modulate the light sensitivity of seedlings through the transcriptional regulation of *GATA4* and *BZS1*.

HDA mediates BR repression of gene expression. The Groucho/TLE-mediated transcription repression has been known to engage HDA³³. In addition, previous studies have shown that TPL interacts with HDA6 and HDA19 (refs 25,28), and our proteomic study has identified HDA19 in the BZR1 complex²⁹. We confirmed that BZR1 associates with both HDA19 and TPL *in vivo* (Fig. 7a). These results suggested that the HDA activity may be required for the BR repression of gene expression, and thus we tested this hypothesis using a HDA inhibitor Trichostatin A (TSA). Although BR treatment dramatically repressed expression levels of *DWF4* and *CPD*, their expression levels were elevated by TSA treatment and much less affected by BR in the presence of TSA (Fig. 7b), indicating that the HDA activity is required for BR repression of gene expression. Consistent with the gene expression pattern, the long-hypocotyl phenotype of *bzr1-1D* was largely suppressed by the TSA treatment, whereas the wild-type seedlings were much less affected under this condition (Fig. 7c). These results support the notion that the BZR1–TPL interaction exploits HDA for the BR repression of gene expression and promotion of cell elongation.

BZR1 directly represses the expression of TPL and TPRs. The genome-wide analyses of BZR1 direct targets showed that *TPL* and *TPRs* are direct targets of BZR1 (Fig. 8a)¹⁵. We performed independent ChIP assays using transgenic plants expressing *bzr1-1D-CFP* under the control of native *BZR1* promoter (*BZR1p::bzr1-1D-CFP*). Consistent with the ChIP-Seq result, the promoter regions of *TPL*, *TPR1*, *TPR2* and *TPR3* were significantly enriched by the BZR1 ChIP compared with the negative control *PP2A* and *UBC30* promoter regions (Fig. 8b),




Figure 6 | TPL/TPRs modulate light sensitivity. (a) TPL binds to the promoters of *GATA4* and *BZS1* in a BR-dependent manner. The TPL–DNA binding was determined by ChIP assay using *TPLp::TPL-HA* transgenic plants. One-week-old seedlings grown on the medium containing 2 μM PPZ were treated with mock (M) or 100 nM BL for 4 h before cross-linking. Error bars indicate the s.d. ($n=3$) and $**P<0.01$ by Student's *t*-test. (b) The BZR1 transcriptional repression of *GATA4* and *BZS1* are compromised in the *tpl; tpr1; tpr4* triple mutant. Total RNA were extracted from etiolated seedlings grown on the medium containing 2 μM PPZ. The relative expression fold changes of *GATA4* and *BZS1* were determined by qRT–PCR. Error bars indicate the s.d. ($n=3$) and $**P<0.01$ by Student's test. (c) The *tpl; tpr1; tpr4* seedlings are hypersensitive to red light. Seedlings were grown in various intensities of red light for 5 days. Numbers indicate the percentages of average hypocotyl length relative to that of dark-grown seedling. Error bars indicate the s.d. ($n=10$).




Figure 7 | BR repression of gene expression requires HDA activity. (a) BZR1 associates with both TPL and HDA19 *in vivo*. *Arabidopsis* mesophyll protoplasts were transfected with *bZR1-1D-GFP* together with *TPL-Myc* and *HDA19-Myc*, and the extracted proteins were immunoprecipitated by anti-GFP antibody. Gel blots were probed with anti-Myc or anti-GFP antibody. (b) HDA inhibitor, trichostatin A (TSA) diminishes BR repressive effect on the expression of *DWF4* and *CPD*. Total RNA was extracted from 2-week-old plants treated with 100 nM BL ± 10 μM TSA for 1 h. The relative gene expression levels were determined by qRT-PCR. Error bars indicate the s.d. (n = 3). Numbers indicate the ratios of expression levels (BL/M or TSA + BL/TSA). (c) TSA treatment inhibits BZR1 promotion of hypocotyl elongation. Seedlings were grown on the medium containing 2 μM PPZ with or without 1 μM TSA in the dark. The error bars in the lower graph indicate the s.d. (n = 10 plants). Scale bar, 5 mm.




Figure 8 | BZR1 represses the expression of TPL and TPRs. (a) ChIP-Seq analyses shows BZR1 binding on the promoters of *TPL* and *TPRs*. Arrows indicate the direction of transcription. (b) ChIP assays confirm that BZR1 binds to promoters of *TPL* and *TPRs*. Transgenic plants expressing *bZR1-1D-CFP* from the native *BZR1* promoter were used for the ChIP assays. BZR1-bound DNA was immunoprecipitated using anti-GFP antibody and analysed by qPCR. BZR1 DNA binding was calculated as the ratio between *bZR1-1D-CFP* and wild-type samples, normalized to that of *PP2A*. (c) The expression levels of *TPL* and *TPR3* are reduced by BR treatment. Seedlings grown on the medium containing 2 μM PPZ medium for 7 days were treated with mock (M) or 100 nM BL for 4 h. (d) *TPR3* repression by BZR1 requires *TPL/TPR* activity. Total RNA were extracted from etiolated seedlings grown on the medium containing 2 μM PPZ. The relative gene expression levels were determined by qRT-PCR. Numbers indicate the ratios of expression levels (*bZR1-1D/Col-0* or *bZR1-1D;tpl;tp1;tp4/tpl;tp1;tp4*). The error bars in **b-d** indicate the s.d. (n = 3). **P < 0.01 and *P < 0.05 by Student's *t*-test.

confirming that BZR1 binds to these *TPL/TPR* promoters. BR treatment reduced the expression level of *TPR3* by twofold, *TPL* by ~40% and *TPR1* slightly, but had no significant effect on *TPR2* expression (Fig. 8c), which correlate with their relative binding by BZR1 detected by ChIP-qPCR (Fig. 8b). These results suggest that BZR1 mediates repression of *TPL* and at least two *TPRs*. In addition, BZR1 repression of *TPR3* also requires *TPL/TPRs* activities, as the expression level of *TPR3* was elevated and much less repressed by *bZR1-1D* in the *tpl;tp1;tp4* mutant background (Fig. 8d), suggesting that BZR1-TPL/TPRs interactions mediate a feedback repression of many members of the *TPL/TPR* gene family.

Discussion

BZR1 is a master transcription factor in BR signalling pathway and regulates expression of thousands of genes to elicit various BR responses¹⁵. Although BR signalling pathway upstream of

BZR1 has been relatively well characterized, the molecular mechanism underlying BZR1 regulation of gene expression is still poorly understood. Here we show that the EAR motif at the C terminus of BZR1 is essential for BZR1 regulation of transcription and plant growth, and the EAR motif mediates BR/BZR1 repression of gene expression by recruiting the co-repressor TPL and HDAs.

EAR motifs of several plant transcription factors have been shown to act as a transcription repression domain by recruiting the Groucho/TLE-like transcriptional corepressor TPL³⁴. We found the EAR motif is highly conserved in the BZR1 family of transcription factors across land plants and it plays a central role in BZR1's transcriptional functions. Specific deletion or mutations of the EAR motif abolished BZR1's abilities to regulate gene expression and a wide range of developmental processes, demonstrating an essential role of the EAR motif in BZR1 function.

The series *in vitro* and *in vivo* experiments demonstrated that the EAR motif of BZR1 mediates interaction with TPL. The BZR1–TPL interaction was detected in yeast two-hybrid, co-IP and *in vitro* pull-down assays. TPL was also identified as a component of the BZR1 complex by affinity purification followed by mass spectrometry analyses²⁹. Deletion of the EAR motif abolished the interaction in both yeast two-hybrid and co-IP assays. Both *in vitro* DNA-binding assays and ChIP assays demonstrated that BZR1 recruits TPL to the promoters of BR-repressed genes. The essential role for TPL/TPRs in BR repression of gene expression is supported by several lines of evidence. First, BZR1 showed significantly reduced ability to repress several BR-repressed genes in the triple *tpl; tpr1; tpr4* mutant compared with wild-type plants. The weak phenotypes of the *tpl; tpr1; tpr4* mutant are likely to be due to redundant functions of other members of the TPL/TPR family, as *TPR3* expression is increased in the *tpl; tpr1; tpr4* mutant and yeast two-hybrid assays indicated that BZR1 interacts with all four TPRs. Second, the TPL fusion to the *bzr1-1DmEAR* restored the defect of *bzr1-1DmEAR* in repressing gene expression and promoting hypocotyl elongation, demonstrating that BZR1 interaction with TPL is essential and sufficient for repressing gene expression and promoting cell elongation. Furthermore, TPL and three TPR genes are direct targets of BZR1, and three of them are repressed by BR treatment under our conditions, supporting complex roles of TPL/TPRs in BR responses.

Given that TPL/TPRs are positive regulators in BR responses, the transcriptional repression of TPL/TPRs by BZR1 would provide a negative feedback loop that enables fine-tuning the BR responses. Furthermore, as the BZR1-mediated repression requires TPL/TPRs proteins, this creates another feedback loop that helps maintain the cellular level of TPL/TPRs. As TPL/TPRs are exploited by multiple hormonal and developmental pathways (auxin, jasmonic acid, abscisic acid and ethylene) to repress gene expression^{26–28,35–37}, regulation of TPL/TPRs expression levels by BZR1 potentially allows BR/BZR1 to modulate other signalling pathways. Therefore, TPL/TPRs play important roles in BR responses not only as the transcription repressors of specific BR target genes but also as BR target genes themselves that potentially mediate feedback regulation of BZR1 function and modulate broader secondary gene expression responses.

The TPL-induced transcriptional repression has been reported to be at least partly mediated by HDA³⁴. A role of HDA in the BZR1 repression of gene expression is supported by the effect of HDA inhibitor TSA on *bzr1-1D* repression of *CPD* and *DWF4* expression, and BZR1 association with both TPL and HDA19 *in vivo*. Together, these results demonstrate that BZR1 recruits TPL and HDA to the BR-repressed promoters to mediate transcriptional repression through chromatin modification.

The transcriptional repression activity of BZR1 is important not only for feedback regulation of BR homeostasis but also for BR promotion of plant growth⁵. As essential components for BZR1-mediated transcriptional repression, TPL/TPRs play a key role in the biological functions of BR. In addition to the BR biosynthetic genes, two BR-repressed genes encoding positive regulators of photomorphogenesis, *GATA4* and *BZS1* (refs 31,32), showed BR-induced TPL promoter binding and TPL/TPR-dependent transcriptional repression. Consistent with TPL/TPR's function in BZR1-mediated repression of *GATA4* and *BZS1*, the triple *tpl; tpr1; tpr4* mutant showed not only reduced sensitivity to BR treatment but also increased sensitivity to red light treatment in hypocotyl elongation, which indicate that TPL/TPRs play an important role in BR regulation of growth and photomorphogenesis. The function of TPL in BR promotion of growth is further confirmed by the recovery of growth promoting activity of *bzr1-1DmEAR* with TPL fusion. Together, these results

provide strong evidence supporting a central role for the TPL-mediated transcriptional repression activity of BZR1 in BR regulation of cell elongation and photomorphogenesis, as well as in the feedback inhibition of BR biosynthesis.

Although the EAR motif has been shown to mediate transcription repression in several plant transcription factors as well as in BZR1, our study provides evidence that the EAR motif of BZR1 may also mediate transcriptional activation. First, the expression of BZR1 direct target genes that are activated by BR and *bzr1-1D* were not activated by *bzr1-1DΔEAR* in transgenic plants. Second, when *bzr1-1DΔEAR-GR* was induced to translocate to the nucleus, the expression level of *SAUR15* was not increased, in contrast to *bzr1-1D-GR* that significantly activated *SAUR15* expression. Third, *bzr1-1DΔEAR* did not activate the *PRE5* promoter activity in the transient gene expression assay. Last, the artificial SRDX domain recovered the defect in transcriptional activation of *bzr1-1DΔEAR*. SRDX contains a different EAR motif from that of BZR1 (LDLDELRLGFA instead of LELTL), and thus the results suggest that the loss of activation function in *bzr1-1DΔEAR* is due to the loss of the EAR motif rather than another overlapping domain. The EAR motif exists in a large number of transcription factors and whether the EAR motif can mediate transcription activation in other factors is an important question to be answered in future studies. Our observation also cautions the interpretation of results obtained with SRDX fusion transcription factors.

How the transcriptional activation and repression activities of BZR1 are determined? Apparently, the *cis*-elements of the promoter specify BZR1's transcriptional activity by probably recruiting different co-factors. A recent study showed that a MYB transcription factor MYBL2 is a transcription repressor and interacts with BZR2/BES1 (ref. 20). Interestingly, MYBL2 has an internal LxLxL motif³⁵ but instead uses a C-terminal TLLFR motif for transcription repression³⁸. Nevertheless, MYBL2 was identified as a TPL-interacting protein in the yeast two-hybrid screen³⁵ and thus it is possible that MYBL2 stabilizes BZR1/BZR2 interactions with TPL and thereby enhances transcriptional repression. Another BZR2/BES1-interacting transcription factor HAT1 also represses gene expression, but loss-of-function *hat1; hat2* mutant showed normal fold change of BR-responsive gene expression²¹, and thus HAT1 appears not essential for BZR2/BES1 transcriptional repression. Alternatively, transcriptional repression might be the default activity of BZR1 in the absence of co-activator that blocks the EAR motif, as suggested by the repression of artificial promoter by the BZR1 fusion to the LexA DNA-binding domain⁵. It is conceivable that the EAR motif in BZR1 must be blocked and inaccessible to TPL at the promoter to be activated by BZR1. Consistent with this notion, BR treatment increased TPL association with BR-repressed promoters but not BR-activated promoters. It is possible that the EAR motif in BZR1 is involved in interaction with co-activators at the BR-activated promoters, to both mediate gene activation and prevent recruitment of TPL repressor. BZR1 activation of transcription appears to involve interaction with bHLH factors. Both BZR1 and BZR2/BES1 interact with bHLH factors of the PIF family¹⁶, and their shared target genes are significantly more likely to be activated than repressed by BR. BZR2/BES1 has also been shown to interact with another bHLH factor BIM1 and histone-modifying enzymes to mediate the transcriptional activation^{13,17,19}. Whether the EAR motif of BZR1 is involved in the interactions with these co-transcription activators remains to be determined in future studies.

Transcription factors with dual functions of activation and repression are common in both plants and animals. For example, WUSCHEL activates *AGAMOUS* expression but represses the

expression of *ARR6* and *ARR7* (ref. 39). In metazoan, the T-cell factor (TCF) transcription factors mediate both repression and activation of target gene expression. When the Wnt signalling pathway is off, TCF interacts with Groucho/TLE corepressor to repress its target genes. Wnt signalling inactivates GSK3, leading to accumulation of β -catenin in the nucleus; β -catenin displaces Groucho/TLE from TCF and re-assembles a transcription activator complex, switching from repression to activation⁴⁰. In this case, TCF bound to the same promoter is switched from a repressor to an activator by exchange of interacting proteins. The mode of action of BZR1 appears more similar to that of dorsal, which represses some target genes by recruiting Groucho/TLE but interacts with the bHLH factor twist to activate target genes that contain twist-binding sites^{41–43}. Unlike BZR1's EAR motif, which is required for both repression and activation, the Groucho/TLE-interaction domain at the C terminus of dorsal is required for transcription repression but not activation. It seems that BZR1 shares features of repressor displacement with TCF, *cis*-element-dependent co-activator interaction with dorsal and inactivation by GSK3 with β -catenin. Furthermore, both BZR1 and dorsal are targets of leucine-rich repeat receptor-mediated signalling pathways^{2,3}. Therefore, the BR signalling pathway shares striking similarities with the Wnt and Toll pathways of metazoan, while mechanistic details seem to have evolved to provide appropriate efficiency required for the different levels of developmental plasticity of the organisms.

Methods

Plant materials and growth conditions. *Arabidopsis thaliana* plants were grown in a greenhouse with a 16-h light/8-h dark cycle at 22–24 °C for general growth and seed harvesting. All the plants used in this study are in Col-0 ecotype background. The *tpl1;tpl2;tpl3* triple mutant was provided by Zhang and coworkers²⁸. The *TPLp::TPL-HA* plants used for ChIP assay was provided by Somers and coworkers²⁵.

Constructs. The full-length complementary DNA of *bzr1-1D* (also *bzr1-1DAEAR*) was fused to GR by overlapping PCR. The resulting *bzr1-1D-GR* and *bzr1-1DdEAR-GR* were inserted into pENTR/SD/D-TOPO vector (Invitrogen) and subcloned into gateway compatible pEarleyGate101 vector for transformation into *Arabidopsis*.

For the *Arabidopsis* mesophyll protoplast transfections, the full-length cDNA of *TPL* and *HDA19* (without stop codon) was cloned into the gateway-compatible *pGWBI7* plasmid, and the cDNA (without stop codon) of *bzr1-1D* and *bzr1-1DAEAR* was cloned into the gateway-compatible pEarleyGate101 plasmid.

The full-length cDNA of *bzr1-1D* (*BZR1P234L*) and *bzr1-1DmEAR* (*bzr1-1D^{L328/330A}*) were cloned into plant expression vectors under the control of the 35S *CAPPDK* promoter as previously reported^{10,44}. All mutants of *BZR1* were generated using the QuickChange Site-Directed Mutagenesis kit (Stratagene) according to the manufacturer's instructions. To generate *bzr1-1DmEAR-SRDX-* and *bzr1-1DmEAR-TPL-overexpressing transgenic lines*, the *SRDX* sequence and full-length cDNA of *TPL* were tagged to *bzr1-1DmEAR* at the C terminus region and subcloned into a plant expression binary vector, *pCB302ES*⁴⁴.

Hypocotyl and root length measurements. Sterilized seeds by 70% (v/v) ethanol and 0.01% (v/v) Triton X-100 were plated on MS medium (PhytoTechnology Laboratories) supplemented with 0.75% phytoagar. After 3 days of incubation at 4 °C, seedlings were irradiated by white light for 6 h to promote germination and then incubated in specific conditions for 5–7 days. Seedlings were photocopied and hypocotyl and root lengths were measured by using ImageJ software (<http://rsb.info.nih.gov/ij>).

Arabidopsis protoplast isolation and transfection. Leaves from 4-week-old plants grown in the short photoperiod were cut to 0.5–1 mm leaf strips and digested by enzyme solution (1.5% cellulase R10; 0.4% macerozyme R10; 0.4 M mannitol; 20 mM KCl; 20 mM MES at pH 5.7; 10 mM CaCl₂; 0.1% BSA) for 3 h at room temperature after being vacuum infiltrated for 30 min in the dark. The enzyme/protoplast solution was diluted by an equal volume of W5 buffer (154 mM NaCl; 125 mM CaCl₂; 5 mM KCl; 5 mM glucose; 2 mM MES at pH 5.7). The enzyme solution containing protoplast was filtered through 75- μ m nylon mesh and the protoplasts were collected by centrifugation at 100 g for 1 min. After washing with W5 solution, the protoplasts were resuspended with W5 solution and incubated on ice for 30 min. After removing W5 by centrifugation at 100 g for 30 s, the protoplasts were resuspended with MMG solution (0.4 M mannitol; 15 mM MgCl₂;

4 mM MES at pH 5.7). For DNA transfection, an equal volume of PEG buffer (40% PEG4000; 0.2 M mannitol; 0.1 M Ca(NO₃)₂) was mixed with the protoplasts and DNA, and four volumes of W5 buffer were added to stop the transfection after 15 min. The resulting solution was mixed well by gently inverting the tube and the protoplasts were pelleted by centrifugation at 100 g for 1 min. The transfected protoplasts were resuspended gently in 0.5 ml of WI solution (0.5 M mannitol; 20 mM KCl; 4 mM MES at pH 5.7) and cultured under weak light at room temperature overnight.

Co-immunoprecipitation assays. About 2×10^4 isolated *Arabidopsis* mesophyll protoplast were transfected with a total 20 μ g of DNA and incubated overnight. Total proteins were extracted from the protoplasts using the IP buffer (50 mM Tris-Cl pH7.5, 1 mM EDTA, 75 mM NaCl, 0.1% Triton X-100, 5% glycerol, 1 mM phenylmethyl sulphonyl fluoride, 1 \times Protease Inhibitor). After centrifugation at 20,000 g for 10 min, supernatants were incubated for 1 h with 5 μ g of polyclonal anti-YFP antibody (custom-made) immobilized on protein A/G agarose beads. The beads were then washed for three times with 1 ml of IP buffer and eluted samples were analysed by immunoblotting using Anti-Myc (Cell Signaling Technology, mouse mAb, 1:5,000 dilution) and Anti-GFP (Clontech JL-8, mouse mAb, 1:5,000 dilution) antibodies.

Transient gene expression assays. Isolated *Arabidopsis* mesophyll protoplasts (2×10^4) were transfected with a total 20 μ g of DNA and incubated overnight. Protoplasts were harvested by centrifugation and lysed in 50 μ l of passive lysis buffer (Promega). Firefly and *Renilla* (as an internal standard) luciferase activities were measured by using a dual-luciferase reporter kit (Promega). The reporter promoters (1 kb of *CPD* promoter and 1.5 kb of *PRE5* promoter) were cloned upstream of the firefly luciferase gene in *pGreen-0800II-luc* vector, which also contains the 35S promoter—*Renilla* luciferase gene. Statistical significance was determined by Student's *t*-test.

qRT-PCR gene expression analysis. Total RNA was extracted by using the Spectrum Plant Total RNA kit (Sigma). M-MLV reverse transcriptase (Fermentas) was used to synthesize cDNA from the RNA. Quantitative real-time RT-PCR was performed in LightCycler 480 (Roche). *PP2A* was used to as an internal control. Gene-specific primers are listed in Supplementary Table 1.

Chromatin immunoprecipitation assays. The plants grown under white light for 2 weeks or in the dark for 5 days were harvested and cross-linked for 20 min in 1% formaldehyde under vacuum. The chromatin complex was isolated, resuspended in lysis buffer (50 mM Tris-HCl pH 8.0, 10 mM EDTA, 200 mM NaCl, 0.5% Triton X-100, 1 mM phenylmethyl sulphonyl fluoride) and sheared by sonication to reduce the average DNA fragment size to around 500 bps. The sonicated chromatin complex was immunoprecipitated by affinity-purified polyclonal anti-YFP antibodies (5 μ g, custom-made) or anti-HA antibody (Roche; 3F10), which was incubated with protein A agarose/salmon DNA (Millipore) overnight. The beads were washed with low-salt buffer (50 mM Tris-HCl at pH 8.0, 2 mM EDTA, 150 mM NaCl, 0.5% Triton X-100), high-salt buffer (50 mM Tris-HCl at pH 8.0, 2 mM EDTA, 500 mM NaCl, 0.5% Triton X-100), LiCl buffer (10 mM Tris-HCl at pH 8.0, 1 mM EDTA, 0.25 M LiCl, 0.5% NP-40, 0.5% deoxycholate) and TE buffer (10 mM Tris-HCl at pH 8.0, 1 mM EDTA), and eluted with elution buffer (1% SDS, 0.1 M NaHCO₃). After reverse cross-linking, the DNA was purified with a PCR purification kit (Thermo Scientific) and analysed by ChIP-qPCR. Enrichment of DNA was calculated as the ratio between transgenic plants and wild-type control, normalized to that of the *PP2A* coding region as an internal control. Statistical significance was determined by Student's *t*-test. Primers for qPCR are listed in Supplementary Table 1.

DNA pull-down assays. GST-TPL-N (amino acids 1–344 of TPL cloned into *pDEST15* vector), MBP and MBP-BZR1 (*pMAL-C2*) proteins were expressed in BL21 codon plus *Escherichia coli* cells and purified either using glutathione beads (GE Healthcare) or amylose resin (NEB) according to manufacturer's protocols. The *DWF4* promoter fragment was amplified by PCR using the biotin-labelled primers (Supplementary Table 1). GST-TPL-N was incubated with the biotin-labelled DNA together with MBP or MBP-BZR1 protein, and then DNA-binding proteins were pulled down by streptavidin agarose beads (Sigma) and analysed by immunoblotting.

References

1. Wang, Z. Y., Seto, H., Fujioka, S., Yoshida, S. & Chory, J. BRI1 is a critical component of a plasma-membrane receptor for plant steroids. *Nature* **410**, 380–383 (2001).
2. Valanne, S., Wang, J. H. & Ramet, M. The *Drosophila* Toll signaling pathway. *J. Immunol.* **186**, 649–656 (2011).
3. Kim, T. W. & Wang, Z. Y. Brassinosteroid signal transduction from receptor kinases to transcription factors. *Annu. Rev. Plant Biol.* **61**, 681–704 (2010).

4. Clouse, S. D. Brassinosteroid signal transduction: from receptor kinase activation to transcriptional networks regulating plant development. *Plant Cell* **23**, 1219–1230 (2011).
5. He, J. X. *et al.* BZR1 is a transcriptional repressor with dual roles in brassinosteroid homeostasis and growth responses. *Science* **307**, 1634–1638 (2005).
6. Yin, Y. *et al.* BES1 accumulates in the nucleus in response to brassinosteroids to regulate gene expression and promote stem elongation. *Cell* **109**, 181–191 (2002).
7. He, J. X., Gendron, J. M., Yang, Y., Li, J. & Wang, Z. Y. The GSK3-like kinase BIN2 phosphorylates and destabilizes BZR1, a positive regulator of the brassinosteroid signaling pathway in *Arabidopsis*. *Proc. Natl Acad. Sci. USA* **99**, 10185–10190 (2002).
8. Peng, P., Zhao, J., Zhu, Y., Asami, T. & Li, J. A direct docking mechanism for a plant GSK3-like kinase to phosphorylate its substrates. *J. Biol. Chem.* **285**, 24646–24653 (2010).
9. Gampala, S. S. *et al.* An essential role for 14-3-3 proteins in brassinosteroid signal transduction in *Arabidopsis*. *Dev. Cell* **13**, 177–189 (2007).
10. Ryu, H. *et al.* Nucleocytoplasmic shuttling of BZR1 mediated by phosphorylation is essential in *Arabidopsis* brassinosteroid signaling. *Plant Cell* **19**, 2749–2762 (2007).
11. Kim, T. W. *et al.* Brassinosteroid signal transduction from cell-surface receptor kinases to nuclear transcription factors. *Nat. Cell Biol.* **11**, 1254–1260 (2009).
12. Tang, W. *et al.* PP2A activates brassinosteroid-responsive gene expression and plant growth by dephosphorylating BZR1. *Nat. Cell Biol.* **13**, 124–131 (2011).
13. Yin, Y. *et al.* A new class of transcription factors mediates brassinosteroid-regulated gene expression in *Arabidopsis*. *Cell* **120**, 249–259 (2005).
14. Vert, G. & Chory, J. Downstream nuclear events in brassinosteroid signalling. *Nature* **441**, 96–100 (2006).
15. Sun, Y. *et al.* Integration of brassinosteroid signal transduction with the transcription network for plant growth regulation in *Arabidopsis*. *Dev. Cell* **19**, 765–777 (2010).
16. Oh, E., Zhu, J. Y. & Wang, Z. Y. Interaction between BZR1 and PIF4 integrates brassinosteroid and environmental responses. *Nat. Cell Biol.* **14**, 802–809 (2012).
17. Li, L., Ye, H., Guo, H. & Yin, Y. *Arabidopsis* IWS1 interacts with transcription factor BES1 and is involved in plant steroid hormone brassinosteroid regulated gene expression. *Proc. Natl Acad. Sci. USA* **107**, 3918–3923 (2010).
18. Li, L. *et al.* *Arabidopsis* MYB30 is a direct target of BES1 and cooperates with BES1 to regulate brassinosteroid-induced gene expression. *Plant J.* **58**, 275–286 (2009).
19. Yu, X. *et al.* Modulation of brassinosteroid-regulated gene expression by Jumonji domain-containing proteins ELF6 and REF6 in *Arabidopsis*. *Proc. Natl Acad. Sci. USA* **105**, 7618–7623 (2008).
20. Ye, H., Li, L., Guo, H. & Yin, Y. MYBL2 is a substrate of GSK3-like kinase BIN2 and acts as a corepressor of BES1 in brassinosteroid signaling pathway in *Arabidopsis*. *Proc. Natl Acad. Sci. USA* **109**, 20142–20147 (2012).
21. Zhang, D. *et al.* Transcription factor HAT1 is phosphorylated by BIN2 kinase and mediates brassinosteroid repressed gene expression in *Arabidopsis*. *Plant J.* **77**, 59–70 (2013).
22. Gendron, J. M. *et al.* Brassinosteroids regulate organ boundary formation in the shoot apical meristem of *Arabidopsis*. *Proc. Natl Acad. Sci. USA* **109**, 21152–21157 (2012).
23. Gonzalez-Garcia, M. P. *et al.* Brassinosteroids control meristem size by promoting cell cycle progression in *Arabidopsis* roots. *Development* **138**, 849–859 (2011).
24. Hiratsu, K., Matsui, K., Koyama, T. & Ohme-Takagi, M. Dominant repression of target genes by chimeric repressors that include the EAR motif, a repression domain, in *Arabidopsis*. *Plant J.* **34**, 733–739 (2003).
25. Wang, L., Kim, J. & Somers, D. E. Transcriptional corepressor TOPLESS complexes with pseudoresponse regulator proteins and histone deacetylases to regulate circadian transcription. *Proc. Natl Acad. Sci. USA* **110**, 761–766 (2013).
26. Pauwels, L. *et al.* NINJA connects the co-repressor TOPLESS to jasmonate signalling. *Nature* **464**, 788–791 (2010).
27. Szemenyei, H., Hannon, M. & Long, J. A. TOPLESS mediates auxin-dependent transcriptional repression during *Arabidopsis* embryogenesis. *Science* **319**, 1384–1386 (2008).
28. Zhu, Z. *et al.* *Arabidopsis* resistance protein SNC1 activates immune responses through association with a transcriptional corepressor. *Proc. Natl Acad. Sci. USA* **107**, 13960–13965 (2010).
29. Wang, C. *et al.* Identification of BZR1-interacting proteins as potential components of the brassinosteroid signaling pathway in *Arabidopsis* through tandem affinity purification. *Mol. Cell. Proteomics* **12**, 3653–3665 (2013).
30. Overvoorde, P. J. *et al.* Functional genomic analysis of the AUXIN/INDOLE-3-ACETIC ACID gene family members in *Arabidopsis thaliana*. *Plant Cell* **17**, 3282–3300 (2005).
31. Fan, X. Y. *et al.* BZS1, a B-box protein, promotes photomorphogenesis downstream of both brassinosteroid and light signaling pathways. *Mol. Plant* **5**, 591–600 (2012).
32. Luo, X. M. *et al.* Integration of light- and brassinosteroid-signaling pathways by a GATA transcription factor in *Arabidopsis*. *Dev. Cell* **19**, 872–883 (2010).
33. Chen, G., Fernandez, J., Mische, S. & Courey, A. J. A functional interaction between the histone deacetylase Rpd3 and the corepressor groucho in *Drosophila* development. *Genes Dev.* **13**, 2218–2230 (1999).
34. Kagale, S. & Rozwadowski, K. EAR motif-mediated transcriptional repression in plants: an underlying mechanism for epigenetic regulation of gene expression. *Epigenetics* **6**, 141–146 (2011).
35. Causier, B., Ashworth, M., Guo, W. & Davies, B. The TOPLESS interactome: a framework for gene repression in *Arabidopsis*. *Plant Physiol.* **158**, 423–438 (2012).
36. Krogan, N. T., Hogan, K. & Long, J. A. APETALA2 negatively regulates multiple floral organ identity genes in *Arabidopsis* by recruiting the co-repressor TOPLESS and the histone deacetylase HDA19. *Development* **139**, 4180–4190 (2012).
37. Causier, B., Lloyd, J., Stevens, L. & Davies, B. TOPLESS co-repressor interactions and their evolutionary conservation in plants. *Plant Signal. Behav.* **7**, 325–328 (2012).
38. Matsui, K., Umemura, Y. & Ohme-Takagi, M. AtMYBL2, a protein with a single MYB domain, acts as a negative regulator of anthocyanin biosynthesis in *Arabidopsis*. *Plant J.* **55**, 954–967 (2008).
39. Ikeda, M., Mitsuda, N. & Ohme-Takagi, M. *Arabidopsis* WUSCHEL is a bifunctional transcription factor that acts as a repressor in stem cell regulation and as an activator in floral patterning. *Plant Cell* **21**, 3493–3505 (2009).
40. Daniels, D. L. & Weis, W. I. Beta-catenin directly displaces Groucho/TLE repressors from Tcf/Lef in Wnt-mediated transcription activation. *Nat. Struct. Mol. Biol.* **12**, 364–371 (2005).
41. Szymanski, P. & Levine, M. Multiple modes of dorsal-bHLH transcriptional synergy in the *Drosophila* embryo. *EMBO J.* **14**, 2229–2238 (1995).
42. Shirokawa, J. M. & Courey, A. J. A direct contact between the dorsal rel homology domain and Twist may mediate transcriptional synergy. *Mol. Cell Biol.* **17**, 3345–3355 (1997).
43. Ratnaparkhi, G. S., Jia, S. & Courey, A. J. Uncoupling dorsal-mediated activation from dorsal-mediated repression in the *Drosophila* embryo. *Development* **133**, 4409–4414 (2006).
44. Hwang, I. & Sheen, J. Two-component circuitry in *Arabidopsis* cytokinin signal transduction. *Nature* **413**, 383–389 (2001).

Acknowledgements

We thank Srinivas S. Gampala and Josh M. Gendron for generating the *bzs285* mutant, Yuelin Zhang for providing seeds of the *tpl1;tp1;tp4* triple mutant and David E. Somers for providing *TPLp::TPL-HA* seeds. The research was supported by a grant from NIH (R01GM066258) to Z.-Y.W. H.R. was supported by a grant from the National Research Foundation (2013R1A1A1076010) funded by the Korean Ministry of Science, ICT and Future Planning. I.H. was supported by the Next Generation Biogreen 21 programme (PJ009516).

Author contributions

E.O., J.-Y.Z. and Z.-Y.W. designed the research; E.O. and J.-Y.Z. conducted all the experiments, except that H.R. and I.H. performed the experiments of Figs 5e,f and E.O. and Z.-Y.W. wrote the manuscript.

Additional information

Supplementary Information accompanies this paper at <http://www.nature.com/naturecommunications>

Competing financial interests: The authors declare no competing financial interests.

Reprints and permission information is available online at <http://npg.nature.com/reprintsandpermissions/>

How to cite this article: Oh, E. *et al.* TOPLESS mediates brassinosteroid-induced transcriptional repression through interaction with BZR1. *Nat. Commun.* **5**:4140 doi: 10.1038/ncomms5140 (2014).