ARTICLE

Received 3 Mar 2014 | Accepted 16 May 2014 | Published 18 Jun 2014

DOI: 10.1038/ncomms5149

OPEN

Petawatt laser absorption bounded

Matthew C. Levy^{1,2}, Scott C. Wilks², Max Tabak², Stephen B. Libby² & Matthew G. Baring¹

The interaction of petawatt (10^{15} W) lasers with solid matter forms the basis for advanced scientific applications such as table-top particle accelerators, ultrafast imaging systems and laser fusion. Key metrics for these applications relate to absorption, yet conditions in this regime are so nonlinear that it is often impossible to know the fraction of absorbed light f , and even the range of f is unknown. Here using a relativistic Rankine-Hugoniot-like analysis, we show for the first time that f exhibits a theoretical maximum and minimum. These bounds constrain nonlinear absorption mechanisms across the petawatt regime, forbidding high absorption values at low laser power and low absorption values at high laser power. For applications needing to circumvent the absorption bounds, these results will accelerate a shift from solid targets, towards structured and multilayer targets, and lead the development of new materials.

¹Department of Physics and Astronomy, Rice University, Houston, Texas 77005, USA. ²Lawrence Livermore National Laboratory, Livermore, California 94551, USA. Correspondence and requests for materials should be addressed to M.C.L. (email: levy11@llnl.gov).

absorption is $f_i = 2\xi_1\xi_2^{3/2}/[\sqrt{\xi_1^2\xi_2 + 1} - \xi_1\sqrt{\xi_2}]$ and electron absorption is $f_e = [(1 - \xi_2)\sqrt{\xi_1^2\xi_2 + 1} - (1 + \xi_2)\xi_1\sqrt{\xi_2}]/[\sqrt{\xi_1^2\xi_2 + 1} - \xi_1\sqrt{\xi_2}] + O(\xi_3/\xi_1^2)$, using the convenient control parameters ξ_1 , ξ_2 and ξ_3 . Intensity and density conditions are controlled by $\xi_1 = [Zm_en_c/(2M_1n_0)]^{1/2}a_0$ for uniform interface charge state Z and ion mass M_1 . $\xi_2 = I_{\text{reflected}}/I_{\text{incident}}$ corresponds to net photon flux deposited in the laser-matter interface, and $\xi_3 = n_en_e/[n_i(M_i + Zm_e)] \ll 1$ is a small parameter exhibiting the disparate mass scales that characterize the petawatt laser-solid absorption modes.

Absorption bounds. Absorption bounds reflect the fact that solutions to the kinematic equations for f_e and f_i can become unphysical for values of f between zero and one. These bounds are derived using constrained optimization techniques⁵⁸ with $f = f_i + f_e$ as the objective function. We optimize f over ξ_2 imposing the simple constraint that the electron energy is real, and the minimization equation is written as $f_* = \text{Min}(f_e + f_i)$, s.t. $\gamma_e \geq 1$. Because the utility function is nonlinear in the control variables, minimization is performed numerically by means of cylindrical algebraic decomposition⁵⁹, and the resulting points are fit to a polynomial using interval $\Delta\xi_1 \approx 10^{-3}$ over the physically relevant range in ξ_1 between $[0.01, 0.5]$. For the fully ionized laser-plasma interface, we calculate that $f_* \approx 1.9\xi_1 - 2.75\xi_1^2 + 1.91\xi_1^3$, indicating that the lower limit on laser absorption is closely related to the process of ion acceleration by an intense circularly polarized radiation pressure source^{3,37,60}. Deviations from the absorption associated with this process occur at small ξ_1 as energy is reapportioned into relativistic electrons, highlighting that the kinematic coupling between ions and electrons represents an important feature of the interaction. When the electronic coupling is removed, we confirmed that f_* converges to the well-established ion acceleration result $f_* \rightarrow 2\xi_1/(1 + 2\xi_1)$ (ref. 37). Maximizing the absorption through $f^* = \text{Max}(f_e + f_i)$, again subject to the constraint that $\gamma_e \geq 1$, yields the upper limit to be

$f^* = 1 - \xi_3/(2\xi_1^2)$. In contrast to the lower limit, there is no well-established analytic result that describes absorption along the f^* curve based on a simple physical mechanism. Here we proceed allowing that $n_e \approx n_c$, implying that absorption along $f = f^*$ corresponds to electrons excited with $\gamma_e \approx a_0^2/2$, within a factor of order unity of the full laser ponderomotive potential. Figure 2 presents a comprehensive description of the absorption, showing surfaces corresponding to f_e and f_i , as well as bounding regions corresponding to f_* and f^* . Ions are seen to dominate the absorption along $f = f_*$, the region corresponding to $\gamma_e = 1$. As the target absorbs more of the laser energy, Fig. 2 shows that this energy is predominantly coupled into relativistic electrons. Electrons dominate the absorption along $f = f^*$, with $f > f^*$ causing f_e to be complex.

Comparison between absorption bounds and published data.

Comparison with published data is facilitated by specifying an interaction-averaged density, which is well-represented by a corrected relativistic critical density⁵⁵ given by $a_0^2 \approx (27/64)(n_0/n_c)^4$ for $n_0/n_c \gg 1$. Figure 3 shows these limits applied to experimental data and simulation results published over two decades, spanning a broad range of laser and plasma parameters, obtained at several laser facilities. From this figure, it is clear that experimental data and kinetic particle-in-cell simulations at a variety of realistic conditions^{37,44,48,49} show excellent agreement with the absorption bound predictions.

Absorption bounds in terms of laser and plasma parameters.

Transforming from the control coordinates, the condition $f_* \leq f \leq f^*$ can be written simply in terms of the laser power and unperturbed plasma density as,

$$\frac{\sqrt{I_1\lambda_1^2}}{\sqrt{I_1\lambda_1^2 + 1.5\sqrt{n_0}}} \leq f \leq 1 - \frac{1.2 \times 10^{18}}{I_1\lambda_1^2} \quad (1)$$

where $[I_1\lambda_1^2] = \text{W } \mu\text{m}^2 \text{ cm}^{-2}$, $[n_0] = \text{cm}^{-3}$, and $I_1\lambda_1^2 > 1.3 \times 10^{18}$. Equation (1) bounds the laser-solid interaction through its

Figure 2 | Bounds on absorption and subpartitioning of absorbed light. The lower-right axis corresponds to ξ_1 describing the initial laser and target conditions, and the total absorption $f = 1 - \xi_2$. Two surfaces corresponding to the absorption into electrons f_e and into ions f_i are shown (the former having slight transparency for visualization purposes). Contours of the optimization function γ_e are superimposed on these surfaces using dark grey. The lower limit f_* (blue) and upper limit f^* (red) on absorption are shown bounding f_e and f_i .

Figure 3 | Comparison between absorption bounds and published data. The complete data set compiled in Davies⁴⁹ is reproduced here, spanning experimental and simulation data published over the past two decades, across a variety of laser and plasma conditions. Dashed lines corresponding to fits of selected experimental data are shown to guide the eye. Additional high-intensity simulation data are reproduced from Levy *et al.*⁴⁸ The upper limit on absorption f^* is depicted in red and the lower limit f_l in blue, with forbidden regions indicated using shading. The two outlying data points correspond to simulations of (†) a very thin 0.2 μm pre-deformed target, and (††) an essentially underdense $n_e > n_0$ interaction, both violating assumptions underpinning the laser-solid model (see Methods).

dynamical history for a realistic time-dependent laser envelope and plasma profile^{37,48}.

Discussion

Figures 2 and 3 highlight that f_* becomes increasingly strict with laser power, forbidding 35% of possible absorption values at $I_l \lambda_l^2 \sim 10^{22} \text{W} \mu\text{m}^2 \text{cm}^{-2}$, a regime accessible at laser facilities such as ELI⁶¹ scheduled to come online in the next few years. These results will therefore play a central role in guiding the next generation of multi-petawatt experiments.

By shrinking the design space to be explored, these results will enable research efforts to focus on useful regions of parameter space thus accelerating the development of laser-solid applications. For these applications, the metric f/f_* using equation (1) provides a theoretical baseline enabling efficiency to be measured and ultimately improved⁶².

For applications needing to circumvent the absorption bounds in equation (1), these results will drive a shift towards new interaction paradigms. To see that assumptions underpinning the laser-solid interaction model must be violated to exceed these limits, it is instructive to examine the two outlying points shown in Fig. 3. The data point labelled (†) corresponds to one simulation of a pre-deformed, very thin target of $\lambda_l > d$ where d is the target thickness, realizing a strongly refluxing configuration (see Methods). The point labelled (††) corresponds to one simulation of a laser interacting with 20 μm of $n_e > n_0$ plasma in front of a thin $n_0 = 20n_c$ target, realizing an essentially underdense situation. We thus confirm that very thin and underdense targets allow absorption in excess of f_* at low laser power, as they should. However, several important applications that have conventionally used solid targets also depend on high absorption at relatively low laser power. These applications include laser-based anti-matter generation for scaled astrophysical studies^{63,64}, ultrafast charged-particle imaging systems²⁷, where increasing absorption reduces noise and improves imaging resolution, and certain approaches to electron-driven fast ignition laser fusion¹⁶. Recently works that

have shifted from solid targets have started to report enhanced results in these areas^{52–54}. The results presented here will accelerate this shift across the petawatt field, and lead the development of novel low density, structured and multilayer targets.

Methods

The essential kinematic relations forming the basis of the optimization analysis reported in this article were published in ref. 48. The general consideration of optimal couplings under the constraint of phase space conservation motivates the present studies⁶⁵. Radiation-hydrodynamic simulations show that particle density in interactions can often be approximated by an exponential distribution $n_0(x) \propto e^{-x/\ell_p}$ for scalelength ℓ_p , due to amplified spontaneous emission (ASE) associated with laser pulse compression generating a ‘pre-plasma’¹. Petawatt laser-solids satisfy $\ell_p [\mu\text{m}] < 1.1 a_0 \tau_l [\text{ps}]$ for pulse duration τ_l such that the primary interaction occurs in the classically overdense $n_0/n_c > 1$ region, while small-scale underdense regions are swept away by the strong laser ponderomotive force, as indicated by energy balance between electron acceleration in the underdense and overdense regions. The laser temporal envelope $I_l (\partial I_l / \partial t)^{-1} \gg 2\pi \omega_{pe}^{-1}$ and the plasma density profile are subject to $n_0 (\partial n_0 / \partial x)^{-1} \gg \ell_s$, both readily satisfied under realistic conditions. Damping of transient momentum effects requires that $\tau_l \omega_{pi} > 2\pi A$ where $\omega_{pi} = (4\pi e^2 Z n_0 / M_i)^{1/2}$ and $A \simeq 3 - 5$, and the target thickness d should exceed the hole punching depth and the effective refluxing hot electron range, $d > c\tau_l/2 + \int_0^{\tau_l} V_{int} dt$ for motion of the laser-matter interface at velocity V_{int} . Deviations from $\beta_e \cdot \hat{k}_l \approx 1$ are second order in the angle $\tan^{-1} [|\beta_e \times \hat{k}_l| / |\beta_e \cdot \hat{k}_l|]$ and therefore do not substantively affect the absorption bound results for realistic scenarios. Energy apportionment into ions increases with this angle but qualitative trends in f_e and f_i are maintained⁶⁶. For electrons following the ponderomotive energy scaling⁴², the cooling length ℓ_c associated with the radiation reaction force within ℓ_s can be estimated⁶⁷ as $\ell_c \approx 2.1 \times 10^{-2} \sqrt{n_0 [\text{cm}^{-3}]} a_0^{-7/2} \ell_s$ due to Debye shielding. Thus, $\ell_c/\ell_s \sim 1$ only at the 10-petawatt level $\sim 10^{23} \text{W} \mu\text{m}^2 \text{cm}^{-2}$, confirming the selection of absorption modes characterizing the petawatt-scale interaction.

References

- Macchi, A. A *Superintense Laser-Plasma Interaction Theory Primer* (SpringerBriefs in Physics, 2013).
- Pukhov, A. Strong field interaction of laser radiation. *Rep. Prog. Phys.* **66**, 47–101 (2003).
- Wilks, S. C. & Kruer, W. L. Absorption of ultrashort, ultra-intense laser light by solids and overdense plasmas. *IEEE J. Quantum Electron.* **33**, 1954–1968 (1997).

4. Daido, H., Nishiuchi, M. & Pirozhkov, A. S. Review of laser-driven ion sources and their applications. *Rep. Prog. Phys.* **75**, 056401 (2012).
5. Robinson, A. P. L., Arefiev, A. V. & Neely, D. Generating superponderomotive electrons due to a Non-wake-field interaction between a laser pulse and a longitudinal electric field. *Phys. Rev. Lett.* **111**, 065002 (2013).
6. Hegelich, B. M. *et al.* Laser acceleration of quasi-monoenergetic MeV ion beams. *Nature* **439**, 441–444 (2006).
7. Fuchs, J. *et al.* Laser-driven proton scaling laws and new paths towards energy increase. *Nat. Phys.* **2**, 48–54 (2005).
8. Toncian, T. *et al.* Ultrafast laser-driven microlens to focus and energy-select mega-electron volt protons. *Science* **312**, 410–413 (2006).
9. Esirkepov, T., Borghesi, M., Bulanov, S., Mourou, G. & Tajima, T. Highly efficient relativistic-ion generation in the laser-piston regime. *Phys. Rev. Lett.* **92**, 2–5 (2004).
10. Bulanov, S. V. *et al.* Unlimited ion acceleration by radiation pressure. *Phys. Rev. Lett.* **135003**, 1–4 (2010).
11. Roth, M. *et al.* Energetic ions generated by laser pulses: a detailed study on target properties. *Phys. Rev. Special Topics* **5**, 061301 (2002).
12. Sentoku, Y., Cowan, T. E., Kemp, A. & Ruhl, H. High energy proton acceleration in interaction of short laser pulse with dense plasma target. *Phys. Plasmas* **10**, 2009 (2003).
13. Flippo, K. A. *et al.* Increased efficiency of short-pulse laser-generated proton beams from novel flat-top cone targets. *Phys. Plasmas* **15**, 056709 (2008).
14. Petrov, G. M. & Davis, J. Laser acceleration of light ions from high-intensity laser-target interactions. *Appl. Phys. B* **96**, 773–779 (2009).
15. Kar, S. *et al.* Ion acceleration in multispecies targets driven by intense laser radiation pressure. *Phys. Rev. Lett.* **109**, 185006 (2012).
16. Tabak, M. *et al.* Ignition and high gain with ultrapowerful lasers. *Phys. Plasmas* **1**, 1626–1634 (1994).
17. Kodama, R. *et al.* Plasma devices to guide and collimate a high density of MeV electrons. *Nature* **432**, 1005–1008 (2004).
18. Theobald, W. *et al.* Hot surface ionic line emission and cold K-inner shell emission from petawatt-laser-irradiated Cu foil targets. *Phys. Plasmas* **13**, 043102 (2006).
19. Johzaki, T. *et al.* Pre-plasma effects on core heating and enhancing heating efficiency by extended double cone for FIREX. *Nucl. Fusion* **51**, 073022 (2011).
20. Bartal, T. *et al.* Focusing of short-pulse high-intensity laser-accelerated proton beams. *Nat. Phys.* **8**, 139–142 (2011).
21. Robinson, A. *et al.* Theory of fast electron transport for fast ignition. *Nucl. Fusion* **54**, 054003 (2014).
22. Norreys, P. *et al.* Fast electron energy transport in solid density and compressed plasma. *Nucl. Fusion* **54**, 054004 (2014).
23. Kemp, A. *et al.* Laserplasma interactions for fast ignition. *Nucl. Fusion* **54**, 054002 (2014).
24. Ryutov, D. *et al.* Similarity criteria for the laboratory simulation of supernova hydrodynamics. *Astrophys. J.* **518**, 821–832 (1999).
25. Remington, B., Drake, R. & Ryutov, D. Experimental astrophysics with high power lasers and Z pinches. *Rev. Mod. Phys.* **78**, 755–807 (2006).
26. Kugland, N. L. *et al.* Self-organized electromagnetic field structures in laser-produced counter-streaming plasmas. *Nat. Phys.* **8**, 809–812 (2012).
27. Wilks, S. C. *et al.* Energetic proton generation in ultra-intense lasersolid interactions. *Phys. Plasmas* **8**, 542 (2001).
28. Barty, C. *et al.* An overview of LLNL high-energy short-pulse technology for advanced radiography of laser fusion experiments. *Nucl. Fusion* **44**, S266–S275 (2004).
29. Zylstra, A. B. *et al.* Using high-intensity laser-generated energetic protons to radiograph directly driven implosions. *Rev. Sci. Instrum.* **83**, 013511 (2012).
30. Nakamura, T. *et al.* High-power γ -ray flash generation in ultraintense laser-plasma interactions. *Phys. Rev. Lett.* **108**, 195001 (2012).
31. Norreys, P. A. *et al.* Efficient extreme UV harmonics generated from picosecond laser pulse interactions with solid targets. *Phys. Rev. Lett.* **76**, 1832–1835 (1996).
32. Gibbon, P. *Short Pulse Laser Interactions with Matter* (Imperial College Press, 2005).
33. Gordienko, S., Pukhov, A., Shorokhov, O. & Baeva, T. Coherent focusing of high harmonics: a new way towards the extreme intensities. *Phys. Rev. Lett.* **94**, 103903 (2005).
34. Dollar, F. *et al.* Scaling high-order harmonic generation from laser-solid interactions to ultrahigh intensity. *Phys. Rev. Lett.* **110**, 175002 (2013).
35. Brunel, F. Not-so-resonant, resonant absorption. *Phys. Rev. Lett.* **59** (1987).
36. Krue, W. L. & Estabrook, K. JB heating by very intense laser light. *Phys. Fluids* **28**, 430 (1985).
37. Naumova, N. *et al.* Hole boring in a DT pellet and fast-ion ignition with ultraintense laser pulses. *Phys. Rev. Lett.* **102**, 1–4 (2009).
38. Kemp, A., Sentoku, Y. & Tabak, M. Hot-electron energy coupling in ultraintense laser-matter interaction. *Phys. Rev. Lett.* **101**, 8–11 (2008).
39. May, J. *et al.* Mechanism of generating fast electrons by an intense laser at a steep overdense interface. *Phys. Rev. E* **84**, 025401 (2011).
40. Taub, A. H. Relativistic Rankine-Hugoniot equations. *Phys. Rev.* **74**, 1–7 (1948).
41. de Hoffmann, F. & Teller, E. Magneto-hydrodynamic shocks. *Phys. Rev.* **80**, 692–703 (1950).
42. Wilks, S. C., Krue, W. L., Tabak, M. & Langdon, A. B. Absorption of ultra-intense laser pulses. *Phys. Rev. Lett.* **69**, 1383–1386 (1992).
43. Haines, M., Wei, M., Beg, F. & Stephens, R. Hot-electron temperature and laser-light absorption in fast ignition. *Phys. Rev. Lett.* **102**, 1–4 (2009).
44. Ping, Y. *et al.* Absorption of short laser pulses on solid targets in the ultrarelativistic regime. *Phys. Rev. Lett.* **100**, 6–9 (2008).
45. Robinson, A. P. L. *et al.* Relativistically correct hole-boring and ion acceleration by circularly polarized laser pulses. *Plasma Phys. Controlled Fusion* **51**, 024004 (2009).
46. Gibbon, P., Andreev, A. A. & Platonov, K. Y. A kinematic model of relativistic laser absorption in an overdense plasma. *Plasma Phys. Controlled Fusion* **54**, 045001 (2012).
47. Kemp, A. & Divol, L. Interaction physics of multipicosecond petawatt laser pulses with overdense plasma. *Phys. Rev. Lett.* **109**, 195005 (2012).
48. Levy, M. C., Wilks, S. C., Tabak, M. & Baring, M. G. Conservation laws and conversion efficiency in ultraintense laser-overdense plasma interactions. *Phys. Plasmas* **20**, 103101 (2013).
49. Davies, J. R. Laser absorption by overdense plasmas in the relativistic regime. *Plasma Phys. Controlled Fusion* **51**, 014006 (2009).
50. Zhao, Z. *et al.* Acceleration and guiding of fast electrons by a nanobrush target. *Phys. Plasmas* **17**, 123108 (2010).
51. Jiang, S., Krygier, A. G., Schumacher, D. W., Akli, K. U. & Freeman, R. R. Effects of front-surface target structures on properties of relativistic laser-plasma electrons. *Phys. Rev. E* **89**, 013106 (2014).
52. Purvis, M. A. *et al.* Relativistic plasma nanophotonics for ultrahigh energy density physics. *Nat. Photonics* **7**, 796–800 (2013).
53. Sgattoni, A., Londrillo, P., Macchi, A. & Passoni, M. Laser ion acceleration using a solid target coupled with a low-density layer. *Phys. Rev. E* **85**, 1–9 (2012).
54. Sarri, G. *et al.* Table-Top laser-based source of femtosecond, collimated, ultrarelativistic positron beams. *Phys. Rev. Lett.* **110**, 255002 (2013).
55. Cattani, F., Kim, A., Anderson, D. & Lisak, M. Threshold of induced transparency in the relativistic interaction of an electromagnetic wave with overdense plasmas. *Phys. Rev. E Stat. Phys. Plasmas. Fluids. Relat. Interdiscip. Topics* **62**, 1234–1237 (2000).
56. Denavit, J. Absorption of high-intensity subpicosecond lasers on solid density targets. *Phys. Rev. Lett.* **69** (1992).
57. Kluge, T. *et al.* Electron temperature scaling in laser interaction with solids. *Phys. Rev. Lett.* **107**, 205003 (2011).
58. Mehrotra, S. On the implementation of a primal-dual interior point method. *SIAM J. Optimization* **2**, 575–601 (1992).
59. Collins, G. E. *Quantifier elimination for Real Closed Fields by Cylindrical Algebraic Decomposition* Vol. 33. Lecture Notes in Computer Science (Springer, 1975).
60. Schlegel, T. *et al.* Relativistic laser piston model: Ponderomotive ion acceleration in dense plasmas using ultraintense laser pulses. *Phys. Plasmas* **16**, 083103 (2009).
61. ELI - extreme light infrastructure, <http://www.eli-beams.eu/> (2014).
62. Gates, B. & Gates, M. *Annual Letter*, <http://www.gatesfoundation.org/Who-We-Are/Resources-and-Media/Annual-Letters-List/Annual-Letter-2013> (2013).
63. Liang, E., Wilks, S. C. & Tabak, M. Pair production by ultraintense lasers. *Phys. Rev. Lett.* **81**, 4887–4890 (1998).
64. Chen, H. *et al.* Relativistic positron creation using ultraintense short pulse lasers. *Phys. Rev. Lett.* **102**, 1–4 (2009).
65. Fisch, N. J. & Rax, J. Free energy in plasmas under wave-induced diffusion. *Phys. Fluids B* **5**, 1754–1759 (1993).
66. Levy, M. C., Wilks, S. C., Tabak, M. & Baring, M. G. Hot electron divergence in the kinematic analysis of relativistic light on solids. *J. Phys.: Conf. Series* (2014).
67. Esarey, E., Ride, S. & Sprangle, P. Nonlinear Thomson scattering of intense laser pulses from beams and plasmas. *Phys. Rev. E* **48**, 3003–3021 (1993).

Acknowledgements

M.C.L. is grateful to B. Breizman and A. Link for discussions. M.C.L. acknowledges the LLNL Lawrence Scholarship for support and the Institutional Computing Grand Challenge for computational resources. S.C.W. acknowledges support from Fusion Energy Sciences in the DOE Office of Science. This work was performed under the auspices of the U.S. Department of Energy by Lawrence Livermore National Laboratory under Contract DE-AC52-07NA27344.

Author contributions

M.C.L. developed the analytical theory with support from S.C.W. and M.T. Assistance with conceptualizing the physical arguments and preparing the manuscript was provided by S.B.L. Theoretical support was provided by M.G.B.

Additional information

Competing financial interests: The authors declare no competing financial interests.

Reprints and permission information is available online at <http://npg.nature.com/reprintsandpermissions/>

How to cite this article: Levy, M. C. *et al.* Petawatt laser absorption bounded. *Nat. Commun.* 5:4149 doi: 10.1038/ncomms5149 (2014).

This work is licensed under a Creative Commons Attribution 4.0 International License. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in the credit line; if the material is not included under the Creative Commons license, users will need to obtain permission from the license holder to reproduce the material. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>