ARTICLE

Received 5 May 2014 | Accepted 21 Nov 2014 | Published 6 Jan 2015

DOI: 10.1038/ncomms6934

A very long-term transient event preceding the 2011 Tohoku earthquake

Yusuke Yokota^{1,†} & Kazuki Koketsu¹

Geodetic transients have been observed in various subduction zones. The 2011 Tohoku earthquake occurred in one of the most active subduction zones globally, the Japan Trench subduction zone (JTSZ). However, no geodetic transient (except afterslip and so on) had been reported in the JTSZ before the Tohoku earthquake. Here we show that a large transient event, with duration longer than any reported previously, occurred in the JTSZ preceding the Tohoku earthquake. We calculate tectonic deformations at Global Positioning System stations along the JTSZ by removing the effects of nearby M_w 6–8 earthquakes. We identify temporal changes in these deformations, deriving 9-year deviation records from regular deformations due to slip deficit at the plate boundary. We perform an inversion of the deviations to obtain the source model of their root event. The relationship between the obtained transient event and Tohoku earthquake is shown through Coulomb stress change and seismic supercycle simulation.

¹Earthquake Research Institute, University of Tokyo, Tokyo 113-0032, Japan. [†]Present address: Hydrographic and Oceanographic Department, Japan Coast Guard, Tokyo 135-0064, Japan. Correspondence and requests for materials should be addressed to K.K. (email: koketsu@eri.u-tokyo.ac.jp).

Geodetic transients have been observed in various subduction zones^{1–3} and have long been believed to be the key phenomena associated with megathrust earthquakes, although the nature of this relationship remains elusive⁴. Indeed, it has been reported that geodetic transients might have preceded megathrust earthquakes⁵. For example, long-term slow slip events (SSEs), which are those with duration longer than 10^7 s (ref. 3) were observed before the 1983 Japan Sea earthquake along the boundary of the Eurasian and North American plates⁶ and before the 2012 Ometepc earthquake in the southern Mexico subduction zone⁷. However, in one of the most active subduction zones in the world, the Japan Trench subduction zone (JTSZ), where the 2011 Tohoku earthquake^{8,9} occurred with a moment magnitude (M_w) of 9.0, no geodetic transient had been reported previously; rather, only afterslip, short-term silent earthquakes and recovery of plate coupling had been reported^{3,10}. In fact, even in the period immediately before the Tohoku earthquake, only short-term transients around the foreshock were reported^{11,12}. Nevertheless, tectonic deformations due to drag of the overriding plate coupled with the subducting plate (slip deficit) were being observed continuously in the JTSZ, and any geodetic transients that occurred in the region should have affected these tectonic deformations.

To investigate this phenomenon, we first analyse the temporal changes in tectonic deformations before the Tohoku earthquake. We use a data set from the dense Global Positioning System (GPS) network GEONET, operated by the Geospatial Information Authority of Japan (GSI). In previous studies^{13–16}, the slip deficit rate in the JTSZ was imaged using partial data sets covering the period of seismic quiescence before the 2003 Tokachi-oki

earthquake. In this study, we use the whole data set from 21 March 1996 to 8 March 2011 (one day before the foreshock).

Results

Time series of deformation. We converted the daily coordinates of the GEONET stations in the Tohoku and Kanto districts along the JTSZ to time series of horizontal deformation (see Methods). Sinusoidal annual variations and the co-seismic and post-seismic signals of nearby M_w 6–8 earthquakes are apparent in the results of this conversion (Fig. 1a,b). In addition, stationary westward trends were obtained at all stations for the seismic quiescence period before 2002 (red lines in Fig. 1b), and we assumed that these represent regular deformations that were due to slip deficit, as in all previous studies^{13–16}. We conducted an additional analysis of the time series; the results shown in Supplementary Note 1 and Supplementary Fig. 1 can also support our assumption.

We removed the annual variations and co-seismic and post-seismic signals from the time series by fitting trigonometric, Heaviside and logarithmic functions. Then, we performed regression analyses for the pre-2002 parts of the time series to determine the trends of regular slip deficit deformations and removed the obtained trends from the time series, as for the Tokai SSE¹ and Bungo Channel SSE¹⁷. If the slip deficit deformations were the only parameters affecting the results, the data should be distributed around the zero line. However, the obtained data began to deviate from the zero line around 2002 (Fig. 1c). This deviation was eastward at the central and southern stations, where movement was accelerated at the time of the 2005 M_w 7.2 Miyagi-oki earthquake, but westward at the northern stations,

Figure 1 | Time series of east-west deformation at GPS stations in the Tohoku district. (a) Selected GPS stations (orange squares) and M_w 6–8 earthquakes (green stars) in the index map. (b) Original time series of east-west deformation obtained from the GEONET F3 solutions³⁴ at the stations and the effects of the M_w 6–8 earthquakes (green lines). (c) Detrended time series obtained by removing the regular trends in 1996–2001 (solid red lines), annual variations and earthquake effects. These time series deviated from the zero lines around 2002 and accelerated at the time of the 2003 or 2005 earthquake (green lines).

Figure 2 | Distribution of total deviations and the result of a two-source inversion. The red and purple contours represent the distributions of the forward slip by the very long-term transient event and the backslip by the northern source, which were obtained through the two-source inversion of the total deviations (pink arrows). The black arrows denote synthetic deviations computed for the inversion result. The co-seismic slip distribution of the 2011 Tohoku earthquake⁸ is also displayed with the epicentre (white star) and Japan Trench (dark green line). The black bar at the bottom right denotes 100 km.

where movement was accelerated at the time of the 2003 M_w 8.3 Tokachi-oki earthquake (green lines in Fig. 1c). The total deviations at all stations were measured from the vertical coordinates on 8 March 2011 in time series diagrams such as that illustrated in Fig. 1c.

Characteristics of the sources of deviations. Figure 2 shows the distribution of measured total deviations (pink arrows). As found in deformation accelerations¹⁸ and deformations including post-seismic signals^{9,19,20}, large eastward deviations were observed along the eastern coast of central and southern Tohoku with a broad extension to the west. Meanwhile, westward deviations are obvious in the data obtained for northern Tohoku, although there was no significant deviation in the Kanto district. Small southward deviations were found extending along the western coast of northern Tohoku, as has been found previously in deformation accelerations¹⁸. This implies that the sources of the large eastward and westward deviations were located far from the western coast. Small southward deviations were also detected along the boundary between central and northern Tohoku, where the large eastward and westward deviations cancelled each other out.

On the basis of the above-mentioned evidence, we assumed the two sources to be located on deeper parts of the plate boundary beneath and off the eastern coast of central and southern Tohoku or northern Tohoku. The northern source for the westward deviations was interpreted previously to be a result of recovery of plate coupling¹⁰ or temporary acceleration of subduction²¹. These interpretations cannot be applied to the southern source, because this source generated deviations in the opposite direction. However, the deviation direction and source position relative to the Tohoku earthquake are similar to those of the Tokai SSE¹ relative to the 1854 M 8.4 Tokai earthquake and the Bungo Channel SSE¹⁷ relative to the 1707 M 8.6 Hoei earthquake. The detrended time series at stations in central and southern Tohoku from during 2002 to August 2005 and from August 2005 to March 2011 are as stationary as those of the Tokai and Bungo Channel SSEs, as shown in Fig. 1c. A more similar case was found in the Alaska subduction zone, where long-term SSEs with duration of several years were observed downdip to the source region of the 1964 M 9.2 Alaska earthquake²². Therefore, we can consider the southern source to be a phenomenon like a long-term SSE, but this would have a total duration of about 9 years, longer than any long-term SSE reported previously; thus, we refer to it as a very long-term transient event, following the terminology in simulation studies^{23,24}.

In contrast to short-term SSEs, long-term SSEs and their relatives are not always accompanied by tremor activities, although, if an episodic tremor and slip (ETS) zone exists in the vicinity of a long-term SSE, the SSE can excite tremor activity in the ETS zone⁴. However, no ETS zones have been found in the JTSZ, and the very long-term transient event was not accompanied by tremor activities.

Inversion of the deviations for finite source models. Because deformation due to slip deficit is westward and slip deficit is traditionally modelled according to a normal faulting mechanism consisting of backslips^{25,26}, the very long-term transient event for eastward deviations can be modelled according to a reverse faulting mechanism consisting of forward slips. Consequently, we located a dipping rectangular reverse fault on the deeper part of the plate boundary in central and southern Tohoku to explore the source characteristics of the very long-term transient event by analysing the deviations. However, we also had to introduce a dipping rectangular normal fault in northern Tohoku, because this northern backslip source affected deviations not only along but also beyond the boundary between northern and central Tohoku.

Then, we performed an inversion of the deviations for finite source models of the very long-term transient event and northern backslip source (see Methods). The resultant slip distribution and M_w of 7.7 are shown in Fig. 2, accompanied by those for the northern backslip source (M_w 7.4) and previous results obtained for the Tohoku earthquake (M_w 9.0) (ref. 8). The synthetic deviations (black arrows in Fig. 2) indicate that southeastward deviations to the northwest of the very long-term transient event and southwestward deviations to the southwest of the northern backslip source resulted in the small southward deviations observed along the western coast of northern Tohoku. The forward slip distribution of the very long-term transient event is slightly different from that of a previous study⁹, because we excluded post-seismic deformations of nearby M_w 6–8 earthquakes from the data, and constrained the location of the very long-term transient event to be similar to the relative locations of the Tokai and Bungo Channel SSEs as mentioned in the previous section.

The lower two time series in Fig. 1b, which were observed at the stations in central and southern Tohoku, never became flat

Figure 3 | Distribution of Coulomb failure function changes. The changes in the Coulomb failure function (ΔCFF)²⁸, due to the very long-term transient event and northern backslip source, were computed for the focal mechanism and depth of the hypocentre (white star) of the 2011 Tohoku earthquake. The zones of positive and negative ΔCFF are indicated in red and blue, respectively. The co-seismic slip distribution of the 2011 Tohoku earthquake⁸ and the Japan Trench are also displayed as black contours and the dark green line, respectively.

even after the very long-term transient event started. This means that, although the transient event was as large as M_w 7.7, the rates of deformations caused by this event never exceeded those due to the slip deficit. However, this is quite typical for a large transient event because the average slip rate is expected to be 6.3 cm per year based on the M_w and the scaling law of $v = 10^{9.5-M_w}$ mm per year (v : slip rate)²⁷. This value is only 74% of the actual subduction velocity, 8.5 cm per year.

Discussion

To examine the relationship between the very long-term transient event and the regular process consisting of the Tohoku earthquake and the slip deficit, we next computed changes in the Coulomb failure function (ΔCFF)²⁸, as was conducted previously for the Tokai SSE¹ and simulated transient events²⁹. The computations were performed for the focal mechanism and hypocentral depth of the Tohoku earthquake. A comparison of the ΔCFF distribution with the slip distribution of the Tohoku earthquake⁸ in Fig. 3 indicates that the northern half zone of the large positive ΔCFF resulting from the very long-term transient event coincides with the main rupture area of the Tohoku earthquake, except for its trenchward extension. However, the

northern backslip source distributed only negative or zero ΔCFF values throughout the rupture area of the Tohoku earthquake. Therefore, only the very long-term transient event affected the Tohoku earthquake. The ΔCFF values of the very long-term transient event, from near its upper edge to the hypocentre of the Tohoku earthquake, varied from +0.2 to +0.02 MPa, with an average of +0.1 MPa, around the mainshock hypocentre. This average is 20 times larger than the value found in the Tokai earthquake source area due to the Tokai SSE¹, which did not lead to a megathrust earthquake.

Finally, we conducted a numerical simulation of megathrust earthquake cycles (seismic supercycles³⁰) to examine the possibility of a very long-term transient event in the JTSZ. We used the two-dimensional mechanical model³¹ in Supplementary Fig. 2 and the method described in Methods. The velocity of the Pacific plate motion relative to the North American plate (V_{pl}) is assumed to be 8.5 cm per year off central Tohoku, according to a plate kinematic model³². As in the original simulation³¹, the result of the simulation in Supplementary Fig. 3 indicates that a megathrust earthquake (large step) occurs every several hundred years, with large slips in the shallower regions around the Tohoku earthquake hypocentre (A, B and C) and smaller slips in the deeper region (D). Under such conditions, slip deficit accumulates (represented by long slopes) and M 7-class earthquakes (small steps in the slopes) occur during a supercycle between megathrust earthquakes. Moreover, several very long-term transient events (flat parts in the slopes) also occur in D during a supercycle, with the last one occurring just before and leading to a megathrust earthquake.

In short, the interseismic period of a supercycle should consist primarily of three phenomena: the slip deficit of the regular process, M 7-class earthquakes and very long-term transient events. Therefore, the 2002 to March 2011 parts of the time series shown in Fig. 1b should represent the sum of the effects of these three phenomena. However, because the crustal and in-slab earthquakes (dark green stars in Fig. 1a) do not belong to the category of M 7-class earthquake defined above, we removed their co-seismic and post-seismic signals (dark green lines in Fig. 1b) from the time series. We measured composite deformations due to the three phenomena from the revised time series, as shown with pink arrows in Supplementary Fig. 4. Then, we performed an inversion of the composite deformations for a source model extending along the Japan Trench, and computed ΔCFF values due to the resultant slip distribution (purple contours in Supplementary Fig. 4), finding the composite ΔCFF around the mainshock hypocentre to be $\sim +1$ MPa.

Considering the results of both the seismic supercycle simulation and the two ΔCFF computations, it is proven that a very long-term transient event occurred over the 9 years preceding the 2011 Tohoku earthquake. We also suggest that this transient event could have led to the advancement of the occurrence time³³ of the Tohoku earthquake, contributing 10% to the composite ΔCFF . The above simulation did not perfectly reproduce these findings; performing an extensive set of simulations with new inputs would provide further understanding of the physics governing such transient events and their relationship to a megathrust earthquake like Tohoku.

Methods

Interseismic deformation. The GSI is analysing GPS data from their GEONET using the Bernese GPS software (version 5.0), IGS ephemerides, and ITRF2005 (ref. 34). The results of the analyses are being published as the F3 solutions for the daily coordinates of the GEONET stations³⁴. The F3 solutions for stations in the Tohoku and Kanto districts from 21 March 1996 to 8 March 2011 were adjusted by removing the motion of the North American plate in the plate kinematic model³². The adjusted coordinates were transformed into the Japan plane rectangular

coordinate system VIII, IX or X (ref. 35). The horizontal components of deformation at each station were calculated from the obtained x - and y -coordinates.

Source inversion scheme. We performed the source inversion using a least-squares method with constraints³⁶, although the positivity constraint on slip vectors was not applied because the observed deviations include the effects of the very long-term transient event and the northern backslip source; thus, we had to model reverse- and normal-faulting sources simultaneously. Smoothness constraints with digital Laplacians remained, and their weights were determined by minimizing Akaike's Bayesian Information Criterion³⁷. Green's functions were computed using the average one-dimensional velocity structure in and off the Tohoku and Kanto districts and the propagator matrix method with dynamic-static separation³⁸.

Seismic supercycle simulation. The simulation was conducted using a two-dimensional mechanical model³¹ and the method for media under the rate- and state-dependent friction law³⁹. The plate boundary shape in the original model was revised to fit to the actual shape in and off central Tohoku. We also modified the distributions of parameters in the rate- and state-dependent friction law ($a - b$, L and σ_n^{eff}) (ref. 39) along the plate boundary. The unstable zone of $a - b \sim 0$ was extended to a depth over 60 km and the effective normal stress σ_n^{eff} was decreased at depths deeper than 40 km (Supplementary Fig. 2). Other minor changes in $a - b$, σ_n^{eff} and L were also applied as shown in Supplementary Fig. 2.

References

- Ozawa, S. *et al.* Detection and monitoring of ongoing aseismic slip in the Tokai region, central Japan. *Science* **298**, 1009–1012 (2002).
- Kostoglodov, V. *et al.* A large silent earthquake in the Guerrero seismic gap, Mexico. *Geophys. Res. Lett.* **30**, 1807 (2003).
- Ide, S., Beroza, G. C., Shelly, D. R. & Uchide, T. A scaling law for slow earthquakes. *Nature* **447**, 76–79 (2007).
- Hirose, H. *et al.* Slow earthquakes linked along dip in the Nankai subduction zone. *Science* **330**, 1502 (2010).
- Roeloffs, E. A. Evidence for aseismic deformation rate changes prior to earthquakes. *Annu. Rev. Earth Planet. Sci.* **34**, 591–627 (2006).
- Iio, Y., Kobayashi, Y. & Tada, T. Large earthquakes initiate by the acceleration of slips on the downward extensions of seismogenic faults. *Earth Planet. Sci. Lett.* **202**, 337–343 (2002).
- Graham, S. E. *et al.* GPS constraints on the 2011–2012 Oaxaca slow slip event that preceded the 2012 March 20 Ometepe earthquake, southern Mexico. *Geophys. J. Int.* **197**, 1593–1607 (2014).
- Yokota, Y. *et al.* Joint inversion of strong motion, teleseismic, geodetic, and tsunami datasets for the rupture process of the 2011 Tohoku earthquake. *Geophys. Res. Lett.* **38**, L00G21 (2011).
- Ozawa, S. *et al.* Preceding, coseismic, and postseismic slips of the 2011 Tohoku earthquake, Japan. *J. Geophys. Res.* **117**, B07404 (2012).
- Ozawa, S., Suito, H., Nishimura, T., Tobita, M. & Munekane, H. Possibility of recovery of slip deficit rate between the North American plate and Pacific plate off Sanriku, northeast Japan. *Geophys. Res. Lett.* **34**, L20308 (2007).
- Kato, A. *et al.* Propagation of slow slip leading up to the 2011 M_w 9.0 Tohoku-Oki earthquake. *Science* **335**, 705–708 (2012).
- Ito, Y. *et al.* Episodic slow slip events in the Japan subduction zone before the 2011 Tohoku-Oki earthquake. *Tectonophysics* **600**, 14–26 (2013).
- Nishimura, T. *et al.* Temporal change of interplate coupling in northeastern Japan during 1995–2002 estimated from continuous GPS observations. *Geophys. J. Int.* **157**, 901–916 (2004).
- Suwa, Y., Miura, S., Hasegawa, A., Sato, T. & Tachibana, K. Interplate coupling beneath NE Japan inferred from three-dimensional displacement field. *J. Geophys. Res.* **111**, B04402 (2006).
- Hashimoto, C., Noda, A., Sagiya, T. & Matsu'ura, M. Interplate seismogenic zones along the Kuril-Japan trench inferred from GPS data inversion. *Nat. Geosci.* **2**, 141–144 (2009).
- Loveless, J. P. & Meade, B. J. Geodetic imaging of plate motions, slip rates, and partitioning of deformation in Japan. *J. Geophys. Res.* **115**, B02410 (2010).
- Ozawa, S., Suito, H., Imakiire, T. & Murakami, M. Spatiotemporal evolution of aseismic interplate slip between 1996 and 1998 and between 2002 and 2004, in Bungo channel, southwest Japan. *J. Geophys. Res.* **112**, B05409 (2007).
- Mavrommatis, A. P., Segall, P. & Johnson, K. M. A decadal-scale deformation transient prior to the 2011 M_w 9.0 Tohoku-oki earthquake. *Geophys. Res. Lett.* **41**, 4486–4494 (2014).
- Suito, H., Nishimura, T., Tobita, M., Imakiire, T. & Ozawa, S. Interplate fault slip along the Japan Trench before the occurrence of the 2011 off the Pacific coast of Tohoku Earthquake as inferred from GPS data. *Earth Planets Space* **63**, 615–619 (2011).
- Nishimura, T. Crustal deformation of northeastern Japan based on geodetic data for recent 120 years [in Japanese]. *J. Geol. Soc. Jpn* **118**, 278–293 (2012).
- Heki, K. & Mitsui, Y. Accelerated Pacific plate subduction following interplate thrust earthquakes at the Japan trench. *Earth Planet. Sci. Lett.* **363**, 44–49 (2013).
- Fu, Y. & Freymueller, J. T. Repeated large slow slip events at the southcentral Alaska subduction zone. *Earth Planet. Sci. Lett.* **375**, 303–311 (2013).
- Liu, Y. & Rice, J. R. Aseismic slip transients emerge spontaneously in three-dimensional rate and state modeling of subduction earthquake sequences. *J. Geophys. Res.* **110**, B08307 (2005).
- Matsuzawa, T., Hirose, H., Shibasaki, B. & Obara, K. Modeling short- and long-term slow slip events in the seismic cycles of large subduction earthquakes. *J. Geophys. Res.* **115**, B12301 (2010).
- Savage, J. C. A dislocation model of strain accumulation and release at a subduction zone. *J. Geophys. Res.* **88**, 4984–4996 (1983).
- Savage, J. C. Interseismic uplift at the Nankai subduction zone, southwest Japan. *J. Geophys. Res.* **100**, 6339–6350 (1995).
- Meade, B. J. & Loveless, J. P. Predicting the geodetic signature of $M_w \geq 8$ slow slip events. *Geophys. Res. Lett.* **36**, L01306 (2009).
- Reasenber, P. A. & Simpson, R. W. Response of regional seismicity to the static stress change produced by the Loma Prieta earthquake. *Science* **255**, 1687–1690 (1992).
- Liu, Y. & Rice, J. R. Spontaneous and triggered aseismic deformation transients in a subduction fault model. *J. Geophys. Res.* **112**, B09404 (2007).
- Sieh, K. *et al.* Earthquake supercycles inferred from sea-level changes recorded in the corals of west Sumatra. *Science* **322**, 1674–1678 (2008).
- Kato, N. & Yoshida, S. A shallow strong patch model for the 2011 great Tohoku-oki earthquake: a numerical simulation. *Geophys. Res. Lett.* **38**, L00G04 (2011).
- Drewes, H. in *Geodetic Reference Frames 95–99* (IAG Symposia, 2009).
- Stein, R. S., Barka, A. A. & Dieterich, J. H. Progressive failure on the North Anatolian fault since 1939 by earthquake stress triggering. *Geophys. J. Int.* **128**, 594–604 (1997).
- Nakagawa, H. *et al.* Development and validation of GEONET new analysis strategy (Version 4) [in Japanese]. *J. Geograph. Surv. Inst.* **118**, 1–8 (2009).
- Geospatial Information Authority. Introduction to the Japan plane rectangular coordinate system [in Japanese]. Available at <https://vldb.gsi.go.jp/sokuchi/patchjgd/download/Help/jpc/jpc.htm> (2010).
- Yoshida, S. *et al.* Joint inversion of near- and far-field waveforms and geodetic data for the rupture process of the 1995 Kobe earthquake. *J. Phys. Earth* **44**, 437–454 (1996).
- Akaike, H. in *Bayesian Statistics*. (eds Bernardo, J. M., DeGroot, M. H., Lindley, D. V. & Smith, M.) 143–166 (Valencia Univ. Press, 1980).
- Zhu, L. & Rivera, L. A. A note on the dynamic and static displacements from a point source in multilayered media. *Geophys. J. Int.* **148**, 619–627 (2002).
- Kato, N. & Tullis, T. E. Numerical simulation of seismic cycles with a composite rate- and state-dependent friction law. *Bull. Seismol. Soc. Am.* **93**, 841–853 (2003).

Acknowledgements

We are grateful to Naoyuki Kato, Kazushige Obara, Paul Segall and Teruyuki Kato for many discussions and advice. We also appreciate colleagues at the GSI for the GEONET F3 solutions and discussions. Shun Higuchi and Hiroaki Kobayashi helped us with additional computation. All figures were drawn using the Generic Mapping Tools. The co-seismic distribution in Figures 2 and 3 was reproduced with permission from Yokota *et al.*⁸ This study was supported by the Grants-in-Aid for Scientific Research 23900002 and 10J06987 from MEXT.

Author contributions

Both the authors equally contributed to this paper.

Additional information

Supplementary Information accompanies this paper at <http://www.nature.com/naturecommunications>

Competing financial interests: The authors declare no competing financial interests.

Reprints and permission information is available online at <http://npg.nature.com/reprintsandpermissions/>

How to cite this article: Yokota, Y. and Koketsu, K. A very long-term transient event preceding the 2011 Tohoku earthquake. *Nat. Commun.* **6**:5934 doi: 10.1038/ncomms6934 (2015).