ARTICLE

Received 14 Jan 2015 | Accepted 11 Mar 2015 | Published 23 Apr 2015

DOI: 10.1038/ncomms7903

OPEN

Direct synthesis of imino-C-nucleoside analogues and other biologically active iminosugars

Milan Bergeron-Brlek¹, Michael Meanwell¹ & Robert Britton¹

Iminosugars have attracted increasing attention as chemical probes, chaperones and leads for drug discovery. Despite several clinical successes, their *de novo* synthesis remains a significant challenge that also limits their integration with modern high-throughput screening technologies. Herein, we describe a unique synthetic strategy that converts a wide range of acetaldehyde derivatives into iminosugars and imino-C-nucleoside analogues in two or three straightforward transformations. We also show that this strategy can be readily applied to the rapid production of indolizidine and pyrrolizidine iminosugars. The high levels of enantio- and diastereoselectivity, excellent overall yields, convenience and broad substrate scope make this an appealing process for diversity-oriented synthesis, and should enable drug discovery efforts.

¹Department of Chemistry, Simon Fraser University, Burnaby, British Columbia, Canada V7G 1S2. Correspondence and requests for materials should be addressed to R.B. (email: rbritton@sfu.ca).

Iminosugars are naturally occurring carbohydrate mimics that inhibit many enzymes of medicinal interest¹. Their biological activity is often attributed to a structural resemblance to the oxacarbenium ion-like transition states that occur during the enzymatic hydrolysis of carbohydrates². As such, many iminosugars are potent inhibitors of glycosidases and glycosyltransferases¹, and have been highlighted as lead candidates for the treatment of a variety of diseases, including cancer, diabetes, viral infections and lysosomal storage disorders (for example, Gaucher and Fabry disease)^{1,3}. The most common naturally occurring iminosugars possess a polyhydroxylated pyrrolidine core and may be additionally annulated as in the pyrrolizidines (for example, **2**, Fig. 1), indolizidines (for example, **3**), or nortropanes⁴. A growing number of unnatural analogues of these compounds have also been reported as leads for drug discovery, including the imino-C-nucleosides developed by Schramm (for example, **1**)^{5,6} and β -hexosaminidase inhibitors developed by Wong^{7,8}. Unfortunately, the incorporation of pyrrolidine iminosugars into chemical screening libraries or diversity-oriented synthesis (DOS) campaigns is problematic, as their syntheses are often lengthy, low-yielding, cost-intensive and limited by reliance on carbohydrate building blocks⁹. Thus, while several such pyrrolidine iminosugars have emerged as clinical candidates or drugs¹, fundamental tools for their high-throughput synthesis are lacking. In fact, much of the success in imino-C-nucleoside synthesis^{10–14} (for example, **1** (ref. 6) and **4** (ref. 5)) has relied on the common building block **5** (refs 14–16). As evidenced by step counts provided in Fig. 1, the synthesis of pyrrolizidine- and indolizidine-based iminosugars (for example, **2**; (ref. 17) and **3**; (ref. 18)) also remains a significant synthetic challenge.

We have reported preliminarily that when mixtures of the dioxanone **8**, an aliphatic aldehyde **6** and *N*-chlorosuccinimide (NCS) are treated with (*S*)-proline, a series of well-orchestrated reactions occur¹⁹. First, the aldehyde undergoes α -chlorination²⁰, producing a racemic mixture of α -chloroaldehydes **7**. Second, an enantioselective proline-catalysed aldol reaction occurs between the dioxanone **8** and the α -chloroaldehyde (*R*)-**7**. Importantly, proline also catalyses racemization of the α -chloroaldehydes **7** and, consequently, this second step effects a dynamic kinetic resolution (DKR)¹⁹. Thus, this one-pot reaction transforms commodity chemicals **6** and **8** into carbohydrate

Figure 1 | Biologically active iminosugars and the common building block **5 for imino-C-nucleoside synthesis.** Immucillin-H (**1**) is a potent transition-state analogue inhibitor of purine nucleoside phosphorylase and a lead for the treatment of human T-cell leukaemia and lymphoma. The structurally related imino-C-nucleoside analogue **4** inhibits nucleoside hydrolase. In addition, the pyrrolizidine and indolizidine iminosugars **2** and **3** inhibit α -mannosidase, an enzyme target for anticancer therapies. The number of steps required to synthesize each of **1–4** and the synthetic building block **5** highlight the challenges faced when incorporating pyrrolidine iminosugars into chemical screening libraries and medicinal chemistry campaigns.

building blocks **9** in excellent yield, diastereoselectivity and enantioselectivity. Considering the spatial relationship between the chloromethine and carbonyl functions in **9**, these aldol adducts may also serve as building blocks^{21–24} for the synthesis of polyhydroxypyrrolidines via a reductive amination–annulation sequence (see grey box, Fig. 2). Such a strategy would allow for the conversion of virtually any acetaldehyde derivative **6** into an iminosugar **10** in two straightforward transformations from commodity chemicals, thus enabling their integration with modern high-throughput screening technologies.

Here we demonstrate that the reductive amination of a wide range of ketochlorohydrins **9** provides a rapid route to pyrrolidine iminosugars^{8,11,25–32}, such as those depicted in Fig. 1. Importantly, this unique two- or three-step process requires no cryogenic, anhydrous or otherwise complicated experimental conditions. The demonstration of this strategy in several short syntheses of biologically active imino-C-nucleoside analogues, and indolizidine and pyrrolizidine iminosugars highlights its adaptability for DOS and the rapid preparation of iminosugar-based screening libraries²⁶.

Results

Reductive amination of α -chlorination-DKR aldol products.

The utility of the synthetic strategy outlined in Fig. 2 relies intimately on a diastereoselective reductive amination of aldol adducts **9**. Enders has reported²⁷ that the reductive amination of related aldol adducts that lack a chloromethine function were non selective (*dr* < 2:1) using NaB(OAc)₃H. Likewise, Madsen found similar selectivities in the reductive amination of the corresponding *syn*-aldol adduct³³. Bearing this in mind, we began by screening solvents and reducing agents, as well as the addition of acetic acid to the reductive amination of ketochlorohydrin **11** (Table 1) (ref. 19). In all cases, an excess of amine was required for complete imine formation and avoidance of competing ketone reduction (entry 1). As indicated in entry 2, the conditions reported by Enders²⁷ delivered the amino alcohols **12a** and **12b** in good yield (82%), albeit low diastereoselectivity. The relative stereochemistry of **12a** was assigned based on analysis of ³J_{H,H} coupling constants and NOESY spectra recorded on the cyclic carbamate derived from the reaction of **12a** with carbonyldiimidazole. Use of NaB(CN)H₃ resulted in an improved diastereomeric ratio of these products (*dr* ~ 6:1) in both CH₂Cl₂ and MeCN (entries 3 and 4) and in tetrahydrofuran (THF) the 1,3-*syn* amino alcohol **12a** was produced as the only detectable diastereomer in near quantitative yield (entry 5). As

Figure 2 | A convenient synthesis of polyhydroxypyrrolidine iminosugars. Organocatalytic tandem α -chlorination-DKR aldol reaction coupled with a reductive amination/annulation sequence to access iminosugars **10**.

Table 1 | Reductive amination of ketochlorohydrin **11.**

Entry	Reducing agent	Solvent	R	Products (% yield)*
1	NaBH ₄ [†]	CH ₂ Cl ₂	Bn	12a (17), 12b (28) [‡]
2	NaB(OAc) ₃ H ^{†,§}	CH ₂ Cl ₂	Bn	12a (36), 12b (46)
3	NaB(CN)H ₃ [§]	CH ₂ Cl ₂	Bn	12a (85), 12b (13)
4	NaB(CN)H ₃ [§]	MeCN	Bn	12a (83), 12b (15)
5	NaB(CN)H ₃ [§]	THF	Bn	12a (98), 12b (<2)
6	NaB(CN)H ₃ [§]	THF	allyl	13a (85), 13b (<2)
7	NaB(CN)H ₃ [§]	THF	propargyl	14a (83), 14b (<2)

a) 2.5 Equivalents of amine is added to stirred solution of **11** at room temperature, after 1 h reducing agent is added and mixture is stirred for an additional 1 h.

*Isolated yield.

[†]Reaction stirred for 12 h.

[‡]Accompanied by significant amounts of carbonyl reduction products.

[§]1 equiv. of HOAc added.

^{||}Not detected in ¹H-NMR spectra recorded on crude reaction product.

summarized in entries 6 and 7, this optimized protocol proved general and also provided access to the corresponding *N*-allyl and *N*-propargyl amines **13a** and **14a**, in excellent yield and diastereoselectivity.

Synthesis of pyrrolidine iminosugars. While the amino-chlorohydrin **12a** did not cyclize directly, its high-yielding conversion into the pyrrolidine iminosugar **15** simply required heating in methanol, which also promoted acetonide removal (Fig. 3). Alternatively, this cyclization could be effected by heating **12a** in toluene with excess NaHCO₃, which provided the orthogonally protected iminosugar **16**. Anticipating that the increased reactivity of a benzylchlorohydrin would favour a one-pot reductive amination–annulation process, the readily available aldol adduct **17** (ref. 19) was also treated with NaB(CN)H₃ in a mixture of THF/HOAc. Following this optimized procedure, the orthogonally protected iminosugar **18** was produced directly and in excellent yield. Removal of both the acetonide and benzyl-protecting groups by hydrogenolysis in acidic methanol gave the imino-C-nucleoside analogue **4**. Considering the aldol adduct **17** is available in one step from phenyl acetaldehyde¹⁹, this three-step synthesis of **4**, a potent (*K*_i = 170 nM) transition-state analogue inhibitor of nucleoside hydrolase⁵, represents a significant advance.

Scope of direct iminosugar synthesis. To further evaluate the scope of this direct iminosugar synthesis, we repeated the reactions described in Fig. 3 with several additional alkyl- and aryl-substituted ketochlorohydrins prepared in one step using our (*S*)-proline-catalysed α -chlorination-DKR aldol reaction¹⁹. It is noteworthy that the enantiomeric ketochlorohydrins are also readily prepared using the corresponding (*R*)-proline-catalysed reaction¹⁹. As indicated in Fig. 4, the reductive amination–annulation process is general and delivers a wide range of polyhydroxypyrrolidines **22–43** in good to excellent overall yield. A number of the orthogonally protected iminosugars depicted in Fig. 4 are crystalline and their structures were confirmed by X-ray crystallographic analysis (Supplementary Information). As expected (*vide supra*), the synthesis of alkyl-substituted iminosugars required an additional cyclization step, whereby

Figure 3 | Synthesis of pyrrolidine iminosugars and imino-C-nucleoside analogues.

A highly diastereoselective reductive amination of chlorohydrin aldol adducts followed by brief heating in methanol or toluene with NaHCO₃ provides rapid access to native or differentially protected iminosugars. Reductive amination of the benzyl chloride-containing aldol adduct **17** leads directly to the protected iminosugar **18**, a precursor to the potent nucleoside hydrolase inhibitor **4**.

the product of reductive amination was heated in MeOH or toluene with NaHCO₃. Thus, this convenient process can be tailored for the production of orthogonally protected (**22**, **24**, **26**, **28** and **30**) or native iminosugars (**23**, **25**, **27**, **29** and **31**). Conversely, reductive amination of aryl-substituted chlorohydrins provided the corresponding iminosugars **18** and **36–39** directly and in excellent yield. It is notable that both the one- and two-step iminosugar syntheses proved tolerant of various functional groups. For example, electron deficient aryl (**36** and **37**), electron rich aryl (**38**) and heteroaryl substituents (**39**) were readily incorporated. Likewise, alkyl (**22–35**), branched alkyl (**24**), silyloxy alkyl (**32**), allyl (**26**), propargyl (**28**), primary alkyl chloride (**33**) and benzyl (**30**) groups were all compatible with the reaction sequence. The use of benzyl amine (**22–39**), allyl amine (**40–41**) or propargyl amine (**42–43**) also highlights the utility of this process for DOS and the potential for further elaboration of these iminosugars through metathesis, click, or cross coupling reactions. Finally, the mild reaction conditions and the protecting-group compatibility deserves note, as the 1,3-dioxane function in **34** and **35**, silyl protecting group in **32** and **43**, and acetonide function (conditions a)) in all substrates remained intact throughout the reaction sequence. Importantly, these readily available iminosugars share many features considered optimal for lead identification^{34–37}, including MW < 350 DA, CLogP < 2, multiple chiral centres, heterocyclic rings and H-bond donors/acceptors. In addition, the ease with which the amine function and ring substituent (*R*¹ and *R*² in **20**) can be differentiated provides unique opportunities for further diversification.

Short syntheses of polyhydroxy pyrrolizidines and indolizidines.

Figure 5 highlights the further application of this convenient strategy to the rapid preparation of several structurally complex polyhydroxy indolizidine and pyrrolizidine alkaloids, including analogues of the glycosidase inhibitors hyacinthacine and steviamine. While several strategies could be exploited for

Figure 4 | Scope of iminocyclitol synthesis. (a) amine (2.5 equivalents), AcOH, 4-Å mol sieves, THF; then NaB(CN)H₃, room temperature; (b) NaHCO₃, PhMe, 105 °C; (c) MeOH, 120 °C, (microwave reactor).

the second annulation event, ready access to the *N*-allyl pyrrolidine **40**, alkyl chloride **33** and protected ketones **34** and **35** suggested annulation events involving ring closing metathesis^{38,39}, alkylation^{24,40,41} or reductive amination^{17,42}. For example, heating the dienylpyrrolidine **40** with the Hoveyda-Grubbs 2nd generation catalyst⁴³ in toluene provided the unsaturated indolizidine **45** in excellent overall yield from 4-pentenal (**44**). Alternatively, starting with 6-chloropentanal (**46**) or 5-chloropentanal (**48**), α -chlorination-DKR aldol reactions¹⁹ followed by reductive amination and cyclization provided the chloroalkylpyrrolidine **33** and pyrrolizidine **49**, respectively. Conversion of **33** into the corresponding indolizidine **47** (ref. 44) required hydrolytic removal of the benzyl-

Figure 5 | Total syntheses of indolizidine and pyrrolizidine iminocyclitols.

(a) NCS, dioxanone **8**, (*S*)-proline (80 mol%), CH₂Cl₂, room temperature (RT); (b) benzyl amine or allyl amine (2.5 equivalents), AcOH, 4-Å mol sieves, THF; then NaB(CN)H₃, RT; (c) NaHCO₃, PhMe, 105 °C; (d) Hoveyda-Grubbs cat (2nd generation, 5 mol%), PhMe, 60 °C, 2 h; (e) H₂ (90 bar), MeOH, 60 °C (H-Cube); (f) NaHCO₃, MeOH, 80 °C, 16 h; then PPTS; (g) PPTS (cat), H₂O, MeOH, 100 °C, 0.5 h (microwave reactor). NCS, *N*-chlorosuccinimide, PPTS, pyridinium *p*-toluenesulfonate.

protecting group and brief treatment with base. Completion of the total synthesis of 7a-*epi*-hyacinthacine A₁ (**50**)^{45,46} simply involved hydrogenolysis of **49** in acidic media.

The reductive amination strategy was explored in short syntheses of the hyacinthacine and steviamine analogues **2** (ref. 17) and *ent*-**3** (ref. 18). In both cases, the ketone function in the readily available pyrrolidines **34** and **35** was unveiled in concert with hydrogenolytic cleavage of the *N*-benzyl group, and the resulting iminium species (not shown) was reduced *in situ* to afford the products depicted as single diastereomers. Importantly, each of the total syntheses depicted in Fig. 5 requires 5 steps or less, originates with inexpensive and readily available chemicals,

and is completed in a matter of days, which compares well with the reported syntheses of these and related compounds (see for example, Fig. 1).

Discussion

In summary, a highly convergent synthesis of iminosugars has been developed that converts a wide range of acetaldehyde derivatives into polyhydroxypyrrolidines in two or three straightforward reactions and does not rely on carbohydrate building blocks. The application of this cost-effective process to the rapid synthesis of indolizidine and pyrrolizidine iminosugars also highlights its utility for the preparation of more structurally complex natural products and their analogues. Importantly, the excellent overall yields, diastereoselectivity and enantioselectivity, coupled with tunability of pharmacophoric features make this process well suited for chemical screening library and DOS campaigns.

Methods

Representative example of reductive amination/annulation sequence.

Synthesis of aminochlorohydrin **18** and iminocyclitol **4**. To a stirred solution of **17** (ref. 19; 130 mg, 0.457 mmol) in THF (4.55 ml) was added BnNH_2 (125 μl , 1.15 mmol) and glacial acetic acid (27.0 μl , 0.457 mmol), and the resulting mixture was stirred at 20 °C for 1 h. $\text{NaB}(\text{CN})\text{H}_3$ (72 mg, 1.15 mmol) was then added and the mixture was stirred for one additional hour. The reaction mixture was then diluted with CH_2Cl_2 to a concentration of 0.05 M and treated with water. The layers were separated and the organic layer was washed with brine, dried (MgSO_4) and concentrated under reduced pressure. Purification of the crude product by flash chromatography (pentane-EtOAc 8:2) afforded pyrrolidine **18** (126 mg, 81% yield) as a crystalline solid. mp = 108–111 °C (EtOH); R_f (pentane-EtOAc 6:4) 0.81; $[\alpha]_D^{20} = +11$ (c 0.70 in CHCl_3); infrared (neat): $\nu = 3444, 2988, 2874, 1454, 1381, 1210, 1048, 853, 753$ and 700 cm^{-1} ; ^1H -NMR (600 MHz, CDCl_3): $\delta = 7.53$ (d, $J = 7.4 \text{ Hz}$, 2H), 7.38 (t, $J = 7.5 \text{ Hz}$, 2H), 7.31–7.20 (m, 6H), 4.03 (d, $J = 4.4 \text{ Hz}$, 1H), 3.90 (d, $J = 12.9 \text{ Hz}$, 1H), 3.74 (s, 1H), 3.74 (dd, $J = 4.5 \text{ Hz}$, $J = 9.6 \text{ Hz}$, 1H), 3.47 (d, $J = 12.8 \text{ Hz}$, 1H), 3.46 (dd, $J = 10.5 \text{ Hz}$, 1H), 3.25 (dd, $J = 4.1 \text{ Hz}$, $J = 10.5 \text{ Hz}$, 1H), 2.89 (ddd, $J = 4.1 \text{ Hz}$, $J = 10.5 \text{ Hz}$, 1H), 2.31 (s, 1H), 1.42 (s, 3H) and 1.40 p.p.m. (s, 3H); ^{13}C -NMR (151 MHz, CDCl_3): $\delta = 141.3, 139.2, 128.9, 128.5, 128.2, 127.4, 127.4, 127.3, 100.3, 76.9, 76.8, 74.1, 67.2, 59.6, 58.7, 29.2$ and 19.8 p.p.m. ; HRMS ESI (high-resolution mass spectrometry electrospray ionization) m/z calcd (calculated) for $\text{C}_{21}\text{H}_{26}\text{NO}_3$ $[\text{M} + \text{H}]^+$ 340.1907, found 340.1886.

Preparation of the imino-C-nucleoside analogue 4. A solution of **18** (20 mg, 0.059 mmol) and pyridinium *p*-toluenesulfonate (15 mg, 0.059 mmol) in 1:1 $\text{H}_2\text{O}/\text{MeOH}$ (4.0 ml) was added to a microwave vial. The vial was sealed in a CEM Discover LabMate microwave reactor and the resulting mixture was heated at 100 °C (as monitored by a vertically focused infrared temperature sensor) for 30 min. The resulting solution was concentrated under reduced pressure and the crude product was used in the next reaction without further purification. A solution of the crude iminocyclitol *p*-toluenesulfonate salt in MeOH (20 ml) was passed twice through an H-Cube continuous-flow reactor using a 30 mm 10% Pd/C cartridge. Conditions: temperature = 35 °C; flow rate = 0.8 ml min^{-1} ; H_2 pressure = 40 bar. The resulting mixture was stirred with DOWEX 1X8-100 (HO^- form) for a further 30 min and the resin was removed by filtration. Concentration and purification of the crude product by flash chromatography on C_{18} silica gel (H_2O) afforded iminoribitol **4** (10 mg, 83% yield over 2 steps) as a colourless oil. $[\alpha]_D^{20} = -31$ (c 0.48 in MeOH); infrared (neat): $\nu = 3306, 2918, 1560, 1494, 1454, 1406, 1347, 1081, 951, 757$ and 699 cm^{-1} ; ^1H -NMR (400 MHz, CD_3OD): $\delta = 7.45\text{--}7.42$ (m, 2H), 7.37–7.32 (m, 2H), 7.27 (ddt, $J = 1.4 \text{ Hz}$, $J = 6.4 \text{ Hz}$, $J = 8.5 \text{ Hz}$, 1H), 4.01 (d, $J = 7.2 \text{ Hz}$, 1H), 3.97 (dd, $J = 4.7 \text{ Hz}$, $J = 6.0 \text{ Hz}$, 1H), 3.86 (dd, $J = 6.1 \text{ Hz}$, $J = 7.2 \text{ Hz}$, 1H), 3.74 (d, $J = 4.5 \text{ Hz}$, 2H) and 3.15 p.p.m. (q, $J = 4.5 \text{ Hz}$, 1H); ^{13}C -NMR (151 MHz, CD_3OD): $\delta = 142.6, 129.5, 128.5, 128.1, 79.2, 73.6, 68.1, 66.8$ and 63.2 p.p.m. ; HRMS (ESI) m/z calcd for $\text{C}_{11}\text{H}_{15}\text{NO}_3$ $[\text{M} + \text{H}]^+$ 210.1125, found 210.1111.

References

- Compain, P. & Martin, O. *Iminosugars from Synthesis to Therapeutic Applications* (Wiley, 2007).
- Lillelund, V. H., Jensen, H. H., Liang, X. & Bols, M. Recent developments of transition-state analogue glycosidase inhibitors of non-natural product origin. *Chem. Rev.* **102**, 515–553 (2002).
- Asano, N. Naturally occurring iminosugars and related compounds: structure, distribution, and Biological activity. *Curr. Top. Med. Chem.* **3**, 471–484 (2003).
- Asano, N. in *Glycoscience*. (eds Fraser-Reid, B., Tatsuta, K. & Thiem, J.) 1887–1911 (Springer-Verlag, 2008).
- Horenstein, B. A., Zabinski, R. F. & Schramm, V. L. A new class of C-nucleosides. 1-(S)-aryl-1,4-dideoxy-1,4-imino-D-ribitol, transition state analogue inhibitors of nucleoside hydrolase. *Tetrahedron Lett.* **34**, 7213–7216 (1993).
- Kicska, G. A. *et al.* Immucillin H, a powerful transition-state analog inhibitor of purine nucleoside phosphorylase, selectively inhibits human T lymphocytes. *Proc. Natl Acad. Sci. USA* **98**, 4593–4598 (2001).
- Liu, J. S., Shikhman, A. R., Lotz, M. K. & Wong, C.-H. Hexosaminidase inhibitors as new drug candidates for the therapy of osteoarthritis. *Chem. Biol.* **8**, 701–711 (2001).
- Liang, P.-H. *et al.* Novel five-membered iminocyclitol derivatives as selective and potent glycosidase inhibitors: new structures for antivirals and osteoarthritis. *ChemBioChem* **7**, 165–173 (2006).
- Hong, Z., Liu, L., Sugiyama, M., Fu, Y. & Wong, C.-H. Concise synthesis of iminocyclitols via petasis-type aminocyclization. *J. Am. Chem. Soc.* **131**, 8352–8353 (2009).
- Compain, P., Chagnault, V. & Martin, O. R. Tactics and strategies for the synthesis of iminosugar C-glycosides: a review. *Tetrahedron: Asymm.* **20**, 672–711 (2009).
- Stocker, B. L., Dangerfield, E. M., Win-Mason, A. L., Haslett, G. W. & Timmer, M. S. M. Recent developments in the synthesis of pyrrolidine-containing iminosugars. *Eur. J. Org. Chem.* **9**, 1615–1637 (2010).
- Yokoyama, M. & Momotake, A. Synthesis and biological activity of azanucleosides. *Synthesis* **9**, 1541–1554 (1999).
- Merino, P., Tejero, T. & Delso, I. Current developments in the synthesis and biological activity of aza-C-nucleosides: immucillins and related compounds. *Curr. Med. Chem.* **15**, 954–967 (2008).
- Evans, G. B., Furneaux, R. H., Hausler, H., Larsen, J. S. & Tyler, P. C. Imino-C-nucleoside synthesis: heteroaryl lithium carbanion additions to a carbohydrate cyclic imine and nitron. *J. Org. Chem.* **69**, 2217–2220 (2004).
- Evans, G. B., Furneaux, R. H., Gainsford, G. J., Schramm, V. L. & Tyler, P. C. Synthesis of transition state analogue inhibitors for purine nucleoside phosphorylase and *N*-riboside hydrolases. *Tetrahedron* **56**, 3053–3062 (2000).
- Evans, G. B. *et al.* Addition of lithiated 9-deazapurine derivatives to a carbohydrate cyclic imine: convergent synthesis of the aza-C-nucleoside immucillins. *J. Org. Chem.* **66**, 5723–5730 (2001).
- Izquierdo, I., Plaza, M. T., Tamayo, J. A., Rodríguez, M. & Martos, A. Polyhydroxylated pyrrolizidines. Part 8: enantiospecific synthesis of looking-glass analogues of hyacinthacin A_5 from DADP. *Tetrahedron* **62**, 6006–6011 (2006).
- Ansari, A. A. & Vankar, Y. D. Synthesis of dihydroxymethyl dihydroxypyrrolidines and steviamine analogues from C-2 formyl glycals. *J. Org. Chem.* **78**, 9383–9395 (2013).
- Bergeron-Breke, M., Teoh, T. & Britton, R. A tandem organocatalytic α -chlorination-aldol reaction that proceeds with dynamic kinetic resolution: a powerful tool for carbohydrate synthesis. *Org. Lett.* **15**, 3554–3557 (2013).
- Halland, N., Braunton, A., Bachmann, S., Marigo, M. & Jørgensen, K. A. Direct organocatalytic asymmetric α -chlorination of aldehydes. *J. Am. Chem. Soc.* **126**, 4790–4791 (2004).
- Britton, R. & Kang, B. α -Haloaldehydes: versatile building blocks for natural product synthesis. *Nat. Prod. Rep.* **30**, 227–236 (2013).
- Draper, J. & Britton, R. A concise and stereoselective synthesis of hydroxypyrrolidines: rapid synthesis of (+)-preussin. *Org. Lett.* **12**, 4034–4037 (2010).
- Dhand, V., Draper, J. A., Moore, J. & Britton, R. A short, organocatalytic formal synthesis of (–)-swainsonine and related alkaloids. *Org. Lett.* **15**, 1914–1917 (2013).
- Dhand, V., Chang, S. & Britton, R. Total synthesis of the cytotoxic anhydrophytosphingosine pachastrissamine (jaspine B). *J. Org. Chem.* **78**, 8208–8213 (2013).
- Crabtree, E. V. *et al.* Synthesis of the enantiomers of XYLNAc and LYXNAc: comparison of β -N-atecylhexosaminidase inhibition by the 8 stereoisomers of 2-*N*-acetylaminio-1,2,4-trideoxy-1,4-iminopentitols. *Org. Biomol. Chem.* **12**, 3932–3943 (2014).
- Saotome, C., Wong, C.-H. & Kanie, O. Combinatorial library of five-membered iminocyclitol and the inhibitory activities against glycol-enzymes. *Chem. Biol.* **8**, 1061–1070 (2001).
- Enders, D., Paleček, J. & Grondal, C. A direct organocatalytic entry to sphingoids: asymmetric synthesis of D-arabino- and L-ribo-phyto sphingosine. *Chem. Commun.* **6**, 655–657 (2006).
- Brandi, A., Cardona, F., Cicchi, S., Cordero, F. M. & Goti, A. Stereocontrolled cyclic nitron cycloaddition strategy for the synthesis of pyrrolizidine and indolizidine alkaloids. *Chem. Eur. J.* **15**, 7808–7821 (2009).
- Medjahdi, M., González-Gómez, J., Foubelo, F. & Yus, M. Stereoselective synthesis of azetidines and pyrrolidines from *N*-tert-butylsulfonyl(2-aminoalkyl)oxiranes. *J. Org. Chem.* **74**, 7859–7865 (2009).
- Davis, F. A., Yang, B. & Deng, J. Asymmetric synthesis of *cis*-5-*tert*-butylproline with metal carbenoid NH insertion. *J. Org. Chem.* **68**, 5147–5152 (2003).

31. Dangerfield, E. M., Timmer, M. S. M. & Stocker, B. L. Total synthesis without protecting groups: pyrrolidines and cyclic carbamates. *Org. Lett.* **11**, 535–538 (2009).
32. Deukeleire, S., D'hooghe, M., Törnroos, K. W. & De Kimpe, N. Stereoselective synthesis of chiral 4-(1-chloroalkyl)- β -lactams starting from amino acids and their transformation into functionalized chiral azetidines and pyrrolidines. *J. Org. Chem.* **75**, 5934–5940 (2010).
33. Lauritsen, A. & Madsen, R. Synthesis of naturally occurring iminosugars from d-fructose by the use of a zinc-mediated fragmentation reaction. *Org. Biomol. Chem.* **4**, 2898–2905 (2006).
34. Lovering, F., Bikker, J. & Humblet, C. Escapade from flatland: increasing saturation as an approach to improving clinical success. *J. Med. Chem.* **52**, 6752–6756 (2009).
35. Walters, W. P. Going further than Lipinski's rule in drug design. *Exper. Opin. Drug. Discov.* **7**, 99–107 (2012).
36. Feher, M. & Schmidt, J. M. Property distributions: differences between drugs, natural products, and molecules from combinatorial chemistry. *J. Chem. Inf. Comput. Sci.* **43**, 218–227 (2003).
37. Pascolutti, M. & Quinn, R. J. Natural products as lead structures: chemical transformations to create lead-like libraries. *Drug Discov. Today* **19**, 215–221 (2014).
38. Verhelst, S. H. L. *et al.* A Short route toward chiral, polyhydroxylated indolizidines and quinolizidines. *J. Org. Chem.* **68**, 9598–9603 (2003).
39. Jiangseubchatveera, N. *et al.* Concise synthesis of (–)-steviamine and analogues and their glycosidase inhibitory activities. *Org. Biomol. Chem.* **11**, 3826–3833 (2013).
40. Parmeggiani, C., Cardona, F., Giusti, L., Reissig, H.-U. & Goti, A. Stereocomplementary routes to hydroxylated nitrogen heterocycles: total syntheses of casuarine, australine, and 7-*epi*-australine. *Chem. Eur. J.* **19**, 10595–10604 (2013).
41. Wardrop, D. J. & Bowen, E. G. Nitrenium ion-mediated alkene bis-cyclofunctionalization: total synthesis of (–)-swainsonine. *Org. Lett.* **13**, 2376–2379 (2011).
42. Randl, S. & Blechert, S. Concise enantioselective synthesis of 3,5-dialkyl-substituted indolizidine alkaloids via sequential cross-metathesis-double-reductive cyclization. *J. Org. Chem.* **68**, 8879–8882 (2003).
43. Garber, S. B., Kingsbury, J. S., Gray, B. L. & Hoveyda, A. H. Efficient and recyclable monomeric and dendritic Ru-based metathesis catalysts. *J. Am. Chem. Soc.* **122**, 8168–8179 (2000).
44. Gómez, L. *et al.* Chemoenzymatic synthesis, structural study and biological activity of novel indolizidine and quinolizidine iminocyclitols. *Org. Biomol. Chem.* **10**, 6309–6321 (2012).
45. Izquierdo, I., Plaza, M. T., Tamayo, J. A., Franco, F. & Sánchez-Cantalejo, F. Total synthesis of natural (+)-hyacinthacine A₆ and non-natural (+)-7a-*epi*-hyacinthacine A₁ and (+)-5,7a-*diepi*-hyacinthacine A₆. *Tetrahedron* **66**, 3788–3794 (2010).
46. Brock, E. A., Davies, S. G., Lee, J. A., Roberts, P. M. & Thomson, J. E. Asymmetric synthesis of polyhydroxylated pyrrolizidines via transannular iodoamination with concomitant *N*-debenzylation. *Org. Lett.* **13**, 1594–1597 (2011).

Acknowledgements

This work was supported by an NSERC Discovery Grant to R.B., a Michael Smith Foundation for Health Research Career Investigator Award to R.B., a Multi-Investigator Research Initiative grant (funded by Brain Canada, Genome BC, Michael Smith Foundation for Health Research; and the Pacific Alzheimer Research Foundation), an NSERC PGSD for M.B.-B.; and a NSERC USRA for M.M.

Author contributions

R.B. and M.B.-B. conceived the experiments and R.B. prepared the manuscript. M.M. assisted with the experiments.

Additional information

Accession codes: The X-ray crystallographic coordinates for structures reported in this study have been deposited at the Cambridge Crystallographic Data Centre (CCDC), under deposition numbers 1038918–1038924. These data can be obtained free of charge from CCDC via www.ccdc.cam.ac.uk/data_request/cif.

Supplementary Information accompanies this paper at <http://www.nature.com/naturecommunications>

Competing financial interests: The authors declare no competing financial interests.

Reprints and permission information is available at <http://npg.nature.com/reprintsandpermissions/>

How to cite this article: Bergeron-Brele, M. *et al.* Direct synthesis of imino-*C*-nucleoside analogues and other biologically active iminosugars. *Nat. Commun.* **6**:6903 doi: 10.1038/ncomms7903 (2015).

This work is licensed under a Creative Commons Attribution 4.0 International License. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in the credit line; if the material is not included under the Creative Commons license, users will need to obtain permission from the license holder to reproduce the material. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>