GLOSSARY

WISCONSIN CARD SORTING TEST (WCST)

Neuropsychological test used to assess rule attainment and cognitive set shifting ability (switching between tasks based on environmental feedback)

VERBAL FLUENCY TASK (VFT)

Neuropsychological test used to measure verbal initiation and ability to retrieve words based on a certain condition such as words beginning with the letter A, or categories such as fruit or animals

STROOP COLOR WORD TEST (SCWT)

Neuropsychological test that measures interference in the reaction time of a task; patients are asked to identify font color of words printed in the color identified by the word (congruent condition), and to identify font color of words printed in a different color to that identified by the word (incongruent condition)

the specificity of MRI BC was high—88.2%—the sensitivity was only 74.3%. The OCGB test had a specificity of 94.1% and a sensitivity of 91.4%, and simultaneous use of the two tests gave 88.2% specificity and 97.1% sensitivity.

The authors conclude that use of the OCGB and MRI BC tests simultaneously can allow neurologists to confidently predict conversion from CIS to CDMS, although they note that their results should be corroborated with results from multiple studies.

Christine Kyme

Original article Masjuan J *et al.* (2005) Clinically isolated syndromes: a new oligoclonal band test accurately predicts conversion to MS. *Neurology* **66**: 576–578

Loss of expectation in Alzheimer's disease reduces the efficacy of analgesics

In a recent study reported in *Pain*, researchers show for the first time that the mechanisms that underlie expectation of drug effects can be disrupted in a neurological disorder, leading to a reduction in drug efficacy.

Benedetti *et al.* used the open-hidden paradigm to investigate the psychological element of the effects of an analgesic in patients with early-stage Alzheimer's disease (AD; mean age 73.5 years) and 16 healthy matched controls. Participants were either openly given a topical treatment of 1% lidocaine for their pain after venipuncture, or were given the drug 'hidden' in a plaster tape. They were asked to rate the severity of their pain before, and 15 min after, lidocaine application. Frontal executive function was determined with the Frontal Assessment Battery (FAB) and electroencephalogram connectivity analysis.

In all participants, open application of lidocaine at the start of the study resulted in a significantly larger analgesic effect than hidden application ($P < 0.001$). A year later, however, the effects of the open application were reduced in AD patients ($P < 0.001$), thought to reflect progression of their disease. Lower FAB scores correlated with reduced analgesia after open application in patients with AD ($P < 0.02$). Patients with reduced prefrontal connectivity showed the lowest FAB scores and the poorest analgesic response.

The authors propose that the loss of expectation mechanisms in patients with AD could also affect the efficacy of other drug therapies

used in these patients, and they suggest that other conditions in which the prefrontal lobes are affected might also merit investigation.

Christine Kyme

Original article Benedetti F *et al.* (2006) Loss of expectation-related mechanisms in Alzheimer's disease makes analgesic therapies less effective. *Pain* **121**: 133–144

Differing cognitive profiles in bipolar disorder and schizophrenia

It is thought that differences in the neural circuitry within the prefrontal cortex might contribute to the different cognitive profiles in bipolar disorder and schizophrenia. To investigate this hypothesis, researchers in the UK have compared the cognitive profiles of patients with these neuropsychological disorders.

In total, 44 patients with stable bipolar disorder, 54 patients with schizophrenia, and 46 healthy controls were enrolled in the study. Each participant performed three tests—the WISCONSIN CARD SORTING TEST (WCST), the VERBAL FLUENCY TASK (VFT) and the STROOP COLOR WORD TEST (SCWT)—all of which engage different neural regions within the frontal lobe.

The schizophrenia group contained a significantly higher proportion of males than the bipolar disorder group ($P = 0.001$). Results from the WCST were similar in all three groups. In the VFT, however, patients with schizophrenia and those with bilateral disorder made more errors than controls; verbal impairment was considerably more marked in schizophrenic patients than in those with bipolar disorder ($P < 0.003$ for all conditions). In the SCWT, patients with schizophrenia performed less well than controls in both congruent and incongruent conditions ($P < 0.001$ and $P = 0.02$, respectively), and less well than those with bipolar disorder in the congruent condition ($P = 0.035$), but not the incongruent condition.

Frangou *et al.* conclude that patients with schizophrenia and bipolar disorder differ in performance in verbal fluency and the SCWT. This variation might be caused by differences in disruption to neural circuitry, but the authors state that neuroimaging studies are needed to determine precise locations of dysfunction.

Rebecca Ireland

Original article Frangou S *et al.* (2006) Frontotemporal function may distinguish bipolar disorder from schizophrenia. *Bipolar Disord* **8**: 47–55