Efaproxiral is an effective addition to WBRT in patients with brain metastases

Metastasis of cancer to the brain is common, affecting as many as 170,000 individuals annually in the US alone. Whole-brain radiation therapy (WBRT) produces local control/response rates of 50–75%, and other treatments have also shown some success; however, post-diagnosis survival is poor, at a median of approximately 4.5 months. Because tumor resistance to radiation is increased by hypoxia, treatments to reduce hypoxia could be valuable. Efaproxiral changes the conformational structure of hemoglobin, thereby assisting oxygen release and improving tumor oxygenation. In the REACH/RT-009 study, Suh *et al.* tested whether addition of efaproxiral to WBRT and supplemental oxygen resulted in longer survival than combined WBRT and oxygen.

The 1:1 randomized, international, phase III study analyzed data from 515 patients with brain metastases originating from solid tumors; of these, 397 had breast cancer (BC) or non-small-cell lung cancer (NSCLC). Patients underwent 2 weeks of treatment with ($n=265$) or without ($n=250$; controls) efaproxiral. Efaproxiral-treated patients survived longer than controls (median 5.4 months vs 4.4 months; 6.0 months vs 4.4 months in the subset of patients with NSCLC/BC), but the treatment benefit was statistically insignificant. By contrast, regression analysis showed a statistically significant reduction in the likelihood of efaproxiral-treated patients to die compared with controls (–26%; $P=0.003$); this reduction in risk of death might, however, be attributable to patients with primary BC, who showed the greatest treatment benefit. Response rates also improved in the efaproxiral arm, with significant improvements seen in those with NSCLC/BC ($P=0.01$).

A further trial, the ENRICH RT-016 study, is therefore underway to investigate efaproxiral-related improvements in survival duration and response to WBRT in patients with brain metastases from BC.

Pippa Murdie

Original article Suh JH *et al.* (2006) Phase III study of efaproxiral as an adjunct to whole-brain radiation therapy for brain metastases. *J Clin Oncol* 24: 106–114

Treatment of malignant glioma with high-dose conformal radiotherapy

Patients with malignant glioma are usually treated with postoperative chemoradiotherapy; however, tumors often recur at the primary location. Improvements in radiation technology aim to minimize damage to surrounding healthy tissue, which might enable higher doses of radiation to be administered without an increase in the risk of cognitive impairment or necrosis. To investigate the effect of radiation dose on tumor recurrence, Tanaka *et al.* compared the outcomes of malignant glioma patients treated with conventional radiotherapy with those treated with high-dose conformal radiotherapy.

Patients treated with high-dose radiotherapy (80–90 Gy) within the years 1990–2002 ($n=90$; 29 with anaplastic astrocytoma and 61 with glioblastoma) were compared with patients treated with conventional radiotherapy (60 Gy) within the years 1979–1989 ($n=94$; 34 with anaplastic astrocytoma and 60 with glioblastoma). After a median follow-up of 12 months, treatment with high-dose radiotherapy resulted in significantly longer patient survival than conventional radiotherapy (adjusted hazard ratio for death 0.30, 95% CI 0.12 to 0.76, $P=0.011$ for anaplastic astrocytoma; and adjusted hazard ratio 0.49, 95% CI 0.28 to 0.87, $P=0.014$ for glioblastoma). High-dose radiotherapy also increased the duration of progression-free survival in patients with anaplastic astrocytoma, but not in those with glioblastoma. White matter abnormalities were more common in the high-dose group than in the low-dose group ($P=0.0002$), but toxic effects were generally tolerable.

In summary, compared with conventional 60 Gy radiotherapy, high-dose 80–90 Gy conformal radiotherapy improved survival. Patients treated with high-dose radiotherapy were less likely to have tumor recurrence at the primary site, indicating that higher doses of radiation might be more effective at suppressing tumor regrowth.

Kate Matthews

Original article Tanaka M *et al.* (2006) High-dose conformal radiotherapy for supratentorial malignant glioma: a historical comparison. *Lancet Oncol* 6: 953–960

GLOSSARY

REACH

Radiation Enhancing
Allosteric Compound for
Hypoxic Brain Metastases