GLOSSARY

LATE-REACTING
NORMAL TISSUES

Tissues with a low cell turnover rate (e.g. lung, kidney and spinal cord); these tissues do not regenerate as quickly as early-reacting tissues and therefore take longer to recover from radiation damage

RECIST CRITERIA

Criteria for measuring tumor shrinkage as an indicator of antitumor activity, as evaluated by the Response Evaluation Criteria in Solid Tumors Group (RECIST)

78% chance of remaining progression-free during that time.

In conclusion, a small proportion of NSCLC patients receiving palliative radiotherapy experience prolonged survival, or possibly cure. The authors are unsure why this phenomenon occurs, but speculate that these patients might have particularly radiosensitive tumors or tumors with low metastatic tendencies. They stress that their study does not indicate that lower radiotherapy doses should be used when trying to cure a patient, but state that radiation oncologists should be aware of the possibility of long-term survival, and keep doses to LATE-REACTING NORMAL TISSUES to within tolerable levels.

Rebecca Ireland

Original article Mac Manus MP *et al.* (2006) Unexpected long-term survival after low-dose palliative radiotherapy for nonsmall cell lung cancer. *Cancer* 106: 1110–1116

^{99m}Tc-DS predicts response to endocrine therapy in advanced breast cancer patients

New research shows promise for ^{99m}Tc-depreotide scintigraphy (^{99m}Tc-DS) in assessing the effectiveness of endocrine therapy in patients with advanced breast cancer. Inactivation of the estrogen receptor by endocrine therapy can downregulate somatostatin receptor 2 expression at the cell surface. Through *in vivo* imaging, ^{99m}Tc-DS can detect this downregulation, which the authors propose is an indicator of response to endocrine treatment.

Eighteen patients were assessed using ^{99m}Tc-DS before initiation of endocrine therapy. After 6 months of treatment, 8 patients had stable disease according to the RECIST CRITERIA (responders; 5 receiving first-line treatment and 3 receiving second-line treatment) and 10 had progressive disease (nonresponders). Pre-therapy scans predicted response with high accuracy: all responders had positive scans and 7 of the 10 nonresponders had negative scans. Follow-up scans, performed in 16 patients 3 weeks after treatment initiation, matched the results of the pre-therapy scans in all cases. The change in ^{99m}Tc-depreotide uptake between the first and second scans was significantly different between responders and nonresponders ($P=0.017$), with uptake decreasing in the former and increasing in the latter. Pre-therapy scan results combined with changes in tracer uptake

between the two scans showed 100% accuracy for predicting endocrine response.

^{99m}Tc-DS can therefore evaluate response to therapy from as little as 3 weeks after treatment initiation. The authors propose this method as a more accurate and noninvasive alternative to *in vitro* assessment of hormone receptor status by ligand-binding assay or immunohistochemistry. A larger trial is needed to validate the results of this small study.

Pippa Murdie

Original article Van Den Bossche B *et al.* (2006) Early prediction of endocrine therapy effect in advanced breast cancer patients using ^{99m}Tc-depreotide scintigraphy. *J Nucl Med* 47: 6–13

Successful antifungal prophylaxis for radiotherapy-induced mucositis

Radiotherapy for head and neck malignancy is associated with oral mucositis, a toxicity that can impact severely on quality of life. Infection of the oral mucosa by *Candida* species is also common during head and neck radiotherapy and is thought to exacerbate the effects of mucositis. A recent paper by researchers in Greece has shown that fluconazole prophylaxis reduces the incidence of both oral mucositis and candidiasis, and consequently the number of treatment interruptions in patients receiving radiotherapy for head and neck cancer.

In this nonblinded study, 34 patients received daily fluconazole from initiation to completion of radiotherapy (group A). A further 29 patients received no prophylactic treatment during radiotherapy, but were administered fluconazole daily for 1 week if candidiasis developed (group B). Smears to assess oral *Candida* burden were performed before and after radiotherapy, and patients underwent weekly oral examinations. Baseline *Candida* load was similar between the two groups. At completion of radiotherapy, the incidence of mucositis was significantly lower in group A than in group B ($P=0.018$). None of the patients had radiotherapy interrupted in group A, whereas 17.2% of patients in group B had treatment interrupted because of mucositis ($P=0.017$). The week of onset of mucositis did not differ significantly between the two groups. Fluconazole prophylaxis prevented candidiasis in all of the patients in group A, whereas 10 patients in group B suffered from this infection.