GLOSSARY

ATBC STUDY

A study conducted in Finland that tested whether alpha-tocopherol or beta-carotene could reduce the incidence of cancer in male smokers

activity ($P < 0.005$); deiodinases 1 and 3 could not be tested in this system. Linkage and sequencing analyses showed that deiodinase gene sequences were normal, but that affected children had a homozygous mutation in *SECISBP2* (selenocysteine-incorporation sequence binding protein 2). The four unaffected siblings and the parents were heterozygous carriers of this mutation, which was not present in controls.

The authors also identified a compound heterozygous mutation in *SECISBP2* in a child of Irish origin, who had similar thyroid-hormone abnormalities to those of the Bedouin children. All four affected children had low levels of two other selenocysteine-containing proteins; however, the effects on thyroid-hormone metabolism seem to predominate. The authors suggest that the relatively mild phenotype could arise because *SECISBP2* function is not totally lost.

Ian Newman

Original article Dumitrescu AM *et al.* (2005) Mutations in *SECISBP2* result in abnormal thyroid hormone metabolism. *Nat Genet* 37: 1247–1252

Role of TSH-releasing hormone in thyroid gland regulation

Secretion of TSH from the pituitary gland is regulated positively by TSH-releasing hormone (TRH) and negatively by thyroid hormone (TH), which also negatively regulates TRH. It has previously been thought that TH was the major regulator of thyroid activity, but this has yet to be demonstrated *in vivo*. Nikrodhanond *et al.* investigated thyroid regulation in knockout mice lacking TRH ($TRH^{-/-}$), the β isoforms of TH receptors ($TR\beta^{-/-}$), or both (double knockout).

TH and TSH levels were elevated in $TR\beta^{-/-}$ mice compared with wild-type mice, because the absence of the TH receptor abolished negative regulation by TH. In $TRH^{-/-}$ mice, the absence of positive regulation by TRH was reflected in the reduced levels of serum TH. The response of double-knockout mice was similar to that of $TRH^{-/-}$ mice, indicating that positive regulation by TRH is the most important element in controlling thyroid function. Unexpectedly, while diet-induced hypothyroidism caused a significant increase in serum TSH levels in all other genotypes, a much lower response was observed in double-knockout mice, which

also behaved similarly to $TR\beta^{-/-}$ mice in being unresponsive to thyroid hormone. This result indicates a possible role for the unliganded TH receptor in TSH expression.

The authors suggest that the hypothyroid TSH response might be mediated by interaction between the TRH and TH pathways. They conclude that reduction in TSH bioactivity in the absence of TRH and the aberrant response to hypothyroidism of double-knockout mice point to a dominant role for TRH in thyroid hormone regulation.

Jim Casey

Original article Nikrodhanond AA *et al.* (2005) Dominant role of thyrotropin-releasing hormone in the hypothalamic–pituitary–thyroid axis. *J Biol Chem* [doi: 10.1074/jbc.M511530200]

Insulin and risk of pancreatic cancer

Type 2 diabetes has been associated with pancreatic cancer, but it is unclear whether diabetes is a causative factor or the result of subclinical malignancy. Insulin has been shown to have mitogenic effects on pancreatic cancer cells, and hyperinsulinemia is known to occur in the early stages of type 2 diabetes, suggesting a mechanism by which insulin might have a carcinogenic effect in the pancreas. Stolzenberg-Solomon *et al.* sought to determine whether prediagnostic glucose and insulin levels and insulin resistance were linked to subsequent incidence of pancreatic cancer in a cohort of Finnish male smokers participating in the ATBC STUDY. Only cancer cases that occurred after the fifth year of follow-up were included in the study.

Adjusting for age, smoking and BMI, participants whose fasting insulin concentration was in the highest quartile were found to have double the risk of pancreatic cancer of those in the lowest quartile. Diabetes was found to double the risk of pancreatic cancer, while high serum glucose levels and insulin resistance were also positively associated with this cancer. Association of risk factors with cancer was found to be even stronger in cases that occurred after more than 10 years of follow-up.

The authors point to several *in vitro* and animal studies that suggest a carcinogenic effect of insulin on pancreatic cells, and conclude that the association between type 2 diabetes, hyperglycemia and pancreatic

cancer observed in previous studies might be explained by the effects of high insulin concentrations in the pancreas.

Jim Casey

Original article Stolzenberg-Solomon RZ *et al.* (2005) Insulin, glucose, insulin resistance, and pancreatic cancer in male smokers. *JAMA* 294: 2872–2878

Are renoprotection guidelines followed in elderly patients with diabetes?

US guidelines recommend that diabetic patients with nephropathy, hypertension, or both, use angiotension-converting-enzyme (ACE) inhibitors or angiotension-II receptor blockers (ARBs) to help slow progression of chronic kidney disease. Winkelmayer *et al.* used Medicare data to explore application of these guidelines to treatment of elderly diabetic patients (≥ 65 years) who were eligible to receive comprehensive drug treatment from the state of Pennsylvania in 2002. Patients with end-stage renal disease were excluded from the analysis.

The primary outcome was the filling of a prescription for either an ARB or an ACE inhibitor during January–March 2003. Of the 30,750 eligible diabetic patients (mean age 78.4 ± 6.4 years, 81% female), 21,138 had received a diagnosis of hypertension, nephropathy, or both. Although these patients were eligible for comprehensive drug treatment with minimal copayments, only 50.7% of diabetic patients with hypertension and/or proteinuria filled a prescription for an ACE inhibitor or ARB during the 3-month period. Older patients (75–84 years and ≥ 85 years) were significantly less likely to fill a prescription for the drugs than patients aged 65–74 years ($P=0.01$ and $P<0.001$, respectively). Patients with proteinuria but not hypertension were less likely to fill a prescription for the drugs than patients with hypertension only ($P=0.03$).

The authors conclude that renoprotection guidelines were not being followed in the elderly diabetic patients they studied, but acknowledge that the study did not assess whether this was because drugs were not prescribed by physicians or because prescriptions were not filled by patients.

Rebecca Ireland

Original article Winkelmayer WC *et al.* (2005) Underuse of ACE inhibitors and angiotensin II receptor blockers in elderly patients with diabetes. *Am J Kidney Dis* 46: 1080–1087

Cortisol levels can predict future cognitive decline in older individuals

Two prospective longitudinal studies recently reported results indicating that levels of cortisol in cognitively intact older people can predict their possible future cognitive decline. Previous studies have implicated elevated circulating cortisol levels in the pathogenesis of impaired cognition.

In a study by Karlamangla and colleagues, researchers measured the resting overnight urinary cortisol levels of 538 individuals (55% female, 45% male) aged 70–79 years with high cognitive function. Neuropsychological testing was used to assess their baseline cognitive function, and follow-up testing was carried out 3 and 7 years later to assess any changes. Cortisol measurements for participants were divided into quartiles for analysis.

Those in the upper three quartiles were found to be at higher risk of incident cognitive impairment than those in the lowest quartile (16% vs 9%), with similar risks for men and women. Adjustment for potential confounders, such as age at baseline, did not greatly affect the association.

In a separate study, Li *et al.* collected salivary cortisol samples from 79 cognitively intact participants aged 63 years or over. Samples were taken at 08:00 h, 15:00 h and 23:00 h, and the procedure was repeated annually for 3 years. Neuropsychological tests were carried out 2–3 days after collection of samples. For the 46 participants (52% female, 48% male) who completed the study, a higher baseline cortisol concentration at 23:00 h predicted a decline in memory as measured by delayed paragraph recall performance.

In the effort to develop strategies for halting cognitive decline in older adults, it is suggested that further research should be undertaken to develop methods of attenuating the detrimental effects of higher cortisol levels in this population.

Christine Kyme

Original articles Karlamangla AS *et al.* (2005) Urinary cortisol excretion as a predictor of incident cognitive impairment. *Neurobiol Aging* 26 (Suppl): S80–S84
Li G *et al.* (2005) Salivary cortisol and memory function in human aging. *Neurobiol Aging* [doi: 10.1016/j.neurobiolaging.2005.09.031]