GLOSSARY

D3 LYMPH-NODE DISSECTION

A D1 dissection with additional removal of the lymph nodes around the blood vessels supplying the stomach that arise from the celiac axis, those in the hepatoduodenal ligament and retropancreatic region, and those surrounding the superior mesenteric vein

D1 LYMPH-NODE DISSECTION

Removal of the perigastric lymph nodes along the lesser and greater curvatures of the stomach

BORRMANN'S CLASSIFICATION

System for classifying gastric carcinomas as polypoid (type I), ulcerating (type II), ulcerating-infiltrating (type III) or infiltrating (type IV)

increased acute toxicities seen in the concomitant arm are a possible cause for concern.

Mitoxantrone is no longer available as an adjuvant therapy for breast cancer, and Rouëssé and his group advocate that concomitant adjuvant chemoradiotherapy with other drugs should be studied further. In particular, the taxanes recently approved for adjuvant therapy have already shown promise for chemoradiotherapy in some trials.

Pippa Murdie

Original article Rouëssé J *et al.* (2006) A phase III randomized trial comparing adjuvant concomitant chemoradiotherapy versus standard adjuvant chemotherapy followed by radiotherapy in operable node-positive breast cancer: final results. *Int J Radiat Oncol Biol Phys* **64**: 1072–1080

Extensive lymph-node dissection improves survival in patients with gastric cancer

A single-center, randomized controlled trial conducted by researchers from Taiwan has shown that patients with gastric cancer who undergo extensive, D3 LYMPH-NODE DISSECTION have higher survival rates than those who receive limited, D1 LYMPH-NODE DISSECTION.

The study by Wu *et al.* randomized 221 patients with gastric adenocarcinoma to receive gastrectomy with either a D1 or D3 dissection at Taipei Veterans General Hospital. Each procedure was carried out by one of three surgeons, all of whom had received extensive training and had experience of at least 25 D3 dissections.

After a median follow-up of 94.5 months, 5-year disease-free survival was 59.5% in patients who underwent D3 surgery, compared with 53.6% in the D1 group ($P=0.041$). Hazard analysis revealed that patients randomized to D3 surgery were almost 50% less likely to die within 5 years than those assigned to D1 surgery. After 5 years, disease had recurred in 50.6% of patients in the D1 group, compared with 40.3% in the D3 group, but the difference was not statistically significant. Multivariate analysis showed that D1 surgery was associated with poor survival, as were nodal disease, tumors occupying the whole stomach, and type III or IV tumor appearance by BORRMANN'S CLASSIFICATION. The authors conclude that D3 lymph-node dissection improves both overall and disease-free survival in patients with gastric cancer, but emphasize that, to minimize

morbidity, the procedure should be performed only by well-trained, experienced surgeons.

Alexandra King

Original article Wu C-W *et al.* (2006) Nodal dissection for patients with gastric cancer: a randomised controlled trial. *Lancet Oncol* **7**: 309–315

Dose-escalated IMRT improves survival and local control in T3–T4 nasopharyngeal carcinoma

Radiation-dose escalation has the potential to reduce local disease recurrence rates in patients with locally advanced nasopharyngeal carcinoma. Previous studies using conventional radiotherapy, however, have been largely unsuccessful, with considerable levels of radiation toxicity. Kwong and colleagues have addressed the need for more-precise treatment by studying dose escalation using the simultaneous modulated accelerated radiation therapy (SMART) boost technique with intensity-modulated radiotherapy (IMRT) in 50 patients with stage T3–T4 nasopharyngeal carcinoma.

All patients received a 70Gy dose to the planning target volume and a 76Gy dose to the gross tumor volume delivered in 35 fractions over 7 weeks. Thirty-four patients also received concurrent cisplatin chemotherapy. Patients were assessed by regular CT or MRI scans, and underwent nasopharyngoscopy and biopsy at 6 and 8 weeks after completion of IMRT. After a median follow-up of 25 months, two patients had local disease recurrence and two had distant recurrence; all four patients had undergone IMRT without concurrent chemotherapy. The 2-year disease-free survival, overall survival, distant-metastases-free survival and loco-regional control rates were 93.1%, 92.1%, 94.2%, and 95.7%, respectively. Toxicities were common, but manageable, and 92% of patients completed IMRT without interruption. Notable late complications included two cases of carotid pseudoaneurysm. The authors conclude that this chemoradiotherapy regimen controls local disease recurrence effectively and improves survival rates in patients with locally advanced nasopharyngeal carcinoma, and therefore recommend the regimen in this setting.

Alexandra King

Original article Kwong DLW *et al.* (2006) Preliminary results of radiation dose escalation for locally advanced nasopharyngeal carcinoma. *Int J Radiat Oncol Biol Phys* **64**: 374–381