GLOSSARY

PSA VELOCITY (PSAV)

The rate at which serum PSA level increases over time

PSA DOUBLING TIME (PSADT)

The time taken for the detected PSA level to double. This test is used as an indicator of the rate of disease progression in patients with prostate cancer

Ultrasound-targeted needle biopsy could improve detection of prostate cancer

In the diagnosis of prostate cancer, it is common for needle biopsies to be performed in a systematic distribution using a sextant approach (6 cores), a systematic pattern of up to 12 cores throughout the prostate, or up to 40 cores in a 'saturation' approach. A targeted method that reduces the number of biopsies taken might decrease cost and patient morbidity. Halpern *et al.* therefore compared an ultrasound-targeted protocol (biopsies taken from areas of greatest enhancement on ultrasonography) with a modified sextant approach (a laterally distributed systematic set of cores taken from the area of greatest blood flow in each sextant) for the detection of prostate cancer.

Cancer was detected in 104 of 301 men (363 biopsy cores). Ultrasound-targeted biopsies were twice as likely as the modified sextant approach to detect cancer when it was present ($P < 0.001$); although a 'by patient' analysis showed that both techniques had similar rates of false-negative results, ultrasound-targeted biopsy was more effective at the base of the prostate, and modified sextant biopsy at the apex ($P = 0.006$). In terms of ultrasound imaging, modalities using intermittent harmonic imaging were superior to baseline gray-scale and Doppler imaging in highlighting affected areas for biopsy ($P < 0.05$).

As both biopsy techniques were targeted to some degree, the authors suggest the advantage of ultrasound-targeted biopsy over traditional, nondirected sextant biopsy is even greater than that reported here. So that carcinomas in the apex of the prostate are not missed, they recommend that contrast-enhanced ultrasound-targeted biopsy be complemented by additional systematic cores in the apex.

Rebecca Doherty

Original article Halpern EJ *et al.* (2005) Detection of prostate carcinoma with contrast-enhanced sonography using intermittent harmonic imaging. *Cancer* **104**: 2373–2383

Stone-to-skin distance predicts success of shock-wave lithotripsy for kidney stones

A simple and easily reproducible predictor of the success of extracorporeal shock-wave lithotripsy (ESWL) in the treatment of urinary stone disease

is needed, to determine which patients might be better treated by other methods. Pareek *et al.* propose that measurement of stone-to-skin distance (SSD) using non-contrast-enhanced CT might be such a tool.

This retrospective study included 64 patients who had undergone ESWL between March 2000 and April 2004 for single lower-pole kidney stones 5–15 mm in diameter. SSD was calculated from the mean of three measurements, taken at true posterior, 45° lateral and true lateral. Stone density and patient BMI were also calculated. Following radiographic assessment 6 weeks after treatment, patients were classified as being stone-free ($n = 30$) or having residual stones ($n = 34$).

The mean SSD, stone density and BMI were all lower in the stone-free than in the residual-stone group ($P < 0.01$, $P < 0.05$ and $P < 0.05$, respectively); however, after logistic regression analysis, SSD was the best statistically significant predictor of ESWL outcome (odds ratio 0.32; $P < 0.01$). SSD in patients in whom ESWL failed was most commonly ≥ 10 cm.

The authors postulate that the shock wave emitted by the lithotripter becomes less effective as the distance between the stone and skin increases. They suggest the maximal treatment distance recommended by the manufacturer of a lithotripsy unit may be inaccurate in some cases, and that each unit should have its specific optimal distance determined. Further investigation of the predictive value of SSD is warranted, particularly in larger, prospective studies.

Rebecca Doherty

Original article Pareek G *et al.* (2005) Shock wave lithotripsy success determined by skin-to-stone distance on computed tomography. *Urology* **66**: 941–944

Preoperative PSA kinetics predict prostate cancer outcomes

Several studies have suggested that PSA kinetics might predict prostate cancer outcomes, although results have been conflicting. The increase in PSA over time (PSA VELOCITY; PSAV) is thought to reflect cancer growth; however, in men with untreated prostate cancer, PSA rises exponentially, rather than linearly. PSA DOUBLING TIME (PSADT) might, therefore, be a more accurate reflection of cancer growth.