Corrigendum: A transactivation-deficient mouse model provides insight into p53 regulation and function

G S Jimenez, M Nister, M Beeche, J S Stommel, E Barcarse, S O’Gorman & G M Wahl

Nat. Genet. **26**, 37–43 (2000).

This article described the generation of a mouse in which homologous recombination was used to change codons 25 and 26 of the gene *Trp53* to glutamine and serine. The initial sequencing of the full-length cDNA from mouse embryonic fibroblasts expressing this allele was considered to be wild-type at all codons except for 25 and 26. A reanalysis of the sequence showed that this mutant also contains a valine at codon 135, which is referred to as a “provisional wild type codon” in the National Center for Biotechnology Information’s LocusLink tool (accession numbers: *Trp53* mRNA, NM_011640; p53 protein, NP_035770). Sequence analysis showed that the lambda genomic clone containing an *EcoRI* fragment encompassing *Trp53* (*Nature* **356**, 215–221; 1992) that was used to prepare the targeting construct contained the Val135 codon, whereas *Trp53* in laboratory mice (*e.g.*, C57Bl6) encodes alanine at codon 135 (K. Krummel, F. Toledo, C. Lee & G.M.W., unpublished data). The properties of the two proteins have been investigated by M.N. *et al.* (*Oncogene*, in the press).

Erratum: A treasury of exceptions

Editorial

Nat. Genet. **36**, 1239 (2004).

The ENCODE project (<http://www.genome.gov/ENCODE/>) is funded and managed by the National Human Genome Research Institute at the US National Institutes of Health.