Abstracts: Session III

sequence tags that allows efficient selection of new potential tumor-associated genes. The resulting ranked list of candidates is analyzed in terms of its content in known genes, used as controls, and the new candidates are further tested for selective expression in normal and tumor tissues by real-time polymerase chain reaction with reverse transcription. The most highly tumor-specific candidates are then pursued to full-length cloning. We present the discovery of a series of new genes that are upregulated in colon cancer.

Visvanathan, Jaya

[56]

Genomic aberrations in gastric adenocarcinoma

Jaya Visvanathan¹, Pulivarthi H. Rao², Christina Rudduck³, Wai Keong Wong³, Chow Yin Wong³, Jean Ho³, Choon Wei Wee¹, Oi Lian Kon¹ & Dennis T.H. Lim¹

¹National Cancer Centre, Singapore

²Baylor College of Medicine, Houston, Texas, USA

³Singapore General Hospital, Singapore

Gastric adenocarcinoma (GA) presents a challenge in understanding the complex interactions of environmental factors with genotype in carcinogenesis. The National Cancer Centre has taken a multifaceted approach to this problem with comparative genomic hybridization of freshly resected GAs. Data from nine cases of intestinal and five cases of diffuse histological types showed the common aberrations to be loss of 4pq (four cases) and gains of 17pq (five cases) and 20pq (four cases). We also documented amplifications in 22q (six cases), 20q (five cases) and 1p (two cases). High-level gains were present in 20q11.2q12 (three cases), 10q24.3q25.1 (one case) and 12q14 (one case). Two cases of diffuse GA showed no change. High-level amplifications in 10q24.3q25.1 and 12q14 have not been reported previously. High-level amplification of 12q14 could be associated with overexpression of *MDM2*, *CDK4* and *RAP1B*. Three cytochrome P450 hydroxylases map to 10q24. Loss of 4pq is a common abnormality in cases of GA; its biological significance may be related to loss of function of caspase 3, caspase 6 and PTPN13, which is frequently deleted in liver and ovarian cancers. High-level gain of 20q or 20pq was another striking finding of this preliminary study. Amplified genes of probable importance from this genomic region are those for DNA methyltransferase 3B, E2F transcription factor 1, EHT and matrix metalloproteinase 5. Gain of 17pq or 17q has been frequently reported in cases of GA, and candidate genes of relevance to gastric carcinogenesis are *HER-2/neu*, and those encoding survivin and gastrin.

Volik, Stanislav

[57]

A 20q13.2 cancer amplicon: novel sequence analysis tools

Stanislav Volik, Joe Gray, David Kowbel & Colin Collins

University of California San Francisco Cancer Center, San Francisco, California

Progress in mammalian genome sequencing demands development of efficient, high-throughput methods for functional annotation of megabase-scale regions. We report the first complete sequence of a tumor amplicon that is associated with aggressive tumor behavior and occurs in a wide range of tumors. The overwhelming amount of data generated for this region by traditional analysis tools (such as GenScan, BLAST and RepeatMasker) required development of efficient annotation methods. For a comprehensive biological annotation of the amplicon we developed Genome Cryptographer. This tool integrates information on distribution and type of repetitive elements, results of database searches, comparative sequence data, expression data, array comparative genomic hybridization copy

number data and information on CpG dinucleotide distribution. Genome Cryptographer revealed an uneven distribution of genes and repetitive elements in the 1-megabase region. The proximal amplicon peak as defined by array comparative genomic hybridization is composed of 60% repetitive sequence and encodes one protein-coding gene, *ZNF217*. A 14-kb duplication was identified close to the 5' end of *ZNF217*; it is 97% homologous to a 14-kb element on chromosome 22q13; contains one of three identified CpG islands, a conserved putative RNA gene and a single repetitive element; and is present at the murine *ZNF217* locus. The distal amplicon peak contains two genes, *CYP24* and *PFDN4*. Breakpoints for both amplicon peaks are located in regions of local repeat maxima. We have also developed software to create digital expression profiles of multi-megabase regions. This allowed us to confirm that *ZNF217* has the highest relative expression in tumors and to identify three additional putative cancer-related genes.

Watson, Mark

[58]

Expression profiling of ductal carcinoma *in situ* by laser capture microdissection and high-density oligonucleotide arrays

Veronica Luzzi, Victoria Holtschlag & Mark Watson

Department of Pathology and Immunology and Alvin J. Siteman Cancer Center, Washington University School of Medicine, St. Louis, Missouri 63110, USA

Gene expression profiling with nucleic acid arrays is a powerful method for the molecular classification of human neoplasms. Laser capture microdissection is an equally useful technique to isolate defined cell populations selectively from heterogeneous histological tissue sections. We demonstrate how a modest use of laser capture microdissection is sufficient to isolate nanogram quantities of high-quality RNA. Together with several internal standards and microcapillary electrophoresis of input RNA, we used two rounds of linear molecular amplification to generate sufficient quantities of labeled target for hybridization to high-density oligonucleotide expression arrays. Our results demonstrate that the technique is reproducible, generates only modest biasing of the original transcript population and offers sensitivity comparable to that achieved using standard methodology. Using this approach, we have compared the expression profiles of nonmalignant human breast epithelium and adjacent ductal carcinoma *in situ* lesions from breast cancer patients. Several genes, including those previously implicated in human breast cancer progression, demonstrate differential expression among the microdissected cell populations. This study demonstrates the feasibility of applying laser capture microdissection and nucleic acid expression array technology to histologically complex clinical specimens to classify human breast cancer molecularly and gain a better understanding of its molecular pathobiology.

Watts, George

[59]

Microarray analysis of a human multiple myeloma cell line selected with melphalan or doxorubicin identifies a set of commonly affected genes

George Watts, Denise Tye, Mary Gleason-Guzman & Bernard Futscher

Arizona Cancer Center, Tucson, Arizona 85724, USA

Multiple myeloma is a monoclonal plasma cell neoplasm that is frequently treated with melphalan and doxorubicin. The development of resistance to drugs such as melphalan and doxorubicin poses a hurdle to their successful clinical use.

Although the cytotoxic mechanisms of the two drugs differ, both can trigger apoptosis as the result of their cytotoxic actions. We proposed that selection for resistance to either melphalan or doxorubicin would lead to defects in apoptotic signaling and execution. We used complementary DNA microarrays to evaluate the relative expression of approximately 5,700 human genes in a human multiple myeloma cell line (8226/S) and its melphalan-resistant subline (8226/LR5). We then compared the results to those from a previous study of doxorubicin resistance. Overall, 1–2% of the 5,700 genes analyzed were differentially expressed by a factor of two or more. Several genes were found to have altered expression resulting from selection with either doxorubicin or melphalan. Genes affected by selection with either drug included those encoding N1 spermine/spermidine acetyltransferase, hydroxysteroid (11- β) dehydrogenase, paraoxonase and TNF receptor-associated factor 3. In addition, melphalan selection altered the same pathways as doxorubicin by affecting the expression of genes involved in apoptosis, such as those for Fas/CD95 and the anti-apoptotic BCL2-related protein A1. We conclude that melphalan and doxorubicin selection affected a common set of genes despite their differing mechanisms of action. These observations support apoptosis as a common endpoint for the cytotoxic actions of many chemotherapeutic drugs and indicate some of the points of the apoptotic process that can be altered by drug selection.

Weisz, Alessandro

[60]

Gene expression monitoring in hormone-responsive human breast cancer cells during estrogen-induced cell cycle progression

Luigi Cicatiello¹, Angelo Facchiano², Raffaele Calogero³, Michele De Bortoli³, Francesco Bresciani¹ & Alessandro Weisz¹

¹Istituto Patologia generale e Oncologia, Seconda Università degli Studi di Napoli, Naples, Italy

²Centro di Ricerca Interdipartimentale di Scienze Computazionali e Biotecnologiche, Seconda Università degli Studi di Napoli, Naples, Italy

³Università di Torino, Turin, Italy

Estrogens are potent mitogens for a subset of their target tissues, including those in breast cancer, in which they promote growth by interacting with specific intracellular receptors (ER- α and - β) of the nuclear receptor superfamily of transcription factors. Definition of the gene expression patterns, or signatures, characteristic of estrogen in growth-responsive cells would allow identification of the gene networks that mediate the cellular responses to these hormones and the discovery of new hormone-responsive genes and gene products that could be used as markers in experimental and clinical settings. For this purpose, we performed an initial characterization of the gene responses of human breast cancer cells to estrogen by determining the gene expression signatures of these cells before and after hormonal stimulation. We used microarrays containing more than 9,000 unique gene probes, about half of them unnamed. About 40% of the genes analyzed, including close to 1,200 expressed sequence tags, were detectable in RNA extracted from human breast cancer cells. Furthermore, the concentration of 10% messenger RNAs was either increased or decreased in the cell at one time or another after hormonal stimulation, each such change corresponding to a specific cell cycle phase. These included transcripts from known estrogen-responsive genes (such as *pS2*, *c-fos*, *c-myc*, and *cyclin D1*) as well as a significant number of new hormonal targets. Hierarchical clustering shows distinct patterns of gene activation and inhibition, which will be exploited to further investigate intracellular signaling by steroid hormones in human breast cancer cells.

Westbrook, Carol

[61]

A cDNA data set for expression studies of hematopoietic stem cells: characterization and database mining

Carol Westbrook, Ignatius Gomes, Tiffany Le, Jeffrey Kapp & Ronald Hoffman

University of Illinois, Chicago, Illinois

Bone marrow cells that express the CD34+ antigen (CD34+ cells) represent a continuum of early and committed progenitor cells, including the most primitive stem cells. We used gene filter arrays to develop a complementary DNA database that is representative of the CD34+ transcriptome, by comparing expression between human and baboon CD34+ cells to identify the most conserved and highly abundant genes in both species. RNA-based probes were sequentially hybridized to GeneFilters (Research Genetics) containing 25,920 human UniGene cDNAs. The human cells displayed 15,970 (62%) of the cDNAs on the filters, of which 97% varied less than threefold in expression level between species and 60% were expressed sequence tags. Polymerase chain reaction analysis with reverse transcription confirmed that selected genes were expressed at the predicted levels. To demonstrate the utility of this stem cell database for gene discovery, we mined it for transcriptional regulators by searching UniGene cluster descriptors for text strings corresponding to relevant protein motifs, including transcription factor, zinc-finger, POU, homeobox, helix-loop-helix and leucine zipper. By this method we selected 230 cDNAs, of which 186 were known genes, including 71 transcription factors, and the remaining 44 were expressed sequence tags. This preliminary description of transcriptional regulators expressed in CD34+ cells provides a potential wealth of new information about the molecular basis of normal hematopoietic cell proliferation and demonstrates the utility of the stem cell database for gene discovery. This database should prove useful for the preparation of a rationally designed cDNA array for expression analysis of normal hematopoiesis, stem cells and leukemias.

Williams, Kathleen

[62]

Patent strategies and patent disputes: SNPs, diseases, and treatment

Kathleen Williams

Palmer & Dodge, LLP, One Beacon Street, Boston, Massachusetts 02108, USA

Genome research and cancer treatment have presented challenges in defining what is meaningful within the large amounts of information generated in genomics and proteomics research. U.S. patent laws are in the process of clarifying what is patentable in genomics, gene expression, proteomics and bioinformatics. All these fields are the subject of intense patent activity, from patent disputes to the creation of patent strategies for protecting new technologies. What legal challenges might an investigator face as a discoverer of a technology that is key to disease treatment, whether it be a gene sequence, a mutation, a protein sequence or a method of dovetailing information on sequences and function? The law hands the researcher a sword to find legal avenues through patents and a shield to protect newly developed technologies. How can one effectively use these legal tools?