Bleomycin hydrolase is associated with risk of sporadic Alzheimer's disease

Susana E. Montoya *et al.*

Nature Genet. **18**, 211–212 (1998).

Due to a typographical error, the genotype frequencies for the ADRC confirmed cases in Table 1 were reported incorrectly as 0.426 instead of 0.490. The corrected table is shown on the right. We regret this error.

Table 1 • BH-PEN allele frequency and genotype distribution

	Total	Allele frequencies		Genotype frequencies		
		A	G	A/A	A/G	G/G
ADCR clinical cases	131	0.70 ± 0.03	0.30 ± 0.05	0.489	0.412	0.100
IADC confirmed cases	75	0.65 ± 0.05	0.35 ± 0.07	0.426	0.440	0.133
ADRC confirmed cases	151	0.67 ± 0.03	0.33 ± 0.05	0.490	0.358	0.152
AD cases ^a	357	0.68 ± 0.02	0.33 ± 0.03	0.476	0.398	0.127
IADC controls	57	0.69 ± 0.05	0.30 ± 0.08	0.456	0.474	0.070
ADRC controls	124	0.70 ± 0.04	0.30 ± 0.05	0.468	0.460	0.073
Pittsburgh controls ^b	139	0.73 ± 0.03	0.27 ± 0.05	0.511	0.432	0.058
AD controls ^a	320	0.71 ± 0.02	0.29 ± 0.04	0.484	0.450	0.066

^aPooled population. ^bRandom population sample from the Pittsburgh geographical area.