

# Evidence that positive selection drives Y-chromosome degeneration in *Drosophila miranda*

Doris Bachtrog

**Why does the Y chromosome harbor so few functional loci? Evolutionary theory predicts that Y chromosomes degenerate because they lack genetic recombination<sup>1</sup>. Both positive<sup>2</sup> and negative<sup>1</sup> selection models have been invoked to explain this degeneration, as both can result in the recurrent fixation of linked deleterious mutations on a nonrecombining Y chromosome. To distinguish between these models, I investigated patterns of nucleotide variability along 37 kb of the recently formed neo-Y chromosome in *Drosophila miranda*. Levels of nucleotide variability on this chromosome are 30 times lower than in highly recombining portions of the genome. Both positive and negative selection models can result in reduced variability levels, but their effects on the frequency spectrum of mutations differ. Using coalescent simulations, I show that the patterns of nucleotide variability on the neo-Y chromosome are unlikely under deleterious mutation models (including background selection and Muller's ratchet) but are expected under recent positive selection. These results implicate positive selection as an important force driving the degeneration of Y chromosomes; adaptation at a few loci, possibly increasing male fitness, occurs at the cost of most other genes on this chromosome.**

In humans and fruit flies, the X chromosome has several thousand genes, whereas its former homolog, the Y chromosome, contains only a few dozen<sup>3,4</sup>. This heteromorphism in gene content results from the loss of function of most genes on the Y chromosome. The degeneration of the Y is a consequence of its lack of sexual recombination (it is clonally transmitted from father to son)<sup>1</sup>. In nonrecombining genomes, selection at one site interferes with the action of selection at linked sites, reducing the efficacy of natural selection<sup>5</sup>. Two types of natural selection can interfere with the removal of deleterious mutations on a nonrecombining Y chromosome<sup>1</sup>. Models based solely on negative selection posit that the constant removal of deleterious mutations by purifying selection decreases the effective number of Y chromosomes. This, in turn, reduces the overall efficacy of purifying selection by increasing the influence of genetic drift<sup>1</sup>. Thus, deleterious mutations can accumulate on the Y solely as a consequence of

other deleterious mutations (the background selection and Muller's ratchet models<sup>1</sup>). An alternative view is that positive selection on the Y chromosome has a key role in its degeneration<sup>2</sup>. This model posits that recurrent fixations of strongly beneficial mutations (selective sweeps) drag along linked deleterious mutations.

Unfortunately, old Y chromosomes, such as those of humans, retain few traces of the processes that led to their degeneration. The best hope for distinguishing between positive and negative selection as causes of degeneration is to survey patterns of variability on young Y chromosomes that are still in the process of degeneration. Although both positive and negative selection models might account for reduced variability on an evolving Y chromosome<sup>1</sup>, the selective sweep model predicts a marked excess of low-frequency mutations relative to neutral expectations<sup>6,7</sup>, whereas negative selection models produce a less severe distortion in the frequency spectrum of mutations<sup>8-10</sup>.

**Table 1 Genomic regions investigated on the neo-Y chromosome**

Region	Total sites	Silent sites	Segregating sites
<i>CycB</i>	1,827	726	1
<i>dpn-1</i>	4,133	4,133	1
<i>dpn-2</i>	2,612	1,757	1
<i>dpn-3</i>	4,843	3,617	2
<i>eng</i>	361	346	0
<i>eve</i>	1,212	736	1
<i>exu1-1</i>	3,818	2,699	3
<i>exu1-2</i>	3,104	3,023	5
<i>exu1-3</i>	2,724	2,444	0
<i>exu1-4</i>	3,278	3,248	3
<i>mle-1</i>	1,655	1,655	1
<i>mle-2</i>	1,961	1,961	1
<i>robo-1</i>	1,766	798	0
<i>robo-2</i>	2,200	2,200	0
<i>zipper</i>	1,154	266	0
Total*	36,648	29,609	19

\*Nucleotide diversity is 0.012% per site based on the mean number of pairwise differences between alleles ( $\theta_w$ ) and 0.021% per site based on the numbers of segregating mutations ( $\theta_w$ ).

Institute of Cell, Animal and Population Biology, University of Edinburgh, Edinburgh EH9 3JT, UK; Department of Molecular Biology and Genetics, Cornell University, Ithaca, New York 14853, USA. Correspondence should be addressed to D.B. (Doris.Bachtrog@cornell.edu).

Published online 25 April 2004; doi:10.1038/ng1347

The newly formed neo-Y chromosome of *D. miranda* is at an intermediate stage of transition from an ordinary autosome into a degenerate Y chromosome<sup>11–13</sup>. Several types of deleterious mutations are accumulating on the neo-Y, including nonsense or frameshift mutations at some protein-coding genes<sup>11,13</sup>. In addition, there is a general increase in the rate of amino-acid substitution at neo-Y-linked genes<sup>14</sup> as well as an accumulation of repetitive and transposable element-derived DNA<sup>11,13,15</sup>. Thus, the *D. miranda* neo-Y provides a useful snapshot of the early stages of Y chromosome degeneration.

To determine what mode of selection is driving the accumulation of deleterious mutations on the neo-Y, I resequenced 37 kb in a population sample of 12 *D. miranda* neo-Y chromosomes. Levels of neutral variability on the neo-Y were much lower than in highly recombining regions (Table 1 and Supplementary Fig. 1 online). Diversity ( $\theta_{\text{Y}}$ ) on the neo-Y is 0.021% per silent site, ~30 times lower (or one-tenth of the expected neutral value correcting for relative numbers of X and Y chromosomes) than levels of diversity based on 15 neo-X- and X-linked genes (0.57% per site; refs. 12,16). This is in close agreement with a previous comparison of 5 kb of homologous regions on the neo-X and neo-Y chromosomes, which found a reduction in variability relative to the neutral expectation,  $\pi/\pi_0$ , of ~0.10 (95% confidence interval = 0.03–0.37; ref. 12).

Can both positive and negative selection models account for the observed reduction in variability on the neo-Y chromosome? The reduction in neutral variability due to deleterious mutations on the neo-Y depends on the number of Y chromosomes  $N_e$ , the rate of deleterious mutations  $U_d$  and their associated selection coefficient  $s_d$ <sup>8,10</sup>. The expected  $N_e$  for the neo-Y in the absence of selection can be estimated independently from average levels of nucleotide variability at other loci and is ~600,000. Relatively little is known about the rate of deleterious mutations and their associated selection coefficients<sup>17</sup>. To investigate whether deleterious mutations can, in principle, explain both the reduction in variability and the distortion in the frequency spectrum of segregating mutations, I allowed the two parameters  $U_d$  and  $s_d$  to vary over a wide grid of fixed values. The reduction in neutral variability ( $\pi/\pi_0$ ) due to deleterious mutations can be approximated by the equation

$$\frac{\pi}{\pi_0} = \frac{1}{N_e} \sum_{i,j=0}^m f_i f_j T_{ij}$$


in which  $T_{ij}$  is the mean time to the most recent common ancestor of two individuals with  $i$  and  $j$  deleterious mutations and  $f_i$  and  $f_j$  are the frequencies of individuals with  $i$  and  $j$  deleterious mutations given by a Poisson distribution with mean  $U_d/s_d$  (refs. 10,18). For a given value of  $N_e$ , I computed possible combinations of  $U_d$  and  $s_d$  that are consistent

with the observed reduction in variability on the neo-Y chromosome ( $0.03 \leq \pi/\pi_0 \leq 0.37$ ).


The shaded area in Figure 1a indicates parameter combinations of deleterious mutation models (combinations of  $U_d$  and  $s_d$  for a given  $N_e$ ) that are compatible with the observed reduction in diversity on the neo-Y. To the top right of Figure 1a are parameter combinations for which an equilibrium can be established between recurrent deleterious mutations and purifying selection (background selection<sup>1</sup>) are indicated. Under these parameters, all chromosomes in the population ultimately derive from the class of chromosomes free of deleterious mutations (the  $f_0$  class). The  $f_0$  class is stable under background selection, but as  $s_d$  becomes weaker (moving to the left in Fig. 1a), this class can be lost by random drift (a process called Muller's ratchet<sup>1</sup>). Parameter combinations for which Muller's ratchet is operating are indicated to the bottom left of Figure 1a.

Beneficial mutation models can also account for the observed reduction in diversity on the neo-Y (Fig. 1b). For a nonrecombining chromosome, such as the neo-Y, a selective sweep can be modeled as a population bottleneck to size one. Such a 'catastrophic sweep' causes all lineages in the population to coalesce suddenly and wipes out variability completely<sup>7,19</sup>. Under the catastrophic sweep model considered, the reduction in variability on the neo-Y depends solely on the time  $T_S$  since the last fixation of a beneficial mutation. I ran coalescent simulations to obtain values of  $T_S$  that are compatible with the observed diversity reduction. A recent selective sweep between 0.03 and 0.46  $N_e$  generations ago is consistent with the observed reduction in variability; a very recent selective sweep about 0.07  $N_e$  generations ago is most compatible with observed levels of diversity on the neo-Y (Fig. 1b).

Although both models can, in principle, explain the diversity reduction, their effects on the frequencies of linked neutral mutations differ.



**Figure 1** Parameter values for deleterious (a) and beneficial (b) selection models to explain the reduction in variability on the neo-Y chromosome of *D. miranda*. (a) Parameters for the deleterious mutation rate  $U_d$  and their associated selection coefficient  $s_d$  that are compatible with a reduction in diversity of 0.03–0.37. The dark gray area indicates values in the parameter space where the class of chromosomes containing the fewest number of deleterious mutations is stable (background selection), and the light gray area indicates parameters where Muller's ratchet is operating. (b) Estimates of the time since the last selective sweep  $T_S$  on the neo-Y chromosome that are compatible with the observed reduction in variability of the neo-Y chromosome (0.03–0.46  $N_e$  generations ago). Shown is the proportion of simulations that gave a value of  $|\theta_{\pi, \text{obs}} - \theta_{\pi, \text{sim}}| < \delta$ , where  $\theta_{\pi, \text{obs}}$  is 3.44 and  $\delta$  is 0.5.



**Figure 2** Mean value of Tajima's  $D$  under deleterious (a) and beneficial (b) selection models for a given reduction in diversity. (a) Each point shows the average value of Tajima's  $D$ , obtained from 1,000 coalescent simulations, for given parameter values of  $U_d$  and  $s_d$  (see Fig. 1). The dark gray points indicate parameters of background selection ( $N_e f_0 s_d > 10$ ); the white points indicate parameters of Muller's ratchet ( $N_e f_0 s_d < 1$ ); and the light gray points indicate parameters in between. (b) Mean value of Tajima's  $D$  for the positive selection model, based on 10,000 coalescent simulations, for different values of  $T_S$ .

To summarize the frequency spectrum of mutations, I used Tajima's  $D$  statistic<sup>20</sup>.  $D$  is expected to be close to zero in a constant-sized population with no selection (the standard neutral model) and negative when there is an excess of rare mutations. Most of the mutations (17 of 19) observed on the *D. miranda* neo-Y chromosome occur just once in the sample (Supplementary Fig. 1 online), resulting in a strongly negative value of  $D$  (−1.98). Such an excess of low-frequency variants is highly unlikely by chance ( $P < 0.01$ ) under the standard neutral model. I used coalescent simulations to generate samples under both deleterious and beneficial selection models to obtain empirical critical distributions for the test statistic.

Figure 2 shows the average values of  $D$  obtained from coalescent simulations for both positive and negative selection models. Deleterious mutation models can substantially reduce the mean value of  $D$  (Fig. 2a), with deleterious mutations of intermediate strength having the largest effect ( $N_e s_d \approx 10$ –100; Fig. 3). Despite reducing the mean value of  $D$  (as low as −1.3 for some combinations of  $U_d$  and  $s_d$ ), deleterious mutations rarely result in extreme values of  $D$ . For all combinations of  $U_d$  and  $s_d$ , at most 5% of the simulations gave as large a negative value of  $D$  as observed in the actual data (Fig. 4a). This means that each of these parameter combinations, despite being compatible with the observed diversity reduction on the neo-Y, can be rejected at the 5% significance level based on the observed value of  $D$ . Therefore, deleterious mutation models (background selection and Muller's ratchet) are unlikely to account for both the reduction in variability and the frequency spectrum of mutations on the neo-Y chromosome.


Selective sweeps on the neo-Y chromosome, in contrast, are expected to both reduce neutral variability and strongly distort the frequency distribution of neutral mutations<sup>7</sup>. The average value of  $D$  in selective sweep simulations was as low as −1.9 (Fig. 2b). The most likely value of  $T_S$  based on the reduction in variability is close to the most likely estimate based on the frequency distribution of mutations on the neo-Y chromosome (0.07 versus 0.05  $N_e$  generations; Figs. 1b and 4b). Unlike deleterious selection models, a large fraction of the selective sweep simulations (up to 40%) are compatible with the observed frequency distribution of mutations on the neo-Y chromosome (Fig. 4b). This suggests that a recent selective sweep shaped patterns of variability on the neo-Y chromosome of *D. miranda*.

Estimates of the rate and effect of deleterious mutations in *Drosophila* species suggest that beneficial mutations probably drag at least some deleterious alleles to fixation on the neo-Y of *D. miranda*<sup>21</sup>


(Supplementary Note online). In particular, a large number of weakly deleterious mutations can be dragged to fixation by a beneficial allele. For example, a single beneficial mutation of 1–2% selective advantage could drag more than 100 transposable elements ( $s_d = 10^{-4}$ ) to fixation, consistent with the observed accumulation and fixation of a large number of transposable elements on the neo-Y chromosome of *D. miranda*<sup>11,13,15</sup>.

What could cause a selective sweep on the neo-Y? Apart from recent suggestions of a high rate of beneficial mutations in *Drosophila* species, the Y chromosome is susceptible to certain types of adaptive mutations. Perhaps most importantly, the fact that the Y chromosome is male-limited may promote sex-specific adaptation. Because the Y is never transmitted through females, it is a haven for alleles that benefit males but harm females. The neo-Y may be experiencing recurrent selective sweeps acquiring such sexually antagonistic alleles<sup>22</sup>. A related hypothesis is that genes with sex-related function undergo adaptive specialization on newly formed X and Y chromosomes<sup>13</sup>. In particular, sex-related genes might evolve male-specific functions on the neo-Y<sup>13</sup>.

Is positive selection driving the degeneration of Y-chromosomes? Here, I showed that a recent selective sweep has occurred on the *D. miranda* neo-Y. Recurrent selective sweeps on the neo-Y could lead to considerable degeneration by fixing linked deleterious mutations<sup>2</sup>.



**Figure 3** Mean value of Tajima's  $D$  under the deleterious mutation model versus the strength of selection ( $N_e s_d$ ). Deleterious mutations of intermediate effect ( $N_e s_d = 10$ –100) cause the strongest distortion in the frequency spectrum of linked neutral mutations.



**Figure 4** Probability  $P$  of obtaining a Tajima's  $D$  equally or more negative in the simulations as observed from the data under deleterious (a) and beneficial (b) selection models. (a) For all parameter combinations of  $U_d$  and  $s_d$ , at most 5% of all simulations give as large a negative Tajima's  $D$  as observed in the data. (b) A wide range of  $T_S$  values for the selective sweep model are compatible with the Tajima's  $D$  of the data; up to 40% of the simulations, for a sweep 0.05  $N_e$  generation ago, give as large a negative Tajima's  $D$  value as observed on the neo-Y chromosome.

Unfortunately, on a nonrecombining chromosome, the current genealogy only contains information dating back to the last selective sweep event. Thus, the history of selection, both positive and negative, before this event cannot be inferred. In this regard, it will be of considerable interest to investigate Y and neo-Y chromosomes of other species. The approach presented here can be used to establish whether positive selection is frequent enough to have a general role in the process of Y chromosome degeneration. If so, Y chromosome degeneration may reflect adaptation at a few loci at the expense of most other genes on this chromosome.

## METHODS

**Data collection, *D. miranda* lines and DNA isolation.** I collected 36,648 bp of neo-Y-linked sequence per line (Table 1). The sequences of *CycB*, *robo-1*, *eve* and *eng* were published previously (5,166 bp in total<sup>12</sup>). In addition, I collected 31,482 bp of nucleotide sequence information, mostly from noncoding regions. For the newly collected data set, I isolated sequence data from five different genomic regions (containing the genes *mle*, *robo*, *dpm*, *exu1* and *zip*) from a genomic library constructed from *D. miranda*<sup>12</sup>. I used allele-specific primers to amplify the neo-Y-linked regions from male genomic DNA by PCR and then directly sequenced both strands using the ABI BigDye chemistry (Perkin-Elmer) on an ABI 377 automated sequencer. Details on primer sequences and sequencing strategy are available on request. The population survey used twelve *D. miranda* lines described elsewhere<sup>12</sup>. I isolated genomic DNA from individual male flies using the PUREGENE DNA extraction kit (Gentra). Because there is no recombination on the neo-Y chromosome (all loci share a common genealogy), data can be pooled among the neo-Y regions analyzed.

**Deleterious mutation models in the coalescent framework.** To model the dynamics of deleterious mutations in a statistical framework, I used the structured coalescent<sup>8,10,18,23</sup>. A Wright-Fisher model of a randomly mating, haploid population of  $N_e$  breeding individuals is considered. Alleles are classified as coming from different genetic backgrounds, defined by the number of deleterious mutations they carry. The basic model considered here assumes that all deleterious mutations have an identical selection coefficient,  $s_d$ . The simulation algorithm of the structured coalescent used was described in detail previously<sup>8,10</sup>. The first step in the algorithm is to generate a transition matrix  $Q_{ij}$  of the probabilities that a chromosome with  $i$  deleterious mutations in a given generation has an ancestor with  $j$  deleterious mutations in the previous generation<sup>8</sup>. Next, samples of chromosomes are generated randomly from the equilibrium distribution of deleterious mutations<sup>24</sup>. After this preliminary step, the genealogy for a sample of 12 chromosomes with no recombination was generated in a population subject to deleterious mutations. For each generation, the number of deleterious mutations in the ancestor of each individual is obtained randomly by using the probabilities in the transition matrix  $Q_{ij}$  as expected values. Coalescence is allowed to occur between individuals having the same number of deleterious mutations with probability  $k_i(k_i-1)/(N_e f_i)$ , where  $k_i$  is the number of lineages with  $i$  deleterious mutations present in the sample at a given generation and  $N_e f_i$  is the deterministic equilibrium size of class  $i$  after selection<sup>10</sup>. Once the most recent common ancestor of the sample is reached, neutral mutations are distributed over the tree by sampling from a Poisson distribution with mean  $(\pi_0 t/2)$ , where  $t$  is the length of each branch in units of  $2 N_e$  generations. I investigated 29,609 neutral sites on the neo-Y chromosome (Table 1); this corresponds to a neutral mutation rate of  $4.4 \times 10^{-5}$  per generation for the region investigated (29,609 sites ( $1.5 \times 10^{-9}$  per site per generation)) or a neutral  $\pi_0$  ( $N_e$ ) of 53.3 for the neo-Y chromosome.

The average estimate of  $\theta_x$  ( $3N_e$ ) for a total of 15 X-linked and neo-X-linked loci in *D. miranda* is 0.57% per site<sup>12,16</sup>. Assuming a neutral mutation rate of  $1.5 \times 10^{-9}$  per generation, the effective number of breeding individuals  $N$  is estimated to be  $\sim 1.2$  million, corresponding to an effective population size  $N_e$  of the neo-Y of  $\sim 600,000$ . Although this estimate is somewhat crude, the reduction in variability is almost independent of population size for strongly deleterious mutations (background selection<sup>8</sup>) and depends only slightly on the effective population size for weakly selected mutations<sup>10</sup>. Sexual selection in males can reduce the population size of the neo-Y relative to the neo-X chromosome. I investigated possible effects of sexual selection using the maximum value by which variation on the neo-Y can be reduced (one-ninth) or using the most plausible value of sexual selection in *Drosophila* populations (one-fifth; Supplemental Methods online). Ignoring the effects of sexual selection does not affect any of the conclusions drawn (Supplemental Figs. 2–7 online).

I ran 1,000 coalescent simulations for each parameter combination of  $U_d$  and  $s_d$  (a total of 388 combinations) and a population size of 600,000 neo-Y chromosomes. I considered the entire range of possible values of  $U_d$  and  $s_d$  that could explain the reduction in variability; mutations with a smaller  $s_d$  are nearly neutral ( $N_e s_d = 1.2$ ) and do not cause sufficient reduction in  $N_e$ , whereas stronger mutations ( $s_d > 0.1$ ) would require biologically unreasonable mutation rates to account for the reduction in diversity (and would not distort the frequency distribution of neutral mutations).

**Positive selection model in the coalescent.** I investigated the likelihood of the data under a catastrophic selective sweep by varying  $T_S$ , the time since the last selective sweep that is compatible with the observed reduction in variability on the neo-Y chromosome. I carried out coalescent simulations for a sample of 12 chromosomes according to the standard neutral model with constant population size<sup>23</sup> until the accumulated time is  $> T_S$ ; at this time point all lineages were forced to coalesce. This truncates the genealogy at the time of the sweep, resulting in a star-shaped genealogy. This method assumes that selection is strong relative to mutation, such that no variant arising during the process of fixation of the selected allele is likely to be sampled<sup>7</sup>. Neutral mutations are placed on the tree by sampling from a Poisson distribution with mean  $(\pi_0 t/2)$ . Values of  $T_S$  were varied over a range of points spaced every 0.01  $N_e$  generations; 10,000 coalescence simulations were run for each value of  $T_S$ .

**Summary statistics and hypothesis testing.** I used Tajima's  $D^{20}$  to summarize the frequency spectrum of mutations.  $D$  considers the normalized difference

between two estimates of the population mutation rate ( $\theta_D$ , based on the mean numbers of pairwise differences between alleles, and  $\theta_W$ , based on the numbers of segregating sites) and is expected to be negative when there is an excess of rare mutations. For both deleterious and beneficial selection models, I obtained empirical cut-off points for the test statistics to obtain the 5% significance level. For each value of  $U_d$  and  $s_d$  (deleterious mutation models) or each value of  $T_S$  (positive selection models), a  $P$  value for the data was estimated as the proportion of simulated samples with an identical or more extreme average value of the Tajima's  $D$  statistics than observed in the data. The average Tajima's  $D$  over several neo-X- and X-linked loci is close to zero<sup>13</sup>; thus, demographic explanations (for example, a recent population expansion in *D. miranda*) cannot explain the strong distortion in the frequency spectrum on the neo-Y.

Note: Supplementary information is available on the Nature Genetics website.

#### ACKNOWLEDGMENTS

I thank P. Andolfatto, N. Barton, B. Charlesworth, I. Gordo, M. Przeworski and S. Wright for discussion and comments on the manuscript. This work was supported by an Austrian Academy of Science postdoctoral fellowship, a Royal Society research grant to D.B. and a grant to B. Charlesworth from the British Biotechnology and Science Research Council.

#### COMPETING INTERESTS STATEMENT

The author declares that she has no competing financial interests.

Received 24 October 2003; accepted 17 March 2004

Published online at <http://www.nature.com/naturegenetics/>

- Charlesworth, B. & Charlesworth, D. The degeneration of Y chromosomes. *Philos. Trans. R. Soc. Lond. B Biol. Sci.* **355**, 1563–1572 (2000).
- Rice, W.R. Genetic hitchhiking and the evolution of reduced genetic activity of the Y sex chromosome. *Genetics* **116**, 161–167 (1987).
- Skaletsky, H. *et al.* The male-specific region of the human Y chromosome is a mosaic of discrete sequence classes. *Nature* **423**, 825–837 (2003).
- Carvalho, A.B. Origin and evolution of the *Drosophila* Y chromosome. *Curr. Opin. Genet. Dev.* **12**, 664–668 (2002).
- Hill, W.G. & Robertson, A. The effect of linkage on limits to artificial selection. *Genet. Res.* **8**, 269–294 (1966).
- Andolfatto, P. & Przeworski, M. More rare variants in regions of lower crossing over in African *Drosophila melanogaster*. *Genetics* **158**, 657–665 (2001).
- Simonsen, K.L., Churchill, G.A. & Aquadro, C.F. Properties of statistical tests of neutrality for DNA polymorphism data. *Genetics* **141**, 413–429 (1995).
- Charlesworth, D., Charlesworth, B. & Morgan, M.T. The pattern of neutral molecular variation under the background selection model. *Genetics* **141**, 1619–1632 (1995).
- Tachida, H. DNA evolution under weak selection. *Gene* **261**, 3–9 (2000).
- Gordo, I., Navarro, A. & Charlesworth, B. Muller's ratchet and the pattern of variation at a neutral locus. *Genetics* **161**, 835–848 (2002).
- Steinemann, M. & Steinemann, S. Enigma of Y chromosome degeneration: neo-Y and neo-X chromosomes of *Drosophila miranda* a model for sex chromosome evolution. *Genetica* **102–103**, 409–420 (1998).
- Bachtrog, D. & Charlesworth, B. Reduced adaptation of a non-recombining neo-Y chromosome. *Nature* **416**, 323–326 (2002).
- Bachtrog, D. Adaptation shapes patterns of genome evolution in sexual and asexual genomes in *Drosophila*. *Nat. Genet.* **34**, 215–219 (2003).
- Bachtrog, D. Protein evolution and codon usage bias on the neo-sex chromosomes of *Drosophila miranda*. *Genetics* **165**, 1221–1232 (2003).
- Bachtrog, D. Accumulation of *spock* and *worf*, two novel non-LTR retrotransposons on the neo-Y chromosome of *Drosophila miranda*. *Mol. Biol. Evol.* **20**, 173–181 (2003).
- Yi, S., Bachtrog, D. & Charlesworth, B. A survey of chromosomal and nucleotide sequence variation in *Drosophila miranda*. *Genetics* **164**, 1369–1381 (2003).
- Drake, J.W. *et al.* Rates of spontaneous mutation. *Genetics* **148**, 1667–1686 (1998).
- Hudson, R.R. & Kaplan, N.L. Gene trees with background selection. in *Non-neutral Evolution: Theories and Molecular Data* 140–153 (Chapman and Hall, New York, 1994).
- Maynard Smith, J. & Haigh, J. The hitch-hiking effect of a favourable gene. *Genet. Res.* **23**, 23–35 (1974).
- Tajima, F. Statistical method for testing the neutral mutation hypothesis by DNA polymorphism. *Genetics* **123**, 585–595 (1989).
- Mukai, T. *et al.* Mutation rate and dominance of genes affecting viability in *Drosophila melanogaster*. *Genetics* **72**, 335–355 (1972).
- Rice, W.R. Male fitness increases when females are eliminated from gene pool: Implications for the Y chromosome. *Proc. Natl. Acad. Sci. USA* **95**, 6217–6221 (1998).
- Hudson, R.R. Gene genealogies and the coalescent process. in *Oxford Surveys in Evolutionary Biology* 1–44 (Oxford University Press, Oxford, 1990).
- Haigh, J. The accumulation of deleterious genes in a population—Muller's Ratchet. *Theor. Popul. Biol.* **14**, 251–267 (1978).