

**Qing et al. reply**—The conclusion by Qiu et al. that HEp-2 and A431 cells do not express FCGR is wrong. The Muggeridge et al. report<sup>7</sup> they quote clearly shows that the FCGR number per cell is approximately 300.

There are few details of how Qiu et al. generated the data presented in their table. The remarkably high multiplicity of infection used in these experiments is not standard, and it is difficult to reconcile their 60% transduction rate for HeLa cells when others have reported that AAV vectors do not transduce these cells well because of the rate-limiting viral second-strand DNA synthesis<sup>11,12</sup>. Transduction efficiencies of 40% for HEp-2 and 10% for A431, respectively, are cited as proof that these cells can be transduced in the absence of FCGR expression. Yet, as stated above, these cells do indeed express FCGR (ref. 7). Thus, it seems that the analysis of FCGR by Qiu et al. using flow cytometry with a monoclonal antibody is inadequate to draw such a conclusion.


We have compared the transduction efficiency of a recombinant AAV-*lacZ* vector ( $4 \times 10^3$  particles/cell) and found transduction efficiencies in HeLa and 293 cells of approximately 4% and 20%, respectively, and <1% in A431 cells which are known to efficiently bind AAV (ref. 13). The lack of transgene expression in A431 cells has previously been reported to be due to very high levels of expression of the epidermal growth factor receptor (EGFR) protein tyrosine kinase known to limit the viral second-strand DNA synthesis<sup>13</sup>. The observed lack of transduction of M07e cells, which we showed do express FCGR (ref. 1), has previously been shown to be due to lack of expression of heparan sulfate proteoglycan<sup>14</sup> (HSPG), a co-receptor of AAV. The absolute requirement for the deliberate expression of both HSPG and FCGR1 in Raji cells, which are known to lack expression of both of these genes<sup>15</sup>, to render these cells permissive for AAV infection, strongly supports our contention that both HSPG and FCGR1 serve as co-receptors for AAV. Of course, other co-receptors may be used in other cells.

KEYUN QING, JONATHAN HANSEN, AND ARUN SRIVASTAVA

Department of Microbiology and Immunology  
Medical Science Building 247  
Indiana University School of Medicine  
635 Barnhill Drive  
Indianapolis, Indiana 46202-5120, USA

**Samulski et al. replies**—Using the procedure that Mizukami et al. used<sup>8</sup>, we also observed a 150-kDa protein (Fig. 1, lane 1); however,

Fig. 1 Viral overlay analysis of two different membrane preparations. Lanes 1 & 2 represent an AAV-2 overlay on equivalent amounts of membrane purified by the Chong & Rose method and the Hennache & Boulanger method respectively. Arrows point to the 150 kDa (lane 1) and 100 kDa (lane 2) proteins that interact with AAV.



this method (described by Chong and Rose<sup>16</sup>) does not stringently purify plasma membrane proteins. In our paper<sup>2</sup>, we used a method that specifically enriched cell surface proteins by 30-fold (ref. 17), as assessed by 5'-nucleotidase activity. In these more stringent conditions, binding to the 150-kDa protein was not detected. (Fig. 1, lane 2), thus our submitted gel<sup>2</sup> was truncated to save space. This protein may be a non-plasma membrane protein (for example, nucleolin as identified by Qiu and Brown), or a cell surface protein that migrates in a different fraction with our procedure. As the 'fold' enrichment of plasma membrane proteins was not monitored in Mizukami's study<sup>8</sup>, all interpretations are plausible.

As for  $\beta 5$  integrin, we also did not see interaction with the purified form, possibly because of the absence of essential post-translational modification. It should be noted that we observed AAV binding to immunoprecipitated  $\beta 5$  integrin, supporting the specificity of this interaction. Furthermore, we established that there is a role for integrin in AAV-2 infection (ref. 2, Figs. 2 and 3). The presence of integrin influences viral infection, but is not essential, as is the case with adenovirus<sup>10,18</sup>. Figure 3 of our study<sup>2</sup> clearly demonstrates that expression of  $\beta 5$  substantially increases AAV-2 internalization in a time-dependent manner, indicating a role in AAV entry, which may have important consequences *in vivo*<sup>2,10,18</sup>.

As for the transduction data, the 260% enhancement we observed is very similar to that seen for adenovirus (320%), whose use of  $\alpha V\beta 5$  integrin as a co-receptor is well established. In addition, it is not surprising that AAV may interact with integrin in a non-RGD manner. A ligand does not have to use an RGD or RGD-like motif in order to interact with integrin.

Integrin  $\alpha V\beta 3$  and  $\alpha V\beta 5$  facilitate adenovirus infection; however, it is  $\alpha V\beta 5$  inte-

grin that has been shown to have a dual role in facilitating both membrane permeabilization and internalization<sup>17</sup>. In addition, compared with  $\alpha V\beta 3$  integrin,  $\alpha V\beta 5$  internalizes adenovirus at a faster rate and renders cells significantly more susceptible to infection<sup>18</sup>. These studies and our data strongly suggest that both Ad and AAV use  $\alpha V\beta 5$  as a co-receptor to mediate viral entry.

RICHARD JUDE SAMULSKI, JEFFREY S. BARTLETT & CANDANCE SUMMERFORD  
Gene Therapy Center  
University of North Carolina at Chapel Hill  
7119 Thurston-Bowles CB7352  
Chapel Hill, NC 27599

1. Qing, K.Y. et al. Human fibroblast growth factor receptor 1 is a co-receptor for infection by adeno-associated virus 2. *Nature Med.* 5, 71-77 (1999).
2. Summerford, C., Bartlett, J.S. & Samulski, R.J.  $\alpha V\beta 5$  integrin: a co-receptor for adeno-associated virus type 2 infection. *Nature Med.* 5, 78-81 (1999).
3. Coughlin, S.R., Barr, P.J., Cousens, L.S., Fretto, L.J. & Williams, L.T. Acidic and basic fibroblast growth factors stimulate tyrosine kinase activity *in vivo*. *J. Biol. Chem.* 263, 988-993 (1988).
4. Yayon, A., Klagsbrun, M., Esko, J.D., Leder, P. & Ornitz, J.D. Cell surface, heparin-like molecules are required for binding of basic fibroblast growth factor to its high affinity receptor. *Cell* 64, 841-848 (1991).
5. WuDunn, D. & Spear, P.G. Initial interaction of herpes simplex virus with cells is binding to heparan sulfate. *J. Virol.* 63, 52-58 (1989).
6. Kaner, R.J. et al. Fibroblast growth factor receptor is a portal of cellular entry for herpes simplex virus type 1. *Science* 248, 1410-1413 (1990).
7. Muggeridge et al. Herpes simplex virus infection can occur without involvement of the fibroblast growth factor receptor. *J. Virol.* 66, 824-830 (1992).
8. Mizukami, H., Young, N. & Brown, K.E. Adeno-associated virus type 2 binds to a 150-kilodalton cell membrane glycoprotein. *Virology* 217, 124-130 (1996).
9. Qiu, J. & Brown, K.E. *Virology* (in the press).
10. Wicham, T.J., Mathias, P., Cheresch, D.A. & Nemerow, G.R. Integrins  $\alpha V\beta 3$  and  $\alpha V\beta 5$  promote adenovirus internalization but not virus attachment. *Cell* 73, 309-319 (1993).
11. Fisher, K.J. et al. Transduction with recombinant adeno-associated virus for gene therapy is limited by leading strand synthesis. *J. Virol.* 70, 520-532 (1996).
12. Ferrari, F.K., Samulski, T., Shenk, T. & Samulski, R.J. Second-strand synthesis is a rate limiting step for efficient transduction by recombinant adeno-associated virus vectors. *J. Virol.* 70, 3227-3234 (1996).
13. Mah, C. et al. Adeno-associated virus 2-mediated gene transfer: Role of epidermal growth factor receptor protein tyrosine kinase in transgene expression. *J. Virol.* 72, 9835-9843 (1998).
14. Bartlett, J.S. & Samulski, R.J. Fluorescent viral vectors: A new technique for the pharmacological analysis of gene therapy. *Nature Med.* 4, 635-637 (1998).
15. Kiefer, M.C. et al. Ligand-affinity cloning and structure of a cell surface heparansulfate proteoglycan that binds basic fibroblast growth factor. *Proc. Natl. Acad. Sci. USA* 87, 6985-6989 (1990).
16. Chong, L.D. & Rose, J.K. Membrane association of functional vesicular stomatitis virus matrix protein *in vivo*. *J. Virol.* 67, 407-414 (1993).
17. Hennache, B. & Boulanger, P. Biochemical study of KB-cell receptor for Adenovirus. *Biochem. J.* 166, 237-247 (1977).
18. Wicham, T.J., Filardo, E.J., Cheresch, D.A. & Nemerow, G.R. Integrin  $\alpha V\beta 5$  selectively promotes adenovirus mediated cell membrane permeabilization. *J. Cell Biol.* 127, 257-264 (1994).