PROTEIN BIOCHEMISTRY

Paving the path to single-molecule structures

A new generation of brilliant X-ray laser sources will be coming online within the next few years. Researchers now show that using these lasers to determine the structures of single molecules should be possible.

Obtaining the atomic structure of a protein by X-ray diffraction first requires purifying and crystallizing it. Although structural genomics efforts have laid pipelines for streamlining this tedious process, it still takes a great deal of time and effort, and many protein classes, such as membrane proteins, are still intractable to structure determination. A technology allowing researchers to obtain atomic structures of single protein molecules would eliminate the need for tiresome purification and crystallization, and would arguably have an enormous impact on structural biology.

Such a potentially revolutionary technology is not just a pipe dream. A new generation of X-ray sources known as X-ray free-electron lasers (XFELs) is being developed, and the first source, at the Stanford Linear Accelerator Center in the United States, is

The structure of the protein chignolin at 1.8 Å resolution, recovered from 72,000 simulated diffraction snapshots of unknown orientation. Reprinted from *Nature Physics*.

expected to be up and running as early as mid-2009. Researchers believe that these brilliant lasers, which come with up to a billion-dollar price tag, will facilitate single-molecule structure determination. “We think that these X-ray lasers are going to transform structure determination in the way that lasers transformed spectroscopy about 40 to 50 years ago,” says Abbas Ourmazd, a physicist at the University of Wisconsin, Milwaukee.

However, single-molecule structure determination will be a bit more complicated than just putting the molecule in the path of a powerful XFEL beam. The scattering of an X-ray signal by a single molecule is extremely faint, so many diffraction ‘snapshots’ must be collected to boost the signal. “Estimates are that one of these XFELs will deliver somewhere between 100 and 1,000 terabits of data per day, most of which will be zeros because single objects don’t scatter very many photons,” explains Ourmazd. Thus a major question has been how to best use this sparse yet huge amount of data to reconstruct the three-dimensional structure of a single molecule.

Ourmazd and his co-workers found that the ideas proposed by other researchers to use XFEL data to solve single-molecule structures fell orders of magnitudes short. His group now has demonstrated, by applying algorithms that exploit the correlations between photons in a large series of X-ray scattering snapshots of a single object in random orientations, that they can recover a high-resolution atomic structure.

PROTEOMICS

GLOBAL INSTABILITY

Scientists create a high-throughput platform for proteome-scale assessment of protein stability.

Proteins are the work horses of the cell, and fluctuations in their expression are fundamental for metabolism, cell-cycle control and communication between a cell and its environment. Although proteome-scale approaches are increasingly used to understand biological networks, the evaluation of dynamic protein stability has remained inaccessible at this scale.

Stephen Elledge and his colleagues at Brigham and Women’s Hospital and Harvard University Medical School now describe global protein stability (GPS) analysis, a high-throughput approach for assessing proteome-scale protein turnover in mammalian cells (Yen *et al.*, 2008). Elledge and his colleagues developed GPS to clear a major technical hurdle to identify substrates of the Skp1-cullin-F-box (SCF) ubiquitin ligase, cullin 1 (Cul1; Yen & Elledge, 2008). “We discovered this family of ubiquitin ligases called the SCF proteins,” he says. “It turned out that many of their substrates were very important regulatory molecules..., if we could only find them! The methods that were out there to identify them were fairly laborious, not easy to do and they weren’t general.”

The GPS method consists of a fluorescence-based multiplex system for assessing protein stability on a high-throughput scale. The researchers used a reporter construct encoding both the fluorescent protein DsRed and an enhanced GFP (EGFP)-X fusion from a bicistronic message, which allowed them to use the ratio of EGFP:DsRed as an indicator of the stability of protein X. They infected cells with a virus derived from a human cDNA library containing the fluorescent reporters and fractionated the cells by fluorescence-activated cell sorting (FACS) based on their EGFP:DsRed ratios. They determined the distribution of each open reading frame across the FACS fractions using a DNA microarray. Using this distribution they calculated a protein stability index for each protein.

The researchers demonstrated several key applications for GPS including the identification of Cul1 ubiquitin ligase substrates and a real-time evaluation of the stability of 8,000 human proteins. Elledge adds: “We can now look at stability and how it varies with position in the cell cycle, which was extremely laborious before. This is just a simple FACS analysis in an unperturbed population of cells.” But cell cycle is not the only parameter that can be assessed. “You could look

NEWS IN BRIEF

PROTEIN BIOCHEMISTRY

Directed evolution with 21 amino acids

Directed evolution via phage display is a well-established method to generate proteins with new or improved activities. Liu *et al.* now present a phage-display directed-evolution system with an expanded genetic code, by adding orthogonal tRNA–aminoacyl-tRNA synthetase pairs encoding unnatural amino acids to the *Escherichia coli* translational machinery. As proof of principle, they evolved a more effective antibody to gp120 containing the unnatural amino acid sulfotyrosine.

Liu, C.C. *et al. Proc. Natl. Acad. Sci. USA* **105**, 17688–17693 (2008).

BIOINFORMATICS

Software for ChIP

Transcriptional regulation can be studied on a genome-wide scale via chromatin immunoprecipitation (ChIP) followed by microarray (ChIP-chip) or sequencing (ChIP-seq) analysis. Ji *et al.* now present CisGenome software for managing and analyzing data from both ChIP-chip and ChIP-seq experiments. This all-inclusive software package supports data visualization, normalization and peak detection, computes false discovery rates and gene-peak associations, and can perform sequence and motif analysis.

Ji, H. *et al. Nat. Biotechnol.* **26**, 1293–1300 (2008).

CHEMICAL BIOLOGY

Caged quantum dots

Photoactivatable fluorophores, which are ‘turned on’ by light when the experimentalist chooses, are useful for imaging a variety of cellular processes. Han *et al.* now describe a method to synthesize photocaged quantum dots, which only become luminescent upon ultraviolet light stimulation and release of the chemical quenching group. Preliminary tests in mouse cells demonstrate their potential for a broad range of imaging applications as quantum dots are much brighter and more stable than organic fluorophores.

Han, G. *et al. J. Am. Chem. Soc.* **130**, 15811–15813 (2008).

GENOMICS

Mapping genome-wide protein-RNA interactions

Licalosi *et al.* describe an unbiased method, high-throughput sequencing of RNA isolated by crosslinking immunoprecipitation (HITS-CLIP), to comprehensively map protein-RNA interactions *in vivo*. HITS-CLIP uses ultraviolet light to induce cross-linking in RNA-protein complexes, allowing them to be immunopurified by stringent procedures. The RNA targets can then be identified by high-throughput sequencing. They used this method to identify Nova-RNA interactions in the mouse brain on a genome-wide scale.

Licalosi, D.D. *et al. Nature* **456**, 464–469 (2008).

BIOINFORMATICS

Predicting post-translational modifications

Mass spectrometry-based proteomics has brought to light a wealth of information about protein post-translational modification sites, but many, many more sites remain yet undiscovered. Schwartz *et al.* used the motif-x software tool to discover phosphorylation motifs in yeast, fly, mouse and human phosphorylation datasets. They also describe a newly developed companion program, scan-x, which can then predict unknown phosphorylation sites in these organisms.

Schwartz, D. *et al. Mol. Cell. Proteomics*; published online 28 October 2008.

Ourmazd likens their method to reconstructing an ancient Greek vase from a pile of shards by maximizing the correlations between the pottery fragments. For reconstructing a molecular structure from a series of diffraction snapshots, the scattered photon arrangements are changed until the correlations between them are maximized. “Our technique depends primarily not on the number of photons you scatter per shot, but rather on the total number of photons you’ve scattered from all possible orientations of the object,” says Ourmazd. Using this correlation approach, his group reconstructed a high-resolution structure of the small protein chignolin from 72,000 simulated diffraction snapshots of the protein in random orientations at the expected scattered photon intensity. “Remarkably, the only information one needs is the dimensionality of the space in which the object lives,” he says.

This approach should theoretically allow researchers to obtain single-molecule structures for proteins of various sizes and even large ensembles such as viruses or nanoparticles. For looking at objects larger than chignolin, “the only requirement is to have a large enough computer,” notes Ourmazd. Looking at single molecules rather than bulk ensembles may also provide a route to mapping conformational heterogeneity, which is all but impossible with current crystallographic methods.

Though it will likely take several years of working out technical issues even after the first XFELs come online, it will certainly be very exciting to see whether the promise of single-molecule structure determination can be realized. “The proof of the pudding is in the eating, and there’s nothing like taking real experimental data and showing everything works,” says Ourmazd. “We’re looking forward to that.”

Allison Doerr

RESEARCH PAPERS

Fung, R. *et al.* Structure from fleeting illumination of faint spinning objects in flight. *Nat. Phys.* advance online publication 9 November 2008.

at populations of cells and ask if stability varies with some cell-surface marker or whatever you’re interested in,” says Elledge. “All the other uses of fusion technology (chemical screens, RNA interference screens) can be adapted to look for things that affect the stability of your protein.”

Despite the exciting opportunities already offered by GPS, Elledge expects the platform to improve. “By cleaning up the way we look for substrates, enlarging the library and getting rid of the noise, I think that we’ll have it in shape in a couple of years where you can really use a single ligase and hope to find a substrate,” he says. Given the number of unexpected biological findings emerging from their early applications, Elledge hopes the surprises keep coming. “The way that we solved the problem allowed us to ask a lot of other questions that we were not initially planning on because we could look at the whole proteome at once and extract general information out of that,” he says, adding: “I suspect, given the fact that there are over 500 different ubiquitin ligases, that there are a lot of things going on in the cell that we don’t know about.”

Amy Donner

RESEARCH PAPERS

Yen, H.-C.S. *et al.* Global protein stability profiling in mammalian cells. *Science* **322**, 918–923 (2008).

Yen, H.-C.S. & Elledge, S.J. Identification of SCF ubiquitin ligase substrates by global protein stability profiling. *Science* **322**, 923–929 (2008).