

NEWS IN BRIEF

ger the linker, the better the receptors recycled. Even though this observation is intriguing and could shed light on basic trafficking mechanisms, the main application of these synthetic receptors is the transport of molecules into cells. This synthetic chemistry approach for accessing the cellular interior offers great versatility. Chemists may be able to engineer artificial receptors with different binding motifs that facilitate the uptake of virtually anything, to the benefit or destruction of cells. For example, in drug delivery, tumor cells carrying a specific synthetic receptor might be made more susceptible to toxic drugs.

Although such applications hold promise, there are still big hurdles to overcome. One is to show that these synthetic receptors, which have primarily been tested on cells in culture, also work in animals. Peterson is cautiously optimistic; his team has preliminary results from a vertebrate model in which the receptors are incorporated into cellular membranes. Whether they still work as receptors and how toxic they are in an animal over a longer time span is now being tested.

Nonetheless, this approach offers an innovative solution to an old problem. If Odysseus had met Blake Peterson, maybe the Greeks would have dismissed the idea of a horse and instead presented the Trojans with a city gate containing a clandestine portal.

Nicole Rusk

RESEARCH PAPERS

Boonyarattanakalin, S. *et al.* Synthetic mimics of small mammalian cell surface receptors. *J. Am. Chem. Soc.* **126**, 16379–16386 (2004).

can be utilized for detection,” says Seeberger. As the next step to this goal, the Seeberger lab is using the array technology to screen bacterial strains in an attempt to define a unique carbohydrate-binding ‘fingerprint’ for each strain. Intriguingly, if the strength of binding to specific carbohydrates is found to correlate with bacterial pathogenicity, the method could be used to quickly estimate the danger associated with a bacterial infection.

Detection of bacteria is just one possible use for this promising technology. Of more interest to researchers is the possibility of using carbohydrate arrays to examine carbohydrate-protein interactions. Carbohydrates are displayed on the surface of all cells and are involved in important cellular functions such as adhesion, recognition and signaling, in addition to being a route for bacterial infection. It has also been suggested that surface carbohydrates could serve as markers for cancer progression. Although they have yet to publish their results, Seeberger expects these new arrays to work on mammalian cells in addition to bacteria. These new technologies should make it much easier for researchers to exploit the many promising avenues of research embodied in carbohydrate mediated interactions.

Daniel Evanko

RESEARCH PAPERS

Disney, M.D. & Seeberger, P.H. The use of carbohydrate microarrays to study carbohydrate-cell interactions and to detect pathogens. *Chem. Biol.* **11**, 1701–1707 (2004).

Nimrichter, L. *et al.* Intact cell adhesion to glycan microarrays. *Glycobiology* **14**, 197–203 (2004).

GENE REGULATION**Substrate-induced gene expression screening of environmental metagenome libraries for isolation of catabolic genes**

Uchiyama *et al.* present a new technique for the identification of genes induced by a catabolite of interest: substrate-induced gene expression screening (SIGEX). Genomic fragments from an organism of interest are cloned into an operon-trap GFP expression vector, and FACS is used to separate clones in which the presence of a particular catabolite induces expression.

Uchiyama, T. *et al. Nat. Biotechnol.*, **23**, 88–93 (2005).

VIROLOGY**Synchronized infection of cell cultures by magnetically controlled virus**

Current strategies for the retroviral infection of cultured cells rely on the diffusion of viral particles. By coating lentivirus with iron oxide nanoparticles and using a magnetic field to draw the coated viruses to the cell monolayer, Haim *et al.* demonstrate a dramatically enhanced rate of association, suggesting the possibility of tightly synchronized studies of the infection process.

Haim, H. *et al. J. Virol.* **79**, 622–625 (2004).

IMAGING AND VISUALIZATION**Tumor imaging by proteolytic activation of cell-penetrating peptides**

Previous studies have demonstrated that relatively simple cell-penetrating peptides (CPPs) can facilitate passage through cell membranes by an associated ‘cargo’ molecule. Jiang *et al.* further enhance this technology, creating CPPs that rely on proteolytic cleavage for activation and demonstrating their capacity to label tumor cells.

Jiang, T. *et al. Proc. Natl. Acad. Sci. USA* **51**, 17867–17872 (2004).

DNA CLONING AND AMPLIFICATION**Accurate multiplex gene synthesis from programmable DNA microchips**

Oligonucleotide synthesis can be a costly and error-prone process. Tian *et al.* present a microarray-based strategy for the rapid and affordable synthesis, amplification and purification of oligonucleotides. They demonstrate the rapid production of highly accurate oligonucleotides and their application to the synthesis of a variety of full-length genes.

Tian, J. *et al. Nature* **432**, 1050–1054 (2004).

GENOMICS**Global identification of human transcribed sequences with genome tiling arrays**

Bertone *et al.* demonstrate the use of a human genome tiling array for the large-scale investigation of transcriptional activity. cDNA derived from liver mRNA was hybridized against arrays representing roughly 1.5 Gb of nonrepetitive genomic sequence, confirming numerous known and predicted genes, and revealing over 10,000 previously unidentified transcribed sequences.

Bertone, P. *et al. Science* **306**, 2242–2246 (2004).