Corrigendum: A microRNA switch regulates the rise in hypothalamic GnRH production before puberty

Andrea Messina, Fanny Langlet, Konstantina Chachlaki, Juan Roa, Sowmyalakshmi Rasika, Nathalie Jouy, Sarah Gallet, Francisco Gaytan, Jyoti Parkash, Manuel Tena-Sempere, Paolo Giacobini & Vincent Prevot

Nat. Neurosci. 19, 835–844 (2016); published online 2 May 2016; corrected after print 3 June 2016

In the version of this article initially published, the Figure 1e,f legend read, “Circulating levels of LH (left panels) and FSH (right panels) in GnRH cells of control (blue) and Dicer mutants (red)”; as the hormones were not measured in GnRH cells, it should have simply read “Circulating levels of LH (left panels) and FSH (right panels) in control (blue) and Dicer mutants (red).” Figure 2b was missing scale bars and has been replaced. The label “TSB-200” was missing from the rightmost bar in Figure 4d. And the treatment in Figure 5c was misidentified as TSB-200 instead of TSB-155. The errors have been corrected in the HTML and PDF versions of the article.

Corrigendum: Hedgehog signaling promotes basal progenitor expansion and the growth and folding of the neocortex

Lei Wang, Shirui Hou & Young-Goo Han

Nat. Neurosci. 19, 888–896 (2016); published online 23 May 2016; corrected after print 28 June 2016

In the version of this article initially published, the units on the *x* axis in Figure 3c were given as mm; the correct units are μm . At the end of the legend to Figure 7, the error bars were described as s.d.; they are actually s.e.m. in **b** and s.d. in **c**. In the third sentence of the Online Methods section on human cerebral organoids, 10% knockout serum replacement, 1% GlutaMAX and 1% MEM-NEAA should have been 20%, 1 $\times$ and 1 $\times$, respectively. In the sixth sentence, 1% N2 supplement, 1% GlutaMAX and 1% MEM-NEAA should each have been 1 $\times$. In the eighth sentence, 6-mm dishes should have been 6-cm dishes, 0.5% N2 supplement and 0.5% MEM-NEAA should each have been 0.5 $\times$, and 1% B27 without vitamin A, 1% GlutaMAX and 1% penicillin/streptomycin should each have been 1 $\times$. In Supplementary Figure 10b, the graph lacked error bars. The errors have been corrected in the HTML and PDF versions of the article.

Addendum: The schizophrenia risk gene product miR-137 alters presynaptic plasticity

Nat. Neurosci. 18, 1008–1016 (2015); published online 25 May 2015; addendum published after print 8 July 2016

Sandra Siegert, Jinsoo Seo, Ester J Kwon, Andrii Rudenko, Sukhee Cho, Wenyan Wang, Zachary Flood, Anthony J Martorell, Maria Ericsson, Alison E Mungenast & Li-Huei Tsai

It has come to our attention that our results, as presented, have caused some confusion among readers with regard to genetic variation at the *MIR137* locus, gene expression of *MIR137* and schizophrenia risk. It should be noted that it is currently unclear how gene variants at the *MIR137* locus confers disease risk and which are the causative or functional alleles or the gene(s) that mediate risk. Also, it is not clear to what extent a single disease-associated SNP, in isolation, relates to disease. In our study we did not examine the effects of single alleles since they often segregate together. It remains to be determined how the multiple reported disease-associated SNPs in the *MIR137* locus alone and in combination influence gene expression. To clarify this would require genome editing to alter SNPs, one by one, in cell lines. Finally, further studies are required to determine whether the synaptic and behavioral changes observed in this study can be linked to the specific genetic variants within the *MIR137* locus associated with schizophrenia in genome-wide association studies.