BRIEF COMMUNICATION OPEN

Spaceflight modulates gene expression in the whole blood of astronauts

Jennifer Barrila^{1,7}, C Mark Ott^{2,7}, Carly LeBlanc³, Satish K Mehta⁴, Aurélie Crabbé^{1,8}, Phillip Stafford⁵, Duane L Pierson² and Cheryl A Nickerson^{1,3,6}

Astronauts are exposed to a unique combination of stressors during spaceflight, which leads to alterations in their physiology and potentially increases their susceptibility to disease, including infectious diseases. To evaluate the potential impact of the spaceflight environment on the regulation of molecular pathways mediating cellular stress responses, we performed a first-of-its-kind pilot study to assess spaceflight-related gene-expression changes in the whole blood of astronauts. Using an array comprised of 234 well-characterized stress-response genes, we profiled transcriptomic changes in six astronauts (four men and two women) from blood preserved before and immediately following the spaceflight. Differentially regulated transcripts included those important for DNA repair, oxidative stress, and protein folding/degradation, including *HSP90AB1*, *HSP27*, *GPX1*, *XRCC1*, *BAG-1*, *HHR23A*, *FAP48*, and *C-FOS*. No gender-specific differences or relationship to number of missions flown was observed. This study provides a first assessment of transcriptomic changes occurring in the whole blood of astronauts in response to spaceflight.

npj Microgravity (2016) 2, 16039; doi:10.1038/npjmgrav.2016.39; published online 8 December 2016

During spaceflight, astronauts experience a variety of physical and psychological stressors, including acceleration at launch, isolation, confinement, high levels of ionizing radiation, sleep deprivation, and microgravity, which have been linked to a variety of physiological changes, including dampened immune function, bone and muscle loss, and increased viral reactivation.^{1–5} The molecular basis of spaceflight-induced responses remains unclear and may provide critical insights into health outcomes associated with space travel. Targeted spaceflight-associated changes in cytokine and antibody expression have been previously reported for astronauts,^{1–3} and more recently Terada *et al* reported the effects of spaceflight on hair follicle gene expression in astronauts.⁶ To evaluate the potential impact of the spaceflight environment on molecular pathways mediating cellular stress responses, we performed a first-of-its-kind pilot study to assess spaceflight-related changes in the expression of stress-response genes in the whole blood of astronauts in response to spaceflight.

Transcriptional profiling of 234 well-characterized stress-response genes was performed using total RNA isolated from whole blood obtained from six consenting astronauts 10 days before launch aboard the space shuttle and 2–3 h after return to Earth (see Supplemental Methods). Four men between the ages of 38 and 47 (mean = 43 years) and two women between the ages of 38 and 44 (mean = 41 years) participated in the study. Each of these astronauts flew aboard one of four space shuttle missions ranging between 10 and 13 days duration that took place during a 2-year period. Previously published studies using blood and saliva samples from these same crew members before, during and just after spaceflight indicated that all six astronauts displayed increases in Epstein Barr virus reactivation both during and

immediately post flight, which is consistent with these individuals being in a stressed state.^{4,5}

Microarray analysis revealed some variation in the gene-expression patterns displayed across individual crew members. This is not surprising given the smaller study size due to the exceptionally rare opportunity for this type of sample collection, as well as other studies that have shown that there can be a wide range of individual and temporal variability in gene-expression patterns in human blood.⁷ High variability was also observed in gene-expression studies using astronaut hair follicles.⁶ We found six transcripts that displayed significant ($P < 0.05$) changes in expression in the crew in response to spaceflight when compared with pre-flight levels (Table 1). Two additional differentially regulated transcripts were identified following outlier removal. No gender-specific differences or relationship to number of missions flown were observed. It is important to note that as for many human spaceflight studies, the small number of human subjects available for analysis is a limiting factor to the statistical power. False discovery rate algorithms determined no significant genes were expressed, likely due to these low study group numbers. The NASA Twins Study with astronauts Scott Kelly and Mark Kelly performed during the One-Year Mission may face similar challenges in their gene-expression data interpretation, as there is an n of 1 in each condition (spaceflight and ground). However, one advantage in the Twins Study over the present one is the longitudinal sample collection that could be performed during the year that Kelly spent in space. It will be intriguing to compare the results from gene-expression analyses between the two studies.

¹Center for Infectious Diseases and Vaccinology, The Biodesign Institute, Arizona State University, Tempe, AZ, USA; ²Biomedical Research and Environmental Sciences Division, NASA Johnson Space Center, Houston, TX, USA; ³Department of Microbiology and Immunology, Program in Molecular Pathogenesis and Immunity, Tulane University Health Sciences Center, New Orleans, LA, USA; ⁴EASI/Wyle Laboratories, Houston, TX, USA; ⁵Center for Innovations in Medicine, The Biodesign Institute, Arizona State University, Tempe, AZ, USA and ⁶School of Life Sciences, Arizona State University, Tempe, AZ, USA.

Correspondence: CA Nickerson (cheryl.nickerson@asu.edu)

⁷These authors contributed equally to this work.

⁸Current address: Laboratory of Pharmaceutical Microbiology, Ghent University, Ottergemsesteenweg 460, Ghent, Belgium

Received 18 July 2016; revised 16 September 2016; accepted 30 September 2016

Table 1. Differentially regulated genes in astronaut blood in response to spaceflight

Gene	Name/description	Ratio post/pre	P value
<i>Downregulated genes</i>			
<i>HSP27</i>	27 kDa heat-shock protein (HSP27); stress-responsive protein 27 (SRP27); estrogen-regulated 24 kDa protein; heat shock protein family B (small) member 1 (HSPB1)	0.44	0.045
<i>BAG-1</i>	BCL-2-binding athanogene-1; glucocorticoid receptor-associated protein RAP46	0.37	0.015
<i>XRCC1</i>	DNA repair protein – X-ray repair cross-complementing factor 1	0.45	0.044
<i>HSP90AB1</i>	90 kDa heat-shock protein- β ; 84 kDa heat-shock protein- β (HSP84); heat shock protein 90 class B (HSPCB)	0.41	0.023
<i>GPX1</i>	Glutathione peroxidase (GSHPX1; GPX1)	0.52	0.041
<i>HHR23A</i>	UV excision repair protein—human homolog of <i>Saccharomyces cerevisiae</i> RAD23A	0.53	0.003 ^a
<i>FAP48</i>	48 kDa FK506 Binding Protein (FKPB)-associated protein	0.55	0.022 ^b
<i>Upregulated genes</i>			
<i>C-FOS</i>	c-Fos proto-oncogene	3.11	0.021

Abbreviation: UV, ultraviolet.

^aAn outlier was identified for the pre-flight mean. Excluding this outlier yielded a significant *P* value. Inclusion of outlier yielded a *P* value of 0.105.^bAn outlier was identified for the pre-flight and post-flight means. Excluding outliers yielded a significant *P* value. Inclusion of outliers yielded a *P* value of 0.111.

Genes altered in expression encode proteins of known importance for DNA repair (*XRCC1* and *HHR23A*), oxidative stress (*GPX1*), and chaperones which have key roles in protein folding and/or proteasomal degradation (*HSP27* and *HSP90AB1*). The finding that genes encoding DNA-repair proteins are down-regulated supports previous studies which showed post-flight increases in chromosomal aberrations in lymphocytes of astronauts⁸ and indications of increased DNA damage in astronauts after long-duration spaceflight.⁹ Kumari *et al* previously reported that exposure of human lymphocytes to simulated microgravity over the course of 7 days led to increased DNA damage.¹⁰ This damage was accompanied by progressive decreases in the expression of a number of representative DNA-repair genes, leading the authors to postulate that impaired DNA-repair capacity could lead to increased damage and mutations. In a separate study, it was reported that 1 week of hindlimb unloading in male BALB/c mice led to the down-regulation of multiple DNA-repair genes including ERCC 1, ERCC 3, ERCC5, ERCC6, and XPC in the testes of BALB/c mice.¹¹

Of particular interest was the downregulation of *GPX1*, which encodes for glutathione peroxidase (GPX1), an enzyme that protects cells from oxidative damage, modulates the immune response (including delayed type hypersensitivity/DTH), and has been associated with increased viral titers in herpes viral infections.¹² Given that spaceflight depresses the DTH response,² alters cellular oxidative functions,^{9,13–17} and increases herpes viral reactivation in astronauts on orbit,^{4,5} additional detailed studies related to glutathione and oxidative stress may fundamentally advance our understanding of mechanisms underlying risks to crew health during a mission. Several previous studies have indicated the potential for alterations in oxidative stress and antioxidant defense status in response to spaceflight and spaceflight-analog models in humans and animals.^{9,13–16} Although the reported magnitude and direction of the expression/activity levels of specific redox enzymes (including GPX1) were variable across these studies (likely due to differences in experimental parameters such as the use of human versus animal subjects, mission duration, etc) the overall findings support the potential for increased oxidative stress and/or decreased antioxidant defense capacity in response to spaceflight. It is also interesting to note that GPX1 is a selenoprotein and decreased serum selenium levels have been reported in astronauts post landing as compared with pre-launch.⁹ Therefore, additional insight into GPX1 and other oxidative defense mechanisms may serve as a guiding principle for establishing nutritional requirements to ensure health safety

and performance of the crew during human exploration of space.^{9,18} When we compared these transcriptomic findings from blood samples to a previously published study using hair samples,⁶ we did not find similarities in the genes that were differentially regulated pre- and post flight, which could be due to a variety of experimental differences, including sample source (whole blood versus hair follicles) and mission duration (10–13 days in the present study compared with ~6 months in the study by Terada *et al*⁶).

To our knowledge, this study represents the first report of transcriptomic changes occurring in the blood of astronauts in response to spaceflight. Several transcripts encoding stress-response genes were suppressed in the crew after exposure to the microgravity environment, including those important for DNA repair, oxidative stress, detoxification, and protein folding/degradation. These processes are vital for maintaining human health by mediating cellular pathways that serve to protect against both environmental and physiological stressors and have been implicated in a broad spectrum of diseases. Since changes in gene expression in peripheral blood may be attributed to a number of factors, including changes in the distribution of blood cellular subsets (which has been detected post flight in the crew),^{1–3} future studies designed to establish direct links between specific physiological/cellular adaptations underlying these transcriptomic changes in response to the microgravity environment will be important. It is important to note that no direct correlation can be made between gene-expression changes observed in this study and the disease status or potential for the development of disease in these individuals. This study provides an initial foundation into the molecular genetic response profiles of astronauts during spaceflight from which additional research into alterations in crew health and performance can be investigated. This kind of research could lead to human health counter-measures to mitigate risk during the transition from short-to-long-duration flight and could potentially translate to health benefits for the general public.

ACKNOWLEDGMENTS

We thank the NASA astronauts who so generously participated in these flight studies. We apologize to any authors whose work could not be cited due to space limitations. This work was funded by USRA Sponsor Award 09930-286-01 (017991-001). We also thank Andrea Throop for helpful discussions during the writing of this manuscript.

COMPETING INTERESTS

The authors declare no conflict of interest.

REFERENCES

1. Gueguinou, N. et al. Could spaceflight-associated immune system weakening preclude the expansion of human presence beyond Earth's orbit? *J Leukoc Biol* **86**, –1038 (2009).
2. Sonnenfeld, G. The immune system in space, including Earth-based benefits of space-based research. *Curr Pharm Biotechnol* **6**, 343–349 (2005).
3. Yi, B., Crucian, B., Tauber, S., Ullrich, O. & Choukèr, A. In: *Effect of Spaceflight and Spaceflight Analogue Culture on Human and Microbial Cells* (eds Nickerson C. A., Pellis N. R. & Ott C. M.) Ch. 3, 61–79 (Springer-Verlag: New York, NY, 2016).
4. Pierson, D. L., Stowe, R. P., Phillips, T. M., Lugg, D. J. & Mehta, S. K. Epstein-Barr virus shedding by astronauts during space flight. *Brain Behav Immun* **19**, 235–242 (2005).
5. Pierson, D. L., Mehta, S. K. & Stowe, R. P. In: *Psychoneuroimmunology* Vol. II (ed Ader R.) 851–868 (Academic Press, 2007).
6. Terada, M. et al. Effects of a closed space environment on gene expression in hair follicles of astronauts in the International Space Station. *PloS One* **11**, e0150801 (2016).
7. Whitney, A. R. et al. Individuality and variation in gene expression patterns in human blood. *Proc Natl Acad Sci USA* **100**, 1896–1901 (2003).
8. Obe, G. et al. Chromosomal aberrations in blood lymphocytes of astronauts after long-term space flights. *Int J Rad Biol* **72**, 727–734 (1997).
9. Smith, S. M., Zwart, S. R., Block, G., Rice, B. L. & Davis-Street, J. E. The nutritional status of astronauts is altered after long-term space flight aboard the International Space Station. *J Nutr* **135**, 437–443 (2005).
10. Kumari, R., Singh, K. P. & Dumond, J. W. Jr. Simulated microgravity decreases DNA repair capacity and induces DNA damage in human lymphocytes. *J Cell Biochem* **107**, 723–731 (2009).

11. Dumond, J. W., Singh, K. P. & Ramesh, G.T. Increase of oxidative stress and loss in expression of DNA repair genes by modeled microgravity in the testis of mice. *Cancer Res* **66**, 834 (2014).
12. Sumba, P. O. et al. Microgeographic variations in Burkitt's lymphoma incidence correlate with differences in malnutrition, malaria and Epstein-Barr virus. *Br J Cancer* **103**, 1736–1741 (2010).
13. Baqai, F. P. et al. Effects of spaceflight on innate immune function and antioxidant gene expression. *J Appl Physiol* **106**, 1935–1942 (2009).
14. Hollander, J. et al. Spaceflight downregulates antioxidant defense systems in rat liver. *Free Radic Biol Med* **24**, 385–390 (1998).
15. Kaur, I., Simons, E. R., Castro, V. A., Ott, C. M. & Pierson, D. L. Changes in monocyte functions of astronauts. *Brain Behav Immun* **19**, 547–554 (2005).
16. Rizzo, A. M. et al. Blood and oxidative stress (BOS): Soyuz mission 'Eneide'. *Microgravity Sci Technol* **19**, 210–214 (2007).
17. Bradamante, S. et al. Oxidative stress and alterations in actin cytoskeleton trigger glutathione efflux in *Saccharomyces cerevisiae*. *Biochim Biophys Acta* **1803**, 1376–1385 (2010).
18. Fang, Y. Z., Yang, S. & Wu, G. Free radicals, antioxidants, and nutrition. *Nutrition* **18**, 872–879 (2002).

This work is licensed under a Creative Commons Attribution 4.0 International License. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in the credit line; if the material is not included under the Creative Commons license, users will need to obtain permission from the license holder to reproduce the material. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>

© The Author(s) 2016

Supplementary Information accompanies the paper on the *npj Microgravity* (<http://www.nature.com/npjmgrav>)