REVIEW ARTICLE OPEN

Fatigue in Parkinson's disease: report from a multidisciplinary symposium

Joseph H Friedman^{1,2}, James C Beck³, Kelvin L Chou^{4,5}, Gracia Clark³, Christopher P Fagundes^{6,7}, Christopher G Goetz⁸, Karen Herlofson⁹, Benzi Kluger¹⁰, Lauren B Krupp¹¹, Anthony E Lang¹², Jao-Shin Lou¹³, Laura Marsh^{14,15}, Anne Newbould³ and Daniel Weintraub¹⁶

Fatigue is a severe problem for many people living with Parkinson's disease (PD). Best estimates suggest that more than 50% of patients experience this debilitating symptom. Little is known about its etiology or treatment, making the understanding of fatigue a true unmet need. As part of the Parkinson's Disease Foundation Community Choice Research Program, patients, caregivers, and scientists attended a symposium on fatigue on 16 and 17 October 2014. We present a summary of that meeting, reviewing what is known about the diagnosis and treatment of fatigue, its physiology, and what we might learn from multiple sclerosis (MS), depression, and cancer—disorders in which fatigue figures prominently too. We conclude with focused recommendations to enhance our understanding and treatment of this prominent problem in PD.

npj Parkinson's Disease (2016) 2, 15025; doi:10.1038/npjparkd.2015.25; published online 14 January 2016

INTRODUCTION

Fatigue: The patient perspective

Anne Newbould (care partner): My husband of almost 40 years, James, has Parkinson's disease, and fatigue is his most severe and debilitating symptom. At first I thought he had depression. I attempted to get him interested in the things that we so loved to do, but it seemed that he lost his zeal for life. When I asked him about it, he told me that he was extremely tired and it almost hurt to try to get up out of bed or his chair. I started researching all I could on Parkinson's and fatigue, but there was very little information. I felt lost and unbelievably helpless. *We went from being participants to spectators in life.* With James' fatigue, we have had to alter our lifestyle to conform to his level of energy. We go to baseball games instead of fly-fishing. We cruise on the local Southern Bell riverboat instead of racing sailboats. We go to the beach and watch the ocean instead of teaching scuba diving. That is why James and I were ecstatic to have the opportunity to raise this issue through the Parkinson's Disease Foundation's Community Choice Research Awards.

Gracia Clark (patient): Fatigue was the first Parkinson's symptom I noticed; it is still my most debilitating one. Feeling fatigued is different from feeling sleepy, depressed or apathetic. As an anthropologist who has worked in West

Africa, I am reminded of the fatigue I suffered after coming down with malaria. When fighting malaria, I faced an overwhelming need to lie down. But unlike fatigue from malaria, Parkinson's fatigue doesn't go away after taking a few malaria pills. Finding treatment for Parkinson's fatigue can be a huge challenge. Medications approved for other diseases—such as narcolepsy—may provide some benefit, but their efficacy in Parkinson's is under studied, and insurance companies can decline to cover such off-label uses. Furthermore, because many diseases can cause fatigue, receiving an accurate diagnosis can be difficult, as it was in my case. I have heard many people with Parkinson's say fatigue is one of their most debilitating symptoms, and yet the symptom is not well understood. There is an urgent need for increased research to better understand fatigue and find ways to treat it.

Fatigue is a common symptom in almost all medical, neurologic, and psychiatric disorders as well as in the otherwise healthy general population.¹ We considered fatigue a subjective symptom, different from muscle fatigue, which is a physiologically defined entity. There are a number of definitions used to describe subjective fatigue,² and all focus on the concept of an abnormal and excessive lack of energy that interferes with normal function. Although the recognition of the importance of fatigue in Parkinson's disease (PD) is relatively new, it should be noted that Parkinson-related fatigue had been recognized in the nineteenth century by some of the major clinical neurologists of their time.³

¹Movement Disorders Program, Butler Hospital, Providence, RI, USA; ²Department of Neurology, Alpert Medical School of Brown University, Providence, RI, USA; ³Parkinson's Disease Foundation, New York, NY, USA; ⁴Department of Neurology, University of Michigan Health System, Ann Arbor, MI, USA; ⁵Department of Neurosurgery, University of Michigan Health System, Ann Arbor, MI, USA; ⁶Department of Psychology, Rice University, M.D. Anderson Cancer Center, Houston, TX, USA; ⁷Department of Health Disparities, Houston, TX, USA; ⁸Department of Neurological Sciences, Rush University Medical Center, Chicago, IL, USA; ⁹Department of Neurology, Sorlandet Hospital, Arendal, Norway; ¹⁰Department of Neurology, University of Colorado Anschutz Campus, Aurora, CO, USA; ¹¹Department of Neurology, Stony Brook University Medical Center, Stony Brook, NY, USA; ¹²Toronto Western Hospital, Morton and Gloria Shulman Movement Disorders Clinic and Edmond J Safran Program in Parkinson's Disease, Toronto, ON, Canada; ¹³University of North Dakota School of Medicine and Health Services, Sanford Health, Grand Forks, ND, USA; ¹⁴Department of Psychiatry, Baylor College of Medicine, Mental Health Care Line, Michael E. DeBakey Veterans Affairs Medical Center, Houston, TX, USA; ¹⁵Department of Neurology, Baylor College of Medicine, Mental Health Care Line, Michael E. DeBakey Veterans Affairs Medical Center, Houston, TX, USA and ¹⁶Department of Psychiatry, Perelman School of Medicine, University of Pennsylvania, Philadelphia, PA, USA.

Correspondence: JH Friedman (Joseph_friedman@brown.edu)

Received 23 May 2015; revised 28 September 2015; accepted 25 October 2015

DEFINING FATIGUE

There are several difficulties in defining fatigue. Fatigue may occur in more than one domain^{4,5} and different scales include different domains. Defining the point at which normal fatigue crosses into pathological fatigue is also challenging. Operationally, fatigue becomes pathological when it is abnormally severe, restricts activities independently of other constraints induced by PD, and is associated with psychological distress; however, it remains difficult to define an exact diagnostic threshold.⁶ Fatigue may be associated or due to other behavioral disorders such as depression, anxiety, and apathy, which are themselves intrinsic to PD.⁷ In addition, there may be confusion between fatigue, a feeling present at rest, and fatigability, a problem induced by activity.⁵ Although PD fatigue has traditionally been considered a symptom, it may be better considered as a syndrome. Therefore, we propose to define PD fatigue as *a sense of exhaustion unexplained by drug effects, other medical, or psychiatric disorders, present for a defined period, and associated with other fatigue-related symptoms, such as reduced motivation and nonrestorative rest, or constraints on activities.*⁶ Such a syndromic definition may provide several benefits for clinical research by improving comparability across studies but might also inhibit recruitment if the definition is too narrow or inflexible. However, this definition remains quite subjective, and until objective measures are discovered, which reflect the sensation of exhaustion, we will continue to be limited in our ability to understand the pathophysiology of this disorder.

CLINICAL REVIEW OF FATIGUE IN PD

Motor fatigue

Most studies of PD fatigue have focused on physical, or motor, fatigue. Case-control studies^{8,9} and community-based studies¹⁰ show fatigue to be both common and severe, with the prevalence ranging between 33 and 70%.¹¹ This broad range probably reflects different patient populations and different evaluation methodologies. This prevalence makes it one of the most common nonmotor PD problems,¹² even in the early stages of the disease. It is clearly a "pre-motor" feature as well, appearing well before motor symptoms.^{13,14} A UK study spanning 16 years, involving 8,166 patients with PD and 46,755 controls, found that the presence of fatigue 5 years before the diagnosis of PD increased the relative risk of developing the disease to 1.56 (ref. 15).

Patients describe their fatigue as different from what they had experienced previously.¹⁶ Unlike fatigue in the general population, PD fatigue often improves with exercise.^{16,17} Two studies evaluating fatigue in newly diagnosed, untreated patients reported it to be a clinically relevant problem^{3,6,10,14} even when motor symptoms were minimal. Over 50% of patients consider fatigue among their three most disabling symptoms.⁸ In spite of this prominent concern by patients, fatigue is often not recognized by the treating neurologist.¹⁸

The evolution of fatigue has not been well studied. Limited data indicate that, once present, fatigue typically persists and usually worsens over time.^{19,20}

Studies of PD-related fatigue are discordant regarding the correlation between fatigue and motor symptom severity, disease stage, or duration.^{2,14} These conflicting results may be explained by different populations as well as different assessment tools. Multiple, but not all, studies in PD show a significant correlation between depression and fatigue.⁸ Anxiety and apathy also correlate with fatigue^{21,22} but many fatigued patients are not depressed, anxious, or apathetic.⁸ Several studies suggest that cognitive performance, frontal lobe activity, and self-reported cognitive impairment are associated with fatigue.^{23,24} However, fatigability on cognitive tasks may not correlate with subjective fatigue.²⁵ Similar to depression, fatigue is frequently associated

with daytime sleepiness and sleep disorders, but can be distinguished from them by the absence of sleep being restorative, and may occur in patients with normal sleep patterns.^{8,9,26} Finally, objectively measured autonomic dysfunction has been associated with fatigue in PD.^{27,28}

Cognitive fatigue

Cognitive fatigue refers to the subjective experience of feeling weary or exhausted during an intellectually challenging task, along with a decreased capacity to initiate or sustain cognitively challenging activities. Cognitive fatigue is sometimes referred to as "central fatigue." Patients have reduced attention and may have deficits on formal neuropsychological testing in cognitive areas of memory, learning, attention, and information process.²⁹ A limited number of studies have examined cognitive fatigue in PD.³⁰ In some patients, cognitive performance is unimpaired; however, the increased mental effort required to maintain a normal level of performance results in fatigue and distress. Conversely, cognitive impairment, especially executive dysfunction, may be associated with cognitive fatigue, as the patient is less able to maintain concentration or think about complex matters. For example, a longitudinal decline in Mini-Mental State Examination score over 8 years was associated with a concurrent increase in fatigue symptoms.¹⁹ In addition, reduced performance on the Wisconsin Card Sorting Task, reduced frontal lobe perfusion, and depression were all correlated with fatigue.²³ The development and experience of fatigue is also context-dependent; cognitive fatigue emerges as task complexity increases and attentional demands are greater. Even in healthy controls, increased task complexity during a walking exercise affects gait speed; however, the impact on physical fatigue and gait slowness is greater in patients with PD.³¹

INSIGHTS INTO FATIGUE FROM OTHER DISORDERS

Although perceived fatigue is probably similar among different illnesses,³² there may be disease-specific differences. Both similarities and differences may provide insights into fatigue pathophysiology. Similarities in fatigue would be compatible with a hypothesis of shared mechanisms, and thus the possibility of similar interventions. The high prevalence of fatigue in medical and psychiatric disorders points either to some very generalized mechanisms or a limitation in the brain's ability to distinguish perceptions. We limited our review of other disorders with prominent fatigue, but excluded chronic fatigue syndrome because of its frequent association with psychiatric disorders such as personality disorders and post-traumatic stress disorder that might confound physiological interpretations.

CANCER

Fatigue has been most studied in oncology. Cancer specialists have developed a disease-specific definition for "cancer-related fatigue" as a syndrome that is included in the ICD 10.⁶ The goal of such a definition was to accelerate the "development of a research agenda." "Fatigue is the most common symptom experienced by adults and children with cancer" (ibid). The criteria look much like typical criteria used in the Diagnostic and Statistical Manual V,³³ the guide for psychiatric diagnoses, with a requirement for "significant fatigue" plus 5 symptoms from a list of 10, with symptoms present daily or almost daily for a 2-week period in the previous month.

Cancer, of course, is not a single disease. Many of the links found between certain cancers and fatigue have not been evaluated in other cancers. Cytokines, released during inflammation, have been correlated with fatigue in cancer.^{34,35}

Hypothalamic–pituitary axis alterations have been implicated.³⁶ Changes in circadian rhythm³⁷ have been associated with fatigue as has a genetic predisposition.³⁶ Psychosocial factors also have a high correlation with fatigue.³⁸

MULTIPLE SCLEROSIS

The estimated prevalence of fatigue in multiple sclerosis (MS) is ~80%. "Chronic persistent fatigue" is defined for MS as significant fatigue present for more than 50% of each day for a minimum of 6 weeks.³⁹ There are some facets of fatigue that distinguish MS fatigue from fatigue associated with other disorders. Unlike PD, fatigue in MS does not decrease with exercise.¹⁷ Heat worsens fatigue in MS, an unusual but consistent observation compared with other disorders.⁴⁰ This has not been studied in PD but anecdotal experience indicates that PD patients routinely prefer warm to cold climates, suggesting that heat is unlikely to worsen fatigue.

Fatigue severity in MS is linked to disease type (chronic progressive more likely to be affected), but, not clearly linked to disability or disease duration.³⁹ It is closely related to concurrent depression, and the presence of depression or anxiety predicts the later occurrence of fatigue. Similarly, the presence of fatigue is a risk factor for later development of depression and anxiety. Sleep disorders are increased in MS subjects with fatigue. Lower levels of education are associated with worse fatigue. As in PD,⁴¹ the sensation of fatigue does not correlate with motor fatigue, the loss of strength with repeated effort (see below). Fatigue has been associated with a variety of gray matter changes in MS as well as functional magnetic resonance imaging changes during tasks.⁴⁰ There may be associations between fatigue and neuroendocrine, autonomic, or neuroimmune markers.

DEPRESSION

Although fatigue is one of the symptoms used to diagnose major depression in the general population,³³ it is often unrelieved when depression improves.^{42–44} Thus, even when the fatigued PD patient has depression, treatment of the latter may not improve the former, thus implying that fatigue may not always simply result from depression.

The data on cancer and MS fatigue are instructive in terms of identifying areas of research that have not been adequately explored in PD. These areas include pain, socioeconomic and educational status, a variety of psychosocial measures, markers of inflammation and immune function, deconditioning, and autonomic dysfunction; none of these observations in other disorders have resulted in clear benefit to patients. A study comparing consecutive people with MS to those with PD, both groups attending similar academic subspecialty neurology clinics, found that the spectrum of symptoms endorsed in the Fatigue Severity Scale (FSS) were not identical.¹⁷ In addition, patients with PD generally reported that fatigue improved with exercise, whereas those with MS reported the opposite. Whether this counter-intuitive observation about exercise is unique to PD fatigue remains to be seen.

CLINICAL MEASURES OF FATIGUE IN PD

A systematic critique of rating scales for diagnosis and severity of PD fatigue used pre-determined criteria for Recommended, Suggested, or Listed, depending on the quality of data.⁴⁵ The FSS met the necessary criteria to be recommended for both diagnostic screening and severity measurement.^{2,46} The Multi-dimensional Fatigue Inventory (MFI)⁴ was designated as recommended for rating fatigue severity, and may be more sensitive to change with interventions than the FSS.⁴⁶ For diagnostic screening only, two other scales were recommended: Functional

Assessment of Chronic Illness Therapy—Fatigue Scale (FACIT-F)⁴⁷ and Parkinson Fatigue Scale (PFS or PFS-16).¹⁶ Since the MDS review article, the Modified Fatigue Impact Scale has been validated in a study involving 100 PD patients.⁴⁸ This scale involves evaluation of cognitive as well as physical and social functioning.

The above scales measure subjective fatigue in PD. Objective motor fatigue can also be measured in PD with performance changes during sustained muscle contractions, repetitive movements, or timed activities. PD studies generally show heightened motor fatigability, which does not typically correlate with subjective fatigue.⁴⁹ Objective measures of cognitive fatigability focus on performance change during a continuous mental task or performance change in another target task after a cognitive challenge.^{50,51} Performance variability over time may be a sensitive metric of fatigability that correlates with fatigue complaints.⁵¹ A study using the Borg Perceived Exertion Rating⁵² measuring perceived effort documented heightened perception in PD in comparison with controls, despite similar levels of performance.⁵³

Physiology and biomarkers for fatigue in PD

The physiology underlying fatigue symptoms in PD is unknown, whereas much is known about the mechanisms of motor fatigability. The only published study of physiological differences between fatigued and nonfatigued PD patients found no measurable differences in oxygen utilization during exercise,⁵⁴ but some studies have suggested that exercising improves fatigue.^{15,17,55,56} Whether fatigue fluctuates with motor fluctuations was looked at in one study.⁵⁷ Subjects were assessed during their off, whereas all other studies used the subjects' experience over a preceding time interval of weeks. Eighty-eight percent of their subjects were fatigued, and fatigue increased with motor "off." Too few subjects were nonfatigued to determine how often fatigue occurred only during motor "off." The physiological implications are unclear, as many other nonmotor symptoms also increased during the "off" period.

Increased motor fatigability in PD is associated with increased cortical excitability. Motor fatigability is defined as the rate of force deterioration in an exercise paradigm. Cortical excitability may be measured by transcranial magnetic stimulation applied to the motor cortex with the compound muscle action potentials recorded from the muscles of interest such as the abductor pollicis brevis or the first dorsal interosseous muscle. Cortical excitability is increased when the motor-evoked potential amplitude increases while transcranial magnetic stimuli are held constant. The cortical excitability is markedly increased in PD at rest, and during intermittent exercise that correlates negatively with the severity of motor fatigue—the higher the cortical excitability, the less motor fatigability.⁵⁸ The increase in cortical excitability is partially normalized by a small dose of levodopa that also reduces motor fatigability;^{58,59} however, neither cortical excitability nor motor fatigability correlated with subjective fatigue complaints. Thus, cortical excitability can serve as a biomarker for motor fatigability in PD but not for subjective fatigue.

BIOMARKERS

Inflammatory markers such as C-reactive protein and cytokines are potential biomarkers for subjective fatigue. In a study comparing 87 PD subjects to 33 controls, several markers of inflammation were assessed in the cerebrospinal fluid and correlated with a variety of nonmotor symptoms. Elevated C-reactive protein and monocyte chemoattractant protein-1 were significantly correlated with fatigue, but this association requires confirmation before being considered clinically relevant.⁶⁰ How serum markers relate to brain

Table 1. Published trials of treatments for fatigue

Modafinil
Methylphenidate
Pergolide
Bromocriptine
Pramipexole
Doxepin
Rasagiline
L-Dopa
Memantine
Caffeine
Behavioral interventions
Exercise
Amantadine
(from 60)

changes in PD is speculative; however, they may relate to autonomic changes outside the central nervous system.

Imaging of fatigue in PD

Motor symptoms in PD result from nigrostriatal dopaminergic denervation, but dopaminergic dysfunction does not appear to be related to fatigue in PD. In the ELLDOPA cohort, 49 levodopa-naïve PD patients with fatigue had similar [123 I]- β -CIT striatal dopamine transporter uptake as 82 PD patients without fatigue.¹⁴ Another study showed no difference in 18 F-dopa uptake between 10 fatigued PD subjects and 10 nonfatigued PD subjects.^{2,61}

Serotonin transporter uptake has been reported to be reduced in chronic fatigue syndrome,^{3,61} suggesting that nondopaminergic pathways may be involved in PD fatigue. Pavese *et al.*⁶² compared serotonergic transporter uptake (using the positron emission tomography (PET) ligand 11 C-DASB) in seven PD subjects with fatigue and eight PD subjects without fatigue. Serotonin transporter binding in the caudate, putamen, ventral striatum, insula, and thalamus was decreased in the fatigued patients. The relationship of the cholinergic system to PD fatigue has not been investigated. The autonomic nervous system may also be involved in PD fatigue. One study found that pressor responses in norepinephrine and dobutamine infusion tests were greater and MIBG cardiac uptake was decreased in PD subjects with fatigue compared with those without fatigue.^{4,27}

CURRENT TREATMENT APPROACHES

There are no evidence-based guidelines for treating fatigue in PD. A recent review⁶³ found only 14 published randomized, controlled trials using validated fatigue measures in PD. No study of nonpharmacological treatments had showed benefit for fatigue. Only one study showed a clinically and statistically significant improvement with stimulants;⁶⁴ however, when data from other stimulant trials were pooled, no effect was seen. Statistically but clinically insignificant benefits were found in *post hoc* analyses of rasagiline,⁶⁵ and small benefits were seen for levodopa and other drugs when fatigue was one of several nonmotor symptoms assessed in randomized, placebo-controlled trials, where nonmotor symptoms were not the primary end points of the study. Nonpharmacologic approaches to treat cognitive fatigue include exercise, daily scheduling, planned rests, and pacing daily activities, similar to what has been tried in treating motor fatigue. The Table 1 summarizes interventions reported upon.

Given the limited evidence regarding treatment of PD fatigue,⁶⁶ a commonsense approach to treat fatigue is recommended, and it will vary with the treating clinician. When counseling patients, it is important to acknowledge fatigue as an important and common symptom in PD, and not necessarily a reflection of depression.

Many patients will have other potential contributors to fatigue, including depression, anxiety, apathy, sleep disorders and concomitant medical problems, and medications. These disorders are not always treatable. Although exercise has not been demonstrated in rigorous trials to reduce fatigue in PD, an interesting and repeated counterintuitive observation is that PD patients often report that they feel energized *after* exercising.^{16,17} As exercise is generally considered an important therapy for most aspects of PD and most medical disorders, we think exercise should always be suggested as there is no drawback.

CURRENT/FUTURE RESEARCH

Current research on PD-related fatigue is focused on identification of biological markers and correlates of fatigue that may provide insights into the development of effective treatments. The Parkinson Progression Marker Initiative (PPMI) study (www.ppmi-info.org) is a biomarker-rich study focused on the course of early PD including ~425 newly diagnosed PD patients and ~200 matched health controls who undergo detailed assessments at baseline and annually (out to 5 years). Fatigue and its severity, as derived from the MDS-UPDRS fatigue item, will be evaluated for correlations with other measures including the following: serial dopamine transporter imaging, blood drawn for genotyping and assessment of plasma proteins, cerebrospinal fluid, serial magnetic resonance imaging brain imaging (structural and diffusion tensor imaging), and nonmotor phenomena, such as cognitive functioning, depression, anxiety, and daytime sleepiness.

Two additional ongoing studies on fatigue in PD (personal communication) are using PET to examine neurophysiological correlates of fatigue. Preliminary results from FDG PET suggest a role for anterior cingulate, insular, superior temporal, and precuneus regions in PD fatigue (Strafella *et al.*, unpublished results). In a separate large PET study, the microglial ligand phenoxyanilide ([18 F]-FEPPA) is being used to examine the role of neuroinflammation in PD relative to fatigue severity based on the FSS.

Areas for future work include identification of which neurotransmitter systems or neural circuits, if any, are salient to the pathophysiology of fatigue in PD. Imaging of brain activity and other physiological measures will likely be utilized.

Comparing voxel-based morphometry data in PD patients with fatigue as well as older adults without fatigue may delineate brain structures associated with PD-related fatigue and whether these associations are similar to those seen in fatigue in older adults without PD.

Functional magnetic resonance imaging should be performed in fatigued and nonfatigued PD patients as free of confounding problems as possible (e.g., free of psychiatric or medical problems or medications associated with fatigue).

Several nonpharmacologic strategies have shown evidence for improving fatigue in MS but have not been tested in PD, including exercise, energy management strategies, and mindfulness-based stress reduction.⁶⁷

Conclusions

Further research is required to better understand fatigue in PD. The lack of progress in understanding the pathophysiology of fatigue, or its treatment, in more intensively studied disorders and the absence of any effective therapy for fatigue in any of them argues strongly for a greater effort. We believe that research to better understand the syndrome and treatment trials based on rational hypotheses of pathophysiology should proceed in parallel.

ACKNOWLEDGMENTS

We thank the Parkinson's community for their inspiration in addressing this debilitating unmet need. We thank the Parkinson's Disease Foundation for their generous financial and logistical support for our gathering to discuss this issue.

CONTRIBUTIONS

All authors contributed intellectual material, and all edited the final manuscript. JHF, KLC, GC, CGG, KH, AEL, JSL, LM, AN DW wrote sections of the manuscript.

COMPETING INTERESTS

JHF: Teva, Auspex, Osmotica, Acadia, Lundbeck, Avid, Medscape, Theravance. KLC has received research support from the NIH (NS44504-08, NS091856-01) and the Michael J. Fox Foundation, has participated as a site-PI in clinical trials sponsored by the Huntington Study Group (2CARE) and Osmotica, receives royalties from UpToDate and Demos Health and serves as a consultant for Medtronic and Accordant. CGG: Consulting or Advisory Board Membership with honoraria from Acadia (Deborah Wood Associates), AstraZeneca, Avanir, Boston Scientific, Clearview, Health Advances, Chelsea Pharmaceuticals (Link Medical Communications), ICON Pricespective LLC, MED-IQ Educational Services, Neurocrine, Pfizer, Teva, and WPP Group Kantor Health LLC. He receives funding from NIH and Michael J Fox Foundation. CGG directs the Rush Parkinson's Disease Research Center that receives support from the Parkinson's Disease Foundation. He directs the translation program for the MDS-UPDRS and UDysRS and receives funds from the International Parkinson and Movement Disorder Society (IPMDS) for this effort. He receives honoraria from the University of California, University of Luxembourg, University of Rochester, World Parkinson Coalition. He has Royalties of Oxford University Press, Elsevier Publishers, Wolters Kluwer Health-Lippincott, and Wilkins and Wilkins, and receives salary from the Rush University Medical Center. KH: Allergan, Abbvie, Novartis. LKB: Johnson and Johnson; Bristol Myers Squibb. AEL has served as an advisor for Abbvie, Allon Therapeutics, Avanir Pharmaceuticals, Biogen Idec, Boehringer-Ingelheim, Bristol Myers Squibb, Ceregene, Cipla, Intekrin, Lilly, Medtronic, Merck, Novartis, NeuroPhage Pharmaceuticals, Teva, and UCB; received honoraria from Medtronic, Teva, UCB, and AbbVie; received grants from Brain Canada, Canadian Institutes of Health Research, Edmond J Safra Philanthropic Foundation, Michael J. Fox Foundation, the Ontario Brain Institute, National Parkinson Foundation, Parkinson Society Canada, Physicians Services Incorporated (PSI), Tourette Syndrome Association, and W. Garfield Weston Foundation; received publishing royalties from Saunders, Wiley-Blackwell, Johns Hopkins Press, and Cambridge University Press; and has served as an expert witness in cases related to the welding industry. DW: receives research funding or support from Michael J. Fox Foundation for Parkinson's Research, National Institutes of Health, Novartis Pharmaceuticals, Department of Veterans Affairs, Avid Radiopharmaceuticals, and Alzheimer's Disease Cooperative Study; honoraria from AbbVie, Biotie, Teva Pharmaceuticals, Otsuka, UCB, Clintrex LLC, and the CHDI Foundation; license-fee payments from the University of Pennsylvania for the QUIP and QUIP-RS; royalties from Wolters Kluwerland; and fees for legal consultation for lawsuit related to antipsychotic prescribing in a patient with Parkinson's disease. The remaining authors declare no conflict of interest.

REFERENCES

- Wright, J. & O'Connor, K. M. Fatigue. *Med. Clin. N. Am.* **98**, 597–608 (2014).
- Friedman, J. H. et al. Fatigue in Parkinson's disease: a review. *Mov. Disord.* **22**, 297–308 (2007).
- García-Ruiz, P. J., Chaudhuri, K. R. & Martínez-Marín, P. Non-motor symptoms of Parkinson's disease: a review from the past. *J. Neurol.* **338**, 30–33 (2014).
- Smets, E. M. A., Garssen, B., Bonke, B. & de Haes, J. C. J. M. The Multidimensional Fatigue Inventory (MFI): psychometric qualities of an instrument to assess fatigue. *J. Psychosom. Res.* **39**, 312–325 (1995).
- Alexander, N. B. et al. Bedside-to-bench conference: research agenda for idiopathic fatigue and aging. *J. Am. Geriatr. Soc.* **58**, 967–975 (2010).
- Barsevick, A. M. et al. Recommendations for high-priority research on cancer-related fatigue in children and adults. *J. Natl Cancer Inst.* **105**, 1432–1440 (2013).
- Weintraub, D. & Burn, D. M. Parkinson's disease: the quintessential neuropsychiatric disorder. *Mov. Disord.* **26**, 1022–1031 (2011).
- Friedman, J. & Friedman, H. Fatigue in Parkinson's disease. *Neurology* **43**, 2016–2018 (1993).
- van Hilten, J. J. et al. Sleep, excessive daytime sleepiness and fatigue in Parkinson's disease. *J. Neural Transm. Park. Dis. Dement. Sect.* **5**, 235–244 (1993).
- Herlofson, K., Ongre, S. O., Enger, L. K., Tysnes, O. B. & Larsen, J. P. Fatigue in early Parkinson's disease. Minor inconvenience or major distress? *Eur. J. Neurol.* **19**, 963–968 (2012).

- Friedman, J. H., Abrantes, A. & Sweet, L. H. Fatigue in Parkinson's disease. *Exp. Opin. Pharmacother.* **12**, 1999–2007 (2011).
- Barone, P. et al. The PRIAMO study: a multicenter assessment of nonmotor symptoms and their impact on quality of life in Parkinson's disease. *Mov. Disord.* **24**, 1641–1649 (2009).
- Post-Sunyer, C. et al. The onset of nonmotor symptoms in Parkinson's disease (the ONSET PD study). *Mov. Disord.* **30**, 229–237 (2015).
- Schiffitto, G. et al. Fatigue in levodopa-naïve subjects with Parkinson disease. *Neurology* **71**, 481–485 (2008).
- Schrag, A. et al. Pre diagnostic presentations of Parkinson's disease in primary care: a case control study. *Lancet Neurol.* **14**, 57–64 (2015).
- Brown, R. G., Dittner, A., Findley, L. & Wessely, S. C. The Parkinson fatigue scale. *Parkinsonism Relat. Disord.* **11**, 49–55 (2005).
- Cochrane G. S. et al. Fatigue and apathy in patients with either Parkinson's disease or multiple sclerosis. *Parkinsonism Relat. Disord.* **21**, 1093–1095 (2015).
- Shulman, L. M., Taback, L., Rabinstein, A. A. & Weiner, W. J. Non-recognition of depression and other non-motor symptoms of Parkinson's disease. *Parkinsonism Relat. Disord.* **8**, 193–197 (2002).
- Alves, G., Wentzel-Larsen, T. & Larsen, J. P. Is fatigue an independent and persistent symptom in patients with Parkinson disease? *Neurology* **63**, 1908–1911 (2004).
- Friedman, J. H. & Friedman, H. Fatigue in Parkinson's disease: a nine-year follow-up. *Mov. Disord.* **16**, 1120–1122 (2001).
- Shulman, L. M., Taback, R. L., Bean, J. & Weiner, W. J. Comorbidity of the nonmotor symptoms of Parkinson's disease. *Mov. Disord.* **16**, 507–510 (2001).
- Skorvanek, M. et al. The associations between fatigue, apathy, and depression in Parkinson's disease. *Acta Neurol. Scand.* **131**, 80–87 (2014).
- Abe, K., Takanashi, M. & Yanagihara, T. Fatigue in patients with Parkinson's disease. *Behav. Neurol.* **12**, 103–106 (2000).
- Goldman, J. G., Stebbins, G. T., Leung, V., Tilley, B. C. & Goetz, C. G. Relationships among cognitive impairment, sleep, and fatigue in Parkinson's disease using the MDS-UPDRS. *Parkinsonism Relat. Disord.* **20**, 1135–1139 (2014).
- Lou, J. S. Physical and mental fatigue in Parkinson's disease: epidemiology, pathophysiology and treatment. *Drugs Aging* **26**, 195–208 (2009).
- Stocchi, F. et al. Prevalence of fatigue in Parkinson disease and its clinical correlates. *Neurology* **83**, 215–220 (2014).
- Nakamura, T. et al. Does cardiovascular autonomic dysfunction contribute to fatigue in Parkinson's disease? *Mov. Disord.* **26**, 1869–1874 (2011).
- Jain, S. et al. Pupillary unrest correlates with arousal symptoms and motor signs in Parkinson disease. *Mov. Disord.* **26**, 1344–1347 (2011).
- Chaudhuri, A. & Behan, P. O. Fatigue in neurological disorders. *Lancet* **363**, 978–988 (2004).
- Dashtipour, K. et al. Effects of exercise on motor and non-motor symptoms of Parkinson's disease. *Parkinsons Dis.* **2015**, 586378 (2015).
- Rochester, L. et al. Attending to the task: interference effects of functional tasks on walking in Parkinson's disease and the roles of cognition, depression, fatigue and balance. *Arch. Phys. Med. Rehabil.* **85**, 1578–1585 (2004).
- Jones, D. E. J., Gray, J. C. & Newton, J. Perceived fatigue is comparable between different disease groups. *Q. J. Med.* **102**, 617–624 (2009).
- American Psychiatric Association *Diagnostic and Statistical Manual* 5th (edn). American Psychiatric Press, 2014.
- Schubert, C., Hong, S., Natarajan, L., Mills, P. J. & Dimsdale, J. E. The association between fatigue and inflammatory marker levels in cancer patients: a quantitative review. *Brain Behav. Immun.* **21**, 413–427 (2007).
- Seruga, B., Zhang, H., Bernstein, L. J. & Tannock, I. F. Cytokines and their relationship to the symptoms and outcome of cancer. *Nat. Rev. Cancer* **8**, 887–899 (2008).
- Ryan, J. L. et al. Mechanisms of cancer-related fatigue. *Oncologist* **12**, 22–34 (2007).
- Hrushesky, W. J. et al. Circadian clock manipulation for cancer prevention and control and the relief of cancer symptoms. *Integr. Cancer Ther.* **8**, 387–397 (2009).
- Aouizerat, B. E. et al. Preliminary evidence of a genetic association between tumor necrosis factor alpha and the severity of sleep disturbance and morning fatigue. *Biol. Res. Nurs.* **11**, 27–41 (2009).
- Jaremk, L. M. et al. Loneliness predicts pain depression and fatigue: understanding the role of immune dysregulation. *Psychoneuroendocrinology* **38**, 1310–1317 (2013).
- Multiple Sclerosis Council for Clinical Practice Guidelines. *Fatigue and Multiple Sclerosis: Evidence-based Management Strategies for Fatigue in Multiple Sclerosis*. Paralyzed Veterans Assoc of America, 1998.
- Krupp, L. B., Serafin, D. J. & Christodoulou, C. Multiple sclerosis associated fatigue. *Exp. Rev. Neurother.* **10**, 1437–1447 (2010).
- Robinson R. L. et al. The importance of unresolved Fatigue in Depression: Costs and Comorbidities. *Psychosom* 2015; **56**: 274–285.

43. Nierenberg, A. A. *et al.* Residual symptoms after remission in major depressive disorder with citalopram and risk of relapse: a STAR*D report. *Psychol. Med.* **40**, 41–50 (2010).
44. Fava, M. *et al.* A cross sectional study of the prevalence of cognitive and physical symptoms during long term antidepressant treatment. *J. Clin. Psychiatr.* **67**, 1754–1759 (2006).
45. Friedman, J. H. *et al.* Fatigue rating scales critique and recommendations by the Movement Disorders Society task force on rating scales for Parkinson's disease. *Mov. Disord.* **25**, 805–822 (2010).
46. Krupp, L. B., LaRocca, N. G., Muir-Nash, J. & Steinberg, A. D. The Fatigue Severity Scale. Application to patients with multiple sclerosis and systemic lupus erythematosus. *Arch. Neurol.* **46**, 1121–1123 (1989).
47. Cella, D. *et al.* Validation of the Functional Assessment of Chronic Illness Therapy-Fatigue (FACIT-F) Scale relative to other instrumentation in patients with rheumatoid arthritis. *J. Rheumatol.* **32**, 811–819 (2005).
48. Schiehser, D. M. *et al.* Validation of the Modified Fatigue Impact Scale in Parkinson's disease. *Parkinsonism Relat. Disord.* **19**, 335–338 (2013).
49. Lou, J. S. *et al.* Levodopa improves physical fatigue in Parkinson's disease: a double-blind, placebo-controlled, crossover study. *Mov. Disord.* **18**, 1108–1114 (2003).
50. Kluger, B. M., Krupp, L. B. & Enoka, R. M. Fatigue and fatigability in neurologic illnesses: proposal for a unified taxonomy. *Neurology* **80**, 409–416 (2013).
51. Wang, C., Ding, M. & Kluger, B. M. Change in intraindividual variability over time as a key metric for defining performance-based cognitive fatigability. *Brain Cogn.* **85**, 251–258 (2014).
52. Borg, G. *Borg's Perceived Exertion and Pain Scales*. Human Kinetics, 1988.
53. Solomon, N. P. & Robin, D. A. Perceptions of effort during handgrip and tongue elevation in Parkinson's disease. *Parkinsonism Relat. Disord.* **11**, 353–361 (2005).
54. Garber, C. E. & Friedman, J. H. Effects of fatigue on physical activity and function in patients with Parkinson's disease. *Neurology* **60**, 1119–1124 (2003).
55. Kelly, N. A. *et al.* Novel, high intensity exercise prescription improves muscle mass, mitochondrial function and physical capacity in individuals with Parkinson's disease. *J. Appl. Physiol.* **116**, 582–592 (2014).
56. Cugusi L. *et al.* Effects of an adapted physical activity program on motor and non-motor functions and quality of life in patient with Parkinson's disease. *NeuroRehabil* 2014; **35**: 789–794.
57. Storch A. *et al.* Non motor fluctuations in Parkinson's disease: severity and correlation with fatigue motor symptoms. *Neurology* **80**, 300–309 (2013).
58. Lou, J. S., Benice, T., Sexton, G. & Nutt, J. Levodopa normalizes exercise related cortico-motoneuron excitability abnormalities in Parkinson's disease. *Clin. Neurophysiol.* **114**, 930–937 (2003a).
59. Lou, J. S. *et al.* Levodopa improves physical fatigue in Parkinson's disease—a double-blind, placebo-controlled crossover study. *Mov. Disord.* **18**, 1108–1114 (2003b).
60. Lindqvist, D. *et al.* Cerebrospinal fluid inflammatory markers in Parkinson's disease—associations with depression, fatigue, and cognitive impairment. *Brain Behav. Immun.* **33**, 183–189 (2013).
61. Yamamoto, S. *et al.* Reduction of serotonin transporters of patients with chronic fatigue syndrome. *Neuroreport* **15**, 2571–2574 (2004).
62. Pavese, N., Metta, V., Bose, S. K., Chaudhuri, K. R. & Brooks, D. J. Fatigue in Parkinson's disease is linked to striatal and limbic serotonergic dysfunction. *Brain* **133**, 3434–3443 (2010).
63. Franssen, M., Winward, C., Collett, J., Wade, D. & Dawes, H. Interventions for fatigue in Parkinson's disease: a systematic review and meta-analysis. *Mov. Disord.* **29**, 1675–1678 (2014).
64. Mendonca, D. A., Menezes, K. & Jog, M. S. Methylphenidate improves fatigue scores in Parkinson's disease: a randomized controlled trial. *Mov. Disord.* **22**, 2070–2076 (2007).
65. Smith, K. M., Eyal, E. & Weintraub, D. ADAGIO Investigators. Combined rasagiline and antidepressant use in Parkinson's disease in the ADAGIO study: effects on nonmotor symptoms and tolerability. *JAMA Neurol.* **72**, 88–95 (2015).
66. Lamotte, G., Rafferty, M. R., Prodoehl, J. *et al.* Effects of endurance exercise training on the motor and non-motor features of Parkinson's disease: a review. *J. Parkinson Dis* **5**, 21–41 (2015).
67. Pachman, D. R., Price, K. A. & Carey, E. C. Non-pharmacological approach to fatigue in cancer. *Cancer J.* **20**, 313–8 (2014).

This work is licensed under a Creative Commons Attribution 4.0 International License. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in the credit line; if the material is not included under the Creative Commons license, users will need to obtain permission from the license holder to reproduce the material. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>