CORRIGENDUM

Interaction between DRD2 and AKT1 genetic variations on risk of psychosis in cannabis users: a case–control study

Marco Colizzi, Conrad Iyegbe, John Powell, Giuseppe Blasi, Alessandro Bertolino, Robin M Murray and Marta Di Forti

npj Schizophrenia (2015) 1, Article number: 15049; doi:10.1038/npjschz.2015.49; published online 18 November 2015

Correction to: npj Schizophrenia (2015) 1, Article number: 15025. doi:10.1038/npjschz.2015.25In the published version of this article, in Table 1, under the entry 'Cannabis use', some *N* (%) data were incorrect due to a minor

miscalculation. The corrected table appears below. The authors regret the error.

Table 1. Demographic measures and patterns of drug use in FEP patients and control subjects				
	<i>PZ</i>		<i>Controls</i>	
	<i>N</i>	%	<i>N</i>	%
<i>Cannabis use</i>				
Yes	158	71.2	142	62.3
No risk allele	34	21.5	40	28.2
One risk allele	89	56.3	82	57.7
Two risk alleles	35	22.2	20	14.1
Never	64	28.8	86	37.7
No risk allele	20	31.2	17	19.8
One risk allele	35	54.7	51	59.3
Two risk alleles	9	14.1	18	20.9
<i>Frequency of use</i>				
Daily	92	63.9	43	39.8
No risk allele	19	20.7	12	27.9
One risk allele	51	55.4	22	51.2
Two risk alleles	22	23.9	9	20.9
Weekly or less	52	36.1	65	60.2
No risk allele	12	23.1	20	30.8
One risk allele	29	55.7	37	56.9
Two risk alleles	11	21.2	8	12.3
No details	14		34	