

H₃ Receptor Antagonism Enhances NCAM PSA-Mediated Plasticity and Improves Memory Consolidation in Odor Discrimination and Delayed Match-to-Position Paradigms

Andrew G Foley¹, Alison Prendergast¹, Claire Barry¹, Darren Scully¹, Neil Upton², Andrew D Medhurst² and Ciaran M Regan^{*,3}

¹Berand Neuropharmacology, NovaUCD, Belfield Innovation Park, University College Dublin, Dublin, Ireland; ²Neurosciences Centre of Excellence for Drug Discovery, GlaxoSmithKline, Harlow, UK; ³School of Biomolecular and Biomedical Science, UCD Conway Institute, University College Dublin, Belfield, Dublin, Ireland

To further understand the procognitive actions of GSK189254, a histamine H₃ receptor antagonist, we determined its influence on the modulation of hippocampal neural cell adhesion molecule (NCAM) polysialylation (PSA) state, a necessary neuroplastic mechanism for learning and memory consolidation. A 4-day treatment with GSK189254 significantly increased basal expression of dentate polysialylated cells in rats with the maximal effect being observed at 0.03–0.3 mg/kg. At the optimal dose (0.3 mg/kg), GSK189254 enhanced water maze learning and the associated transient increase in NCAM-polysialylated cells. The increase in dentate polysialylated cell frequency induced by GSK189254 was not attributable to enhanced neurogenesis, although it did induce a small, but significant, increase in the survival of these newborn cells. GSK189254 (0.3 mg/kg) was without effect on polysialylated cell frequency in the entorhinal and perirhinal cortex, but significantly increased the diffuse PSA staining observed in the anterior, ventromedial, and dorsomedial aspects of the hypothalamus. Consistent with its ability to enhance the learning-associated, post-training increases in NCAM PSA state, GSK189254 (0.3 mg/kg) reversed the amnesia induced by scopolamine given in the 6-h post-training period after training in an odor discrimination paradigm. Moreover, GSK189254 significantly improved the performance accuracy of a delayed match-to-position paradigm, a task dependent on the prefrontal cortex and degree of cortical arousal, the latter may be related to enhanced NCAM PSA-associated plasticity in the hypothalamus. The procognitive actions of H₃ antagonism combined with increased NCAM PSA expression may exert a disease-modifying action in conditions harboring fundamental deficits in NCAM-mediated neuroplasticity, such as schizophrenia and Alzheimer's disease.

Neuropsychopharmacology (2009) **34**, 2585–2600; doi:10.1038/npp.2009.89; published online 5 August 2009

Keywords: GSK189254; odor discrimination paradigm; delayed match-to-position paradigm; spatial learning; NCAM PSA; neurogenesis

INTRODUCTION

Histamine mediates diverse biological effects through the histamine H₁, H₂, H₃, and H₄ receptor subtypes (Hough, 2001). The H₃ receptor is widely expressed in the mammalian brain, particularly in regions associated with cognition and arousal, such as the cerebral cortex, hippocampus, and hypothalamus (Martinez-Mir *et al*, 1990; Pollard *et al*, 1993; Pillot *et al*, 2002). Activation of H₃ autoreceptors inhibits histamine synthesis and release, whereas activation of H₃ heteroreceptors inhibits the release of transmitters such as acetylcholine, noradrenaline, dopamine, and 5-HT from non-histaminergic neurons (Schlicker *et al*, 1994; Schlicker and Kathmann, 1998;

Brown *et al*, 2001). As a consequence, several strategies are being pursued for the development of selective histamine H₃ receptor antagonists that increase the release of neurotransmitters involved in cognitive processes, such as acetylcholine (Johnson *et al*, 2004; Witkin and Nelson, 2004). The first generation of imidazole-based molecules, including thioperamide (Arrang *et al*, 1987), have recently been superseded with the second generation of non-imidazole H₃ receptor antagonists (Hancock, 2003; Stark, 2003; Celanire *et al*, 2005; Leurs *et al*, 2005) and many of these exert significant cognition-enhancing actions in a variety of rodent models, including object recognition, olfactory recognition, water maze, radial maze, and passive avoidance conditioning paradigms (Esbenshade *et al*, 2003, 2004, 2005; Fox *et al*, 2003, 2005; Medhurst *et al*, 2007).

H₃ receptor antagonists, such as GSK189254 (6-((3-cyclobutyl-2,3,4,5-tetrahydro-1H-3-benzazepin-7-yl)oxy)-N-methyl-3-pyridinecarboxamide hydrochloride), ABT-239 (4-(2-(2-((2R)-2-methylpyrrolidinyl)ethyl)-benzofuran-5-yl)

*Correspondence: Professor CM Regan, School of Biomolecular and Biomedical Science, UCD Conway Institute, University College Dublin, Belfield, Dublin 4, Ireland, Tel: +353 1 7166775, Fax: +353 1 7166920, E-mail: Ciaran.Regan@ucd.ie
Received 3 April 2009; revised 27 May 2009; accepted 20 June 2009

benzonitrile) and ciproxifan, have also been shown to induce *c-fos* activation in cortical, hippocampal, and hypothalamic brain regions (Hancock *et al*, 2006; Medhurst *et al*, 2007), a neuroplastic event associated with processes of memory consolidation (Guzowski, 2002; Kaczmarek *et al*, 2002). GSK189254 can also reverse amnesia induced by scopolamine administered at 6 h after passive avoidance conditioning (Medhurst *et al*, 2007), a period associated with extensive synaptic remodeling during consolidation of avoidance conditioning and spatial learning paradigms within the rodent hippocampal dentate gyrus (O'Malley *et al*, 1998, 2000; Eyre *et al*, 2003). Such synaptic remodeling is accompanied by the activation of neural cell adhesion molecule (NCAM) polysialylation (PSA) state, a mechanism supporting structural plasticity in the adult nervous system (Bonfanti, 2006; Gascon *et al*, 2007; Rutishauser, 2008). Interestingly, NCAM PSA is significantly enhanced by chronic administration (~40 days) of cognition-enhancing drugs such as cholinesterase inhibitors (Murphy *et al*, 2006) and 5-HT₆ receptor antagonists (Foley *et al*, 2008).

In the adult brain, NCAM PSA is primarily associated with regions that undergo structural reorganization in response to physiological and/or behavioral stimuli, such as the hypothalamus and hippocampal formation (Bonfanti *et al*, 1992). The consolidation of many behavioral tasks is now known to require a transient increase in the frequency of polysialylated neurons, notably at the 12 h post-training time, in the hippocampal dentate gyrus (Fox *et al*, 1995 [84]Murphy *et al*, 1996; Foley *et al*, 2003a; Sandi *et al*, 2003, 2004) and associated entorhinal and perirhinal cortex (O'Connell *et al*, 1997; Fox *et al*, 2000). Moreover, disruption of PSA function, by intraventricular infusion endoneuraminidase-N or anti-PSA, impairs task consolidation (Becker *et al*, 1996; Muller *et al*, 1996; Venero *et al*, 2006; Lopez-Fernandez *et al*, 2007; Seymour *et al*, 2008). Many of the NCAM PSA-immunopositive cells located at the infragranular zone of the dentate gyrus have been identified as newly generated granule cells that remain available for integration into the neuronal architecture and before their natural loss by apoptosis (Ge *et al*, 2007; Dupret *et al*, 2007; Toni *et al*, 2008). Similarly, the activity-dependent modifications to hypothalamic synapse and astrocytic coverage of oxytocinergic neurons, which occur in response to physiological stimuli arising during parturition, lactation, or chronic dehydration, are dependent on NCAM PSA, as these are prevented by infusions of endoneuraminidase-N (Hoyk *et al*, 2001; Monlezun *et al*, 2005; Catheline *et al*, 2006; Theodosis *et al*, 2006).

To further explore the influence of H₃ antagonism on the processes of memory consolidation, we investigated the effects of GSK189254 and thioperamide on the modulation of NCAM PSA state, as this has been shown to underpin learning-associated synaptic remodeling, and is a necessary neuroplastic mechanism for memory and learning.

METHODS

Animal Maintenance

Naive male Wistar rats (postnatal day 80, ~350 gm) were employed in all studies. The animals were purpose bred at the Biomedical Facility, University College Dublin, and

maintained in standard laboratory conditions until the time of use. Animals were introduced to the experimental holding rooms 5 days before the commencement of the study, housed in pairs during this period, and maintained at 22–24°C on a standard 12-h light/dark cycle, with food and water available *ad libitum*. Food-restricted animals were maintained at approximately 90% of their free-feeding weight (~20 g/rat/day) with *ad libitum* access to water. For 2 days preceding the commencement of behavioral studies, the animals were handled, and weighed and assessed in an open field arena for locomotor activity, rearing, and general behavior over a 5-min period. All experimental procedures were approved by the Animal Research Ethics Committee of University College Dublin, conformed to EU Council Directive 86–609-EEC, and were carried out by individuals retaining the appropriate license issued by the Irish Department of Health.

Odor–Reward Association Paradigm

The training protocol employed has been described previously (Roulet *et al*, 1997; Foley *et al*, 2003a). Each individual animal was assigned a specific target odor that was always associated with the sponge that contained the food reward and the sponges were interchanged between the three corners of the training apparatus between trials to prevent spatial bias. Training was carried out in a single session of five trials. An inter-trial interval of 5 min was allowed between trials, and latency (s) was defined as the time taken to correctly identify the correct target odor and obtain the associated food reward. Animals were tested for recall in a single trial 24 and 72 h after training. Statistical analysis of the behavioral data employed two-way ANOVA and the Mann–Whitney *U*-test for non-parametric data. In each case, *p*-values less than 0.05 were considered to be significant.

Water Maze Spatial Learning Paradigm

The training protocol employed has been described previously (Murphy *et al*, 1996; Foley *et al*, 2004). Water maze training was initiated at 4 h after the final drug administration. During testing, the platform was hidden in the same quadrant 30 cm from the sidewall. Animals were trained in a single training session consisting of 5 trials, each separated by an inter-trial test interval of 300 s. Computerized tracking software (Watermaze 3.1, Labview written by Matthias Grossmann, Dresden, Germany) was used to track the swim path for each animal. The time taken by the rat to find the hidden platform within a 90-s criterion period was defined as the escape latency time. Escape latencies were measured over five trials in each training session. The effect of drug treatment and trial number on escape latency from the water maze was assessed by repeated measures ANOVA. Specific trials and probe differences were analyzed using Mann–Whitney *U*-test for non-parametric data. In all cases, *p*-values less than 0.05 were considered to be significant.

Delayed Match-to-Position Paradigm

The training paradigm employed was based on a modification of those previously described (Dunnett, 1985, 1993;

Kirkby *et al*, 1995; Sahgal, 1987). Postnatal day 80 Wistar rats were housed in pairs and autoshaped for lever pressing over a period of 40 days. Animals were required to remember which of two levers had previously been presented and select (press) this lever following a random delay period to record a correct response and receive food reinforcement. Animals were considered to have completed the autoshaping training once they had achieved a stable level of daily performance (>85% successful trials) over a period of 3 consecutive days. During training, a single lever was presented and, after defined delay periods of 4–32 s, two levers were introduced and selection of the correct lever resulted in a food reward. Animals were trained each day in 72 trials, 12 of each delay period pseudorandomly presented, over a period of 4 days using a block design that ensured each animal was administered each dose of the test compound and, thereby, served as its own control. Performance of the animals in the delayed match-to-position (DMTP) paradigm was expressed as percent correct lever press at increasing delay periods, and values significantly different between the vehicle- and drug-treated groups were analyzed by two-way ANOVA and the Mann-Whitney *U*-test for non-parametric data. In each case, *p*-values less than 0.05 were considered to be significant.

Quantitative Analysis of BrdU and NCAM PSA Expression

BrdU immunolabeling. After transcardial perfusion with a 4% paraformaldehyde solution (pH 7.4), brains were removed and snap-frozen. For the analysis of bromodeoxyuridine (BrdU)-immunopositive hippocampal dentate granule cells, horizontal sections (50 μ m) were taken at 500- μ m intervals between level 4.1 and 7.6 mm below bregma (Paxinos and Watson, 1986). The sections were then washed in 0.1 M phosphate-buffered saline (PBS) containing 5 mM MgCl₂ and 1 mM CaCl₂, and denatured at 37°C for 1 h in DNase (1000 U/ml). The sections were again washed and blocked with 10% (v/v) normal goat serum (NGS) for 30 min, then incubated for 20 h with the primary antibody (anti-rat IgG BrdU; Harlan, Bicester, UK) diluted 1:100 in PBS containing 10% (v/v) NGS. Subsequently, the sections were washed and incubated at room temperature for 1 h with the secondary antibody (Alexa488 or Alexa647 goat anti-rat IgG; Molecular Probes, Paisley, UK) diluted 1:200 again in PBS containing 10% NGS. The sections were again washed and mounted in Citifluor® (Agar, Essex, UK).

The total number of immunopositive cells within the entire granule cell layer of each section was determined by the Cavalieri method, as previously described (Mirescu *et al*, 2004). The average density of BrdU-labeled cells per granule cell layer at each bregma level was established and used to estimate the total number of BrdU-immunopositive cells in the hippocampus of each treatment group by multiplying the density of immunopositive cells by the estimated volume. Statistical analysis employed ANOVA followed by Bonferroni *post hoc* test and analysis by Student's *t*-test. For qualitative purposes, some sections from 4% paraformaldehyde-perfused brains were employed for double immunolabeling of NCAM PSA and BrdU in the hippocampal formation at a level –5.6 mm from bregma.

The same immunohistochemistry staining protocols were employed but without 70% ethanol post-fixation.

NCAM PSA immunolabeling. The immunohistochemical procedures employed to detect NCAM PSA have been described in greater detail previously (Fox *et al*, 1995). Cryosections were thaw-mounted onto glass slides, fixed for 30 min with 70% ethanol, and incubated overnight with anti-PSA ascitic fluid (generous gift of Prof G Rougon; Rougon *et al*, 1986) diluted 1:500 in 1 M PBS containing 1% (w/v) BSA and 1% (v/v) NGS. The sections were exposed for 1 h to Alexa488- or Alexa647-conjugated goat anti-mouse IgM (Molecular Probes) diluted 1:200 in PBS containing 1% BSA and 1% NGS, and mounted in Citifluor® (Agar). For the analysis of the NCAM PSA-positive hippocampal dentate granule cell layer/hilus border cells, 10 alternate sections were taken at a level equivalent to 5.6 mm below bregma, at which level this cell population was found to be maximal. The frequency of polysialylated neurons in the rat medial temporal lobe was examined in layer II of the entorhinal and perirhinal cortices at bregma levels –7.1, –7.6, –8.1, and –8.6 mm.

The total number of NCAM PSA-immunoreactive neurons on the right dentate granule cell layer/hilar border was counted in seven alternate 12- μ m sections, commencing –5.6 mm from bregma, to preclude double counting of the 5–10 μ m perikarya (*n* = 6). Cell counts were standardized to unit area of the granule cell layer, 0.15 ± 0.01 mm² at this level, and expressed as mean \pm SEM values. In the cortex, cell counts were standardized to unit length of the layer II band taken to be 10 mm (*n* = 3). Hypothalamic NCAM PSA immunostaining was evaluated in coronal cryostat sections (12 μ m), which were cut in the rostro-caudal plane at the level of 1.8 mm, 2.56, and 3.3 mm caudal to bregma. The average gray level intensity of PSA-immunopositive neurons was determined and standardized with respect to nonspecific background staining measured in the corpus callosum, and these values were used to generate mean \pm SEM values for each treatment group. Statistical analysis employed the Student's *t*-test and a significance level of *p* < 0.05 was employed in all cases.

Drug Administration Protocols

All studies were conducted using thioperamide and the hydrochloride salt of GSK189254 (6-((3-cyclobutyl-2,3,4,5-tetrahydro-1*H*-3-benzazepin-7-yl)oxy)-*N*-methyl-3-pyridinecarboxamide hydrochloride. Studies addressing the influence of GSK189254 (0.1–3 mg/kg) and thioperamide (10 mg/kg) on NCAM PSA expression employed animals that had received drug or vehicle (1% methylcellulose solution in dH₂O) by oral gavage once daily for a period of 4 days and were drug free at time of analysis 24 h after the final drug treatment. This 4-day dosing regime was selected as an intermediate protocol based on efficacy observed in cognition models after 1–8 days dosing. Analysis of repeat drug administration on spatial learning was carried out 2 h after the final dose, and the animals were culled for NCAM PSA expression 12 h later. In the protocol employed in analyzing the drug influence on neurogenesis and apoptosis, BrdU (50 mg/kg) was administered by the intraperitoneal route and GSK189254 (0.3 mg/kg) by oral gavage, and

these were separated by 1 h. A second BrdU injection was given 12 h later. One cohort was killed 24 h after the final injections, whereas another cohort was maintained drug free for a further 14 days to evaluate the effects on cell survival. In both the odor discrimination and DMTP tasks, GSK189254 or vehicle was administered acutely by oral gavage 2 h before training. In the odor discrimination task, scopolamine (0.8 mg/kg) was administered by intraperitoneal injection at 20 min before training.

RESULTS

NCAM PSA in the Hippocampal Dentate Gyrus is Enhanced by Repeat Administration of GSK189254

NCAM PSA expression in the adult dentate gyrus was found to be primarily associated with granule cell bodies located at the infragranular zone and their dendritic arbor that extended through the granular cell layer and into the molecular layer (Figure 1a). A 4-day treatment with GSK189254 significantly increased the frequency of these NCAM-polysialylated cells in a dose-dependent manner ($F(4,25) = 6.39$; $p < 0.05$; one-way ANOVA) (Figure 1b). The maximal effect of GSK189254 was observed at doses of 0.03–0.3 mg/kg ($p < 0.05$; Bonferroni *post hoc* analysis) as compared with the vehicle-treated groups, and the higher doses of 1 and 3 mg/kg were without significant effect on polysialylated cell frequency, indicating the drug effect to exhibit a bell-shaped response. Repeat administration of the H₃ receptor antagonist thioperamide for 4 days also significantly increased the frequency of polysialylated neurons (Figure 1c). The thioperamide and GSK189254 studies were carried out at different times and the small variation in baseline NCAM PSA-labeled cell frequency can be found under these circumstances. The effect of GSK189254 on NCAM PSA expression was dependent on repeat drug treatment, as a single administration of GSK189254 at the optimal dose (0.3 mg/kg) was without effect on dentate polysialylated cell frequency (vehicle: 169 ± 6.1 ; drug-treated: 174.0 ± 12.3 cells/unit area).

Spatial Learning-Induced Activation NCAM PSA State in the Dentate Gyrus is Augmented After Repeat Administration of GSK189254

Given that the frequency of dentate polysialylated neurons transiently increase at the 12 h post-training time during the consolidation of a variety of learning paradigms (Fox *et al*, 1995; Murphy *et al*, 1996; Foley *et al*, 2003a; Sandi *et al*, 2003), and that the extent of this increase is commensurate with complexity experienced in consolidating the task (Sandi *et al*, 2004; Murphy *et al*, 2006), we determined whether GSK189254-induced increases in basal polysialylated cell expression could be further enhanced at the 12 h post-training time after water maze training. Separate cohorts of animals were treated with either vehicle or GSK189254 (0.3 mg/kg) for 4 days and trained in the water maze paradigm at 2 h after the final drug treatment. All animals acquired the spatial learning task as indicated by the gradual reduction in latency to reach the hidden platform over the five trials of the single training session

($F(4,65) = 16.47$; $p < 0.0001$; 2-way ANOVA) (Figure 2a). Moreover, a further 2-way ANOVA indicated that a 4-day treatment with GSK189254 induced a significant improvement in the acquisition of the water maze task compared with vehicle-treated controls ($F(1,65) = 5.02$; $P = 0.0285$). Analysis of the hippocampal dentate gyrus in brain tissue collected from the vehicle-treated animals at 12 h after task acquisition revealed the expected increase in polysialylated cell frequency as compared with that in the tissue taken immediately after training ($p < 0.05$; Student's *t*-test) (Figure 2b). Similarly, sections taken from the GSK189254-treated cohort revealed a significant increase in dentate NCAM PSA immunoreactivity between animals killed immediately after training (0 h) and those killed at the 12 h post-training time ($p < 0.05$; Student's *t*-test). However, comparison of vehicle- and drug-treated groups at the 12 h post-training time also showed a significant increase in dentate polysialylated cell frequency ($p < 0.05$; Student's *t*-test), indicating that this learning-induced increase was not saturated and could be further increased during consolidation after administration of GSK189254.

Dentate Neurogenic Rate is Unaffected by GSK189254 but Cell Survival is Enhanced

Many of the NCAM PSA-immunopositive cells located at the infragranular zone of the dentate gyrus have been identified as newly generated granule cells (Gage, 2000; Seki, 2002). These newborn granule cell neurons in the dentate gyrus remain available for integration into the neuronal architecture for 1–2 months after their birth (Ge *et al*, 2007; Kee *et al*, 2007) and before their natural loss by the process of apoptosis (Dupret *et al*, 2007). As this process of cell generation and loss has been argued to provide a substrate for the synaptic remodeling associated with memory consolidation (Lledo *et al*, 2006), we determined whether GSK189254 influenced neurogenic rate in the hippocampal dentate gyrus. Neurogenic rate in the infragranular zone of the dentate gyrus was determined by estimating the number of BrdU-immunolabeled cells immediately after a 4-day treatment with BrdU, and the survival of labeled cells was similarly determined 2 weeks later. BrdU immunoreactivity was located to the cell nucleus and was clearly differentiated from the cell surface labeling of anti-NCAM PSA; however, not all BrdU-immunopositive cells were immunoreactive for NCAM PSA (Figure 3a). BrdU-immunopositive cells were quantified throughout the hippocampal dentate gyrus granule cell layer from a level –4.1 to –7.6 mm with respect to bregma and expressed as total cell number, as previously documented (Foley *et al*, 2008), and showed that no drug-induced modulation of BrdU-positive cell frequency was observed after 4-day administration of GSK 189254 (vehicle: 1325 ± 98.3 ; GSK189254: 1366.7 ± 69.8) (Figure 3b). However, after a 14-day period, during which animals received no additional drug treatments, the number of surviving BrdU-immunopositive cells was significantly increased in the hippocampal dentate gyrus of animals that had been treated with GSK189254 (vehicle: 265 ± 33.3 ; GSK189254: 491.7 ± 69.9 ; $p < 0.05$; Student's *t*-test) (Figure 3c).





Figure 1 The effect of GSK189254 on polysialylated cell frequency in the dentate gyrus of adult Wistar rats. The animals received GSK189254 (0.3 mg/kg, p.o.), thioperamide (10 mg/kg), or vehicle (1% methylcellulose (w/v)) for 4 days and were drug free at the time of killing 24 h after the final treatment. (a) Illustrates qualitative images of PSA immunoreactivity at low- (i-iii) and high-resolution (iv-vi) in the dentate granule cell layer (GCL) at -5.6 mm with respect to bregma. The arrowheads indicate the position of the immunostained cells at the infragranular zone and the scale bars represent 100 μ m. The quantitative, dose-dependent effects of repeat dosing with GSK189254 are shown in (b), and the effects of repeat dosing with thioperamide in (c). Data points represent the mean \pm SEM and group sizes are indicated within the columns. Values significantly different ($p < 0.05$) from the vehicle control are indicated with an asterisk.

NCAM PSA Expression in the Hippocampal Entorhinal and Perirhinal Cortex and Hypothalamus After Treatment with GSK189254

We also investigated the influence of GSK189254 on the expression of NCAM PSA state in brain regions other than the hippocampal dentate gyrus. The entorhinal cortex of the hippocampal formation was selected because previous studies had indicated the cognition-enhancing effects of 5-HT6

antagonists to be associated with a substantial increase in NCAM PSA expression in layer II of the entorhinal and perirhinal cortex (Foley *et al*, 2008). As reported previously (O'Connell *et al*, 1997; Fox *et al*, 2000), a band of PSA-immunopositive neurons was observed in the entorhinal and perirhinal cortex at 7.1 mm below bregma (Figure 4a), and their numerical density significantly increased at each descending level from 7.1 to 8.6 mm below bregma (Figure 4b). However, a 4-day treatment with GSK189254




Figure 2 Influence of repeat administration of GSK189254 on learning-induced activation of dentate polysialylated cell frequency in adult Wistar rats after spatial learning in the water maze paradigm. The animals received GSK189254 (0.3 mg/kg, p.o.) or vehicle (1% methylcellulose (w/v)), for 4 days before training and trained 2 h after the final drug treatment; NCAM PSA expression was analyzed 12 h post-training. (a) GSK189254 to induce a significant improvement of task acquisition in a single trial of five sessions ($n = 3-4$). The significant ($p < 0.05$), learning-induced increase in dentate polysialylated cell frequency between the 0 and 12 h post-training times in these animals is shown by the asterisks in (b). The cross indicates the significant difference in polysialylated cell frequency that exists between the vehicle- and drug-treated animals at the 0 and 12 h time.

(0.3 mg/kg), the dose found to optimally increase dentate polysialylated frequency, or thioperamide (10 mg/kg), had no effect on the dorso-ventral density and distribution of immunostained cells within each treatment group ($F(3,24) = 34.34$; $p < 0.0001$; two-way ANOVA) (Figure 4b).

Given that the pro-vigilant actions of H₃ antagonists could contribute, in part, to their procognitive actions (Le *et al*, 2008), we also analyzed the effect of GSK189254 treatment on NCAM PSA expression in the hypothalamus. Within this brain region, PSA immunoreactivity was diffuse in nature, as would be expected given its dominant association with glia (Theodosios *et al*, 1999). The areas examined for NCAM PSA immunoreactivity in the coronal sections, cut in the rostro-caudal direction at bregma levels 1.8, 2.56, and 3.3 mm, and their relationship to the same region in the rat stereological atlas (Paxinos and Watson, 1986), are shown in Figure 5. The absence of discrete immunostained cell bodies in the areas encompassing the anterior, ventromedial, and dorsomedial aspects of the hypothalamus necessitated the use of gray level analysis to quantify the effect of drug treatment on NCAM PSA expression. Using this procedure, 4-day administration of GSK189254 (0.3 mg/kg) was found to induce a significant and uniform increase in NCAM PSA immunoreactivity (Student's *t*-test, $p < 0.05$) (Figure 6).

GSK189254 Enhances Consolidation of an Odor Discrimination Paradigm

The finding that H₃ antagonism exerted a significant effect on dentate NCAM PSA state supports our previous study in

which GSK189254 was found to reverse the amnesia for avoidance conditioning task when induced by scopolamine administered at the 6 h post-training period of consolidation (Medhurst *et al*, 2007). Given that the training stimulus employed in such tasks is based on the avoidance of stressful experience (Merino *et al*, 2000), we further determined whether GSK189254 (0.3 mg/kg) exerted a similar effect in an odor discrimination paradigm in which the conditioning stimulus is stress free. Animals readily learned to acquire the odor discrimination paradigm, as the latency to locate the correct target odor became significantly reduced over the five trials of the training session. The latency to nose-poke in the sponge with the target odor decreased markedly from trial 1 to trial 2 and remained stable thereafter (Figure 7a). Acute administration of scopolamine alone, or in combination with GSK189254, had no significant effect on task acquisition, as judged by a two-way ANOVA analysis of the individual treatment groups in terms of escape latency times ($F(2,60) = 0.29$, $p = 0.75$). However, a similar analysis revealed all treatment groups to significantly decrease their latencies to locate the sponge containing the correct target odor over the five acquisition trials employed in the paradigm ($F(4,60) = 16.53$, $p < 0.0001$), indicating a rapid and robust acquisition of the task in all treatment cohorts.

Rats treated with vehicle alone showed good task recall at both the 24 and 72 h recall times. By contrast, acute scopolamine treatment alone was found to significantly impair task consolidation, as was evident by the significant increase in latency to locate the correct target odor at the 24 h recall time ($p < 0.05$; Mann-Whitney *U*-test)




Figure 3 Influence of repeat administration of GSK189254 on neurogenesis and cell survival in the adult dentate gyrus of the Wistar rat. BrdU- (red) and NCAM PSA-immunolabeled cells (green) at -5.6 mm to bregma in the dentate granule cell layer (GCL) of animals treated with vehicle (1% methylcellulose (w/v)) and GSK189254 (0.3 mg/kg, p.o.) are shown in (a) and (b), respectively. The animals were treated for 4 days and were drug free at the time of killing 24 h after final drug treatment. The BrdU-positive cells are indicated by the arrowheads and the NCAM PSA-positive cells by asterisks. The scale bar represents 100 μ m. (c) Illustrates the quantitative effect of the drug treatment on the frequency of BrdU- and NCAM PSA-labeled cells at postnatal day 80 (neurogenesis) and postnatal day 94 (cell survival). Data points represent the mean \pm SEM ($n = 3$) and values significantly different ($p < 0.05$) from the vehicle control are indicated with an asterisk.

(Figure 7b). A similar, but not statistically significant, trend to increased latency was also observed with scopolamine at the 72 h recall time. Acute administration of GSK189254 (0.3 mg/kg) significantly reversed the scopolamine-induced recall deficits observed at both the 24 and 72 h recall times ($p < 0.05$; Mann-Whitney *U*-test) (Figure 7b). These results suggested that the primary action of scopolamine in the odor discrimination paradigm emerges in the period of consolidation, as has been observed previously for an avoidance conditioning paradigm (Doyle and Regan, 1993), and that repeat administration of cognition-enhancing drugs, including H₃ antagonists, ameliorate this amnesic action of scopolamine (Foley *et al*, 2004, 2008; Medhurst *et al*, 2007).

Influence of Acute GSK189254 Administration on the DMTP Paradigm

All animals were trained in the DMTP paradigm for a period of 40 days and had achieved a stable level of performance at the chosen delays of 0, 4, 8, 16, 24, and 32 s. During training, baseline performance accuracy was approximately 80% in all groups at the shortest delay periods (0–8 s) but decreased as the delay period was lengthened to 32 s. Analysis of the vehicle-treated group revealed that choice accuracy decreased in a delay-dependent manner ($F(5,216) = 26.31$, $p < 0.0001$; 2-way ANOVA). Acute administration of GSK189254 (0.3 mg/kg) significantly improved

task performance accuracy ($F(1,108) = 6.04$, $p = 0.01$; 2-way ANOVA) (Figure 8); however, this was not observed after administration of the higher doses of 1 and 3 mg/kg, suggesting that the effect of H₃ antagonism on the DMTP task exhibited a bell-shaped response.

DISCUSSION

A major finding in this study was that a 4-day treatment with the H₃ receptor antagonist GSK189254 resulted in a dose-dependent increase in the frequency of polysialylated neurons in the infragranular zone of the hippocampal dentate gyrus. GSK189254 is a highly selective H₃ receptor antagonist that improves cognitive performance in a diverse range of cognition paradigms in rats (Medhurst *et al*, 2007). Our current data show for the first time that, amongst other transmitter systems, regulation of histaminergic function through H₃ receptor blockade can augment neuroplasticity mechanisms necessary for the effective consolidation of memory. Previous studies have shown other cognition-enhancing drugs, such as nefiracetam and tacrine, to increase NCAM PSA expression in the dentate gyrus (Murphy *et al*, 2006); however, their magnitude of effect was much less than that observed with GSK189254. Indeed, the ability of GSK189254 to increase the basal frequency of dentate polysialylated cells was more akin to that observed with a chronic 40-day administration of 5-HT₆ antagonists




Figure 4 Influence of repeat administration of GSK189254 and thioperamide on basal polysialylated cell frequency in layer II of the entorhinal cortex of adult Wistar rats. The animals received GSK189254 (0.3 mg/kg, p.o.), thioperamide (10 mg/kg, p.o.), or vehicle (1% methylcellulose (w/v)) for 4 days before training and were drug free at the time of killing 24 h after final drug treatment. (a) Qualitative images of PSA immunoreactivity after treatment with vehicle (i), GSK189254 (ii), and thioperamide (iii) in layer II of the entorhinal cortex (indicated with bracket). The arrowheads indicate the position of the pia and the scale bar represents 200 μ m. (b) Illustrates the quantitative effect of drug treatment on polysialylated cell frequency at increasing levels below bregma. Data points represent the mean \pm SEM ($n = 3$).

such as SB-271046 and SB-399885 (Foley *et al*, 2008), which suggests that the more recently developed procognitive agents tend to exert a more substantial impact on the neuroplastic mechanisms associated with memory consolidation.

The common ability of agents with differing primary modes of action to enhance NCAM PSA-mediated neuroplasticity remains, however, a complex issue. Glutamatergic excitation, driven by the NMDA receptor, is known to be necessary for the rapid decrease in NCAM PSA expression in the adult vagal complex after stimulation of its afferents (Bouzioukh *et al*, 2001), which suggests that post-translational glycosylation of NCAM is associated with periods of enhanced inhibition and/or neuronal quiescence. This is consistent with the marked downregulation of transcripts associated with neurotransmission, ion channel conductance, and signal transduction that is observed at the 12 h post-training period of memory consolidation when a transient increase in NCAM PSA expression is required

for memory consolidation (O'Sullivan *et al*, 2007). In contrast, however, the potent procognitive actions of NNC-711 (1-(2-((diphenylmethylene) amino) oxy) ethyl)-1,2,4,6-tetrahydro-3-pyridinecarboxylic acid hydrochloride), a GABA reuptake inhibitor with anticonvulsant activity (Suzdak *et al*, 1992), fails to influence polysialylated cell frequency, as does phenytoin, which lacks a procognitive action but dampens neural activity by slowing reactivation of voltage-dependent sodium channels (Rogawski and Porter, 1990; Murphy *et al*, 2006).

In general, the molecular events associated with regulation of the adult form of polysialyltransferase (ST8SiaIV/PST) remains to be fully elucidated; however, it would seem to be regulated by protein kinase C δ (PKC δ ; Gallagher *et al*, 2000, 2001), which, in turn, is influenced by complex cell-signaling mechanisms (Steinberg, 2004). Agents, such as curcumin and α -tocopherol, which induce the degradation of PKC δ , enhance PST activity and PSA of neurons in the infragranular zone of the dentate gyrus and the synaptic




Figure 5 Influence of repeat administration of GSK189254 on NCAM PSA expression in the hypothalamus of adult Wistar rats. PSA immunoreactivity in coronal sections, at an increasing rostro-caudal distance, containing the anterior, ventromedial, and dorsomedial areas of the hypothalamus is shown in (a), (b), and (c), respectively. The animals were treated with vehicle (1% methylcellulose (w/v)) or GSK189254 for 4 days and were drug free at the time of killing 24 h after final drug treatment. The panels show the precise brain region, located by reference to a rat brain atlas (Paxinos and Watson, 1986), and the delineated area used to determine gray level in the vehicle- and drug-treated animals.

remodeling of their dendritic arbor (Conboy *et al*, 2009; Ferri *et al*, 2006; Zingg and Azzi, 2004). The ability of GSK189254 to increase dentate polysialylated cell frequency may, however, be related to its ability to augment the outflow of acetylcholine, as two other cholinergic agents, tacrine and nefiracetam, similarly increase cholinergic drive (Nishizaki *et al*, 2000; Irizarry and Hyman, 2001; Medhurst *et al*, 2007) and all three agents enhance NCAM PSA in a bell-shaped dose-dependent manner (Murphy *et al*, 2006). By contrast, 5-HT₆ antagonists increase polysialylated cell frequency in a linear, dose-dependent manner over the range in which they elicit procognitive actions, but can increase not only acetylcholine release (Rogers and Hagan, 2001; Shirazi-Southall *et al*, 2002; Foley *et al*, 2008) but also

modulate excitatory amino acid neurotransmission (Dawson *et al*, 2001). The cause of the bell-shaped curve observed with GSK189254 on NCAM PSA is unclear. It may reflect the fact that H₃ antagonists can increase the release of multiple neurotransmitters in addition to acetylcholine, such as histamine, dopamine, and noradrenaline (Arrang *et al*, 1988; Fox *et al*, 2005; Medhurst *et al*, 2007), or it may be a consequence of the existence of numerous H₃ receptor splice variants (Hancock *et al*, 2003). The non-overlapping nature of the NCAM PSA bell-shaped dose-response curve with the effective doses in the passive avoidance and aged water maze paradigms may relate to a higher dose requirement be necessary to ameliorate memory loss in these deficit models (Medhurst *et al*, 2007).




Figure 6 Quantitative gray level analysis of NCAM PSA expression in the hypothalamus of adult Wistar rats after repeat administration of GSK189254. The animals were treated with vehicle (1% methylcellulose (w/v)) or GSK189254 (0.3 mg/kg, p.o.) for 4 days and were drug free at the time of killing 24 h after final drug treatment. Data points represent the mean \pm SEM ($n=4$) and values significantly different ($P<0.05$) from the vehicle control are indicated with an asterisk.




Figure 7 Influence of acute GSK189254 administration on the reversal of a scopolamine-induced learning impairment in the acquisition and retention phases of an odor discrimination paradigm in adult Wistar rats. The animals received GSK189254 (0.3 mg/kg, p.o.), or vehicle (1% methylcellulose (w/v)) at 2 h before training. The scopolamine (0.8 mg/kg, i.p.) was administered 20 min before the training. The influence of drug treatment on task acquisition and retention (recall) is shown in (a) and (b), respectively. Data points represent the mean \pm SEM ($n=6$), and values significantly different ($p<0.05$; two-way ANOVA and the Mann–Whitney U -test) from the vehicle control are indicated with an asterisk. Values significantly different between cohorts treated with scopolamine alone and scopolamine co-administered with GSK189254 are indicated with a cross ($p<0.05$; two-way ANOVA and the Mann–Whitney U -test).

The ability of GSK189254 to increase the basal frequency of dentate polysialylated neurons not only improved performance in the acquisition of a water spatial learning paradigm but also resulted in a significant enhancement of the transient increase in NCAM PSA expression at the 12 h post-training time. This has also been observed after separate treatments with tacrine and nefiracetam (Murphy *et al*, 2006), and the 5-HT₆ antagonists SB-271046 and SB-399885 (Foley *et al*, 2008). The improved learning

associated with increased NCAM PSA expression is not surprising given that the numerical frequency of polysialylated dentate neurons is directly correlated with task performance (Sandi *et al*, 2004; Murphy *et al*, 2006). The majority of newly synthesized PSA seems to be associated with the synapse-specific NCAM 180-kDa isoform (Doyle *et al*, 1992), and this would serve to reduce cell–cell signaling and facilitate synapse remodeling (Rutishauser, 2008), a suggestion reinforced by the learning deficits




Figure 8 Influence of acute GSK189254 administration on the delayed match-to-position paradigm in adult Wistar rats. The animals received acute GSK189254 by gavage, at doses of 0.3 mg/kg (a), 1 mg/kg (b), and 3 mg/kg (c) 2 h before training and compared with those receiving vehicle (1% methylcellulose (w/v)) alone. Data points represent the mean \pm SEM ($n = 10$) and values significantly different ($p < 0.05$; two-way ANOVA and the Mann–Whitney U -test) from the vehicle control are indicated with an asterisk.

observed in mice that are deficient for the adult form of polysialyltransferase (ST8SialIV/PST-1) (Markram *et al*, 2007) and the enhanced consolidation of this paradigm by post-training infusions of a cyclic oligopeptide that acts as a non-competitive agonist of PSA (Torregrossa *et al*, 2004; Florian *et al*, 2006). Moreover, facilitation of PSA activation by cognition-enhancing drugs, such as the H₃ receptor antagonist GSK189254, may have implications for the treatment of neurodegenerative conditions as, in Alzheimer's disease, the natural autoprotective response to age-related cognitive deficits is a small, but significant, activation of dentate polysialylated cell frequency (Mikkonen *et al*, 1999).

Many, but not all, of the NCAM PSA-immunopositive cells located at the infragranular zone of the dentate gyrus are newly generated granule cells (Gage, 2000; Seki, 2002) that are proposed to be involved in memory and learning (Gould *et al*, 1999; Shors *et al*, 2001; Dupret *et al*, 2007), and that their rate of production may be modulated by change in neurotransmitter status (Brezun and Daszuta, 2000; Malberg *et al*, 2000). However, in this study, GSK189254 was found to be without effect on neurogenic rate, despite producing marked increases in basal and learning-induced polysialylated cell frequency. This finding was not completely unexpected, as previous studies have consistently failed to implicate increased NCAM PSA-mediated plasticity with neurogenesis (Fox *et al*, 1995; Pham *et al*, 2005; Lopez-Fernandez *et al*, 2007; Duveau *et al*, 2007). Moreover, in a previous study, we have failed to associate chronic treatment with 5-HT₆ antagonists, with any alteration in hippocampal neurogenesis despite the significant increase in polysialylated cell frequency in both the hippocampal dentate gyrus and layer II of the medial temporal lobe (Foley *et al*, 2008), the latter being a brain region that does not sustain neurogenesis into adulthood (Ehninger and Kempermann, 2003). It is likely, however, that the activation of NCAM PSA state facilitates the integration of newborn cells into the dentate neuronal architecture over the 1- to 2-month period that follows their birth (Ge *et al*, 2007; Kee *et al*, 2007; Toni *et al*, 2008). In this respect, it was

interesting to observe that a 4-day treatment with GSK189254 increased the survival time of the newly formed dentate granule cells, as evidence by the significant increase in BrdU-labeled cells remaining at 2 weeks after the final BrdU injection. This would have the consequence of increasing the availability of newly formed cells for incorporation into the structure of the hippocampal formation in response to the neuroplastic demands necessary for information processing. Increased neural activity enhances cell survival (Bruehl-Jungerman *et al*, 2006) and their incorporation into the neural architecture is a complex multistep process that is associated with NCAM PSA expression (Schmidt-Hieber *et al*, 2004; Zhao *et al*, 2006; Toni *et al*, 2008). The ability of GSK189254 to prolong the half-life of these neuronal precursors, by reducing apoptosis, may also explain the general ability of H₃ receptor antagonists to protect against NMDA-induced cell death (Dai *et al*, 2007). PSA acts as a competitive antagonist of the extrasynaptic NR2B glutamate receptor subunit (Hammond *et al*, 2006), and its enhanced expression induced by GSK189254 administration may confer a neuroprotective quality by improving cell survival.

Unlike the hippocampus, we found no evidence for an effect of a 4-day treatment with GSK189254 on the frequency of polysialylated neurons in the rat entorhinal or perirhinal cortex, despite H₃ receptors being widely distributed in brain cortical regions (Pillot *et al*, 2002). This finding was surprising, as H₃ antagonists including GSK189254 are known to activate the intermediate early gene *c-fos* in both the prefrontal and somatosensory cortex (Medhurst *et al*, 2007; Southam *et al*, 2009), and cortical reorganization of sensory information is accompanied by increased glutamatergic drive and *c-fos* activation (Benali *et al*, 2008). In contrast, chronic treatment (40 days) with the 5-HT₆ receptor antagonists SB-271046 and SB-399885 markedly increase NCAM PSA expression in the entorhinal and perirhinal cortex (Foley *et al*, 2008), an effect that may relate to their ability to increase glutamate outflow (Dawson *et al*, 2001), as a similar action is not observed with GSK189254 (Medhurst *et al*, 2007). However, H₃ receptor

antagonism is also known to enhance *c-fos* mRNA expression in the supraoptic and paraventricular nuclei of the hypothalamus (Vizuete *et al*, 1995) and, in this brain region, we found GSK189254 to induce a widespread, uniform, and substantial increase in NCAM PSA expression. Modulation of vasopressin and oxytocin neurons in the hypothalamus is associated with an array of homeostatic functions (Hass *et al*, 2008), which are dependent on the activation of *c-fos* expression (Kjaer *et al*, 1994; Ma *et al*, 2008) and the NCAM PSA-mediated morphological plasticity that modifies astrocytic coverage of oxytocinergic neurons and their synaptic inputs in response to homeostatic functioning of the hypothalamic axis (Theodosis *et al*, 1999, 2006). Histamine acting on the hypothalamus affects the release of many hormones from the pituitary gland (Eriksson *et al*, 2001) that mediate the reaction to stress in the hypothalamic-pituitary axis, which may, in part, support the suggestion that H₃ receptor antagonists have some efficacy as antidepressants (Pérez-García *et al*, 1999). It is currently unclear why GSK189254 increased NCAM PSA in the hippocampus and hypothalamus, but not in cortical areas. This may be due to the involvement of different H₃ receptor splice variants in distinct brain areas (Hancock *et al*, 2003) or the potential heterogeneity of histaminergic nerve projections from the hypothalamus to different brain structures (Giannoni *et al*, 2007).

Given that H₃ antagonism has been associated with an increase in *c-fos* expression in the prefrontal cortex (Medhurst *et al*, 2007; Southam *et al*, 2009) and that modulation of NCAM PSA expression in this brain region is associated with learning-induced neuroplasticity (ter Horst *et al*, 2008), we determined whether GSK189254 influenced the DMTP task, as this paradigm specifically requires activation of the prefrontal cortex and not the hippocampus (Sloan *et al*, 2006). As the delay-dependent deficits increased in parallel with the load on working memory, GSK189254 was found to produce a significant improvement on task accuracy. This finding is of particular interest, as deficits in similar tasks in human subjects have been linked to psychiatric conditions, such as schizophrenia (Goldberg *et al*, 1987; Berman and Weinberger, 1990; Owen *et al*, 1995). Moreover, the ability of GSK189254 to increase cortical outflow of acetylcholine (Medhurst *et al*, 2007) along with the requirement of the cholinergic system in the DMTP task (Herremans *et al*, 1995) and the increase in acetylcholine release induced by atypical antipsychotics (Ichikawa *et al*, 2002) further implicates the role of the histaminergic system in schizophrenia (Arrang, 2007) and the potential of H₃ antagonists as novel antipsychotic agents (Southam *et al*, 2009). It is worth noting that a potential bell-shaped dose-response relationship occurred in the DMTP task similar to that observed in the NCAM PSA studies. However, this is not consistent with efficacy data in several other rodent cognition models with GSK189254 (Medhurst *et al*, 2007), and therefore may be specific to the DMTP paradigm.

Finally, the apparent importance of GSK189254 in the modulation of acetylcholine release and its relationship to NCAM PSA expression in learning was further underscored by its ability to reverse scopolamine-induced deficits in the odor discrimination paradigm, a task involving an increase in hippocampal NCAM PSA-mediated neuroplasticity

(Foley *et al*, 2003a; Knafo *et al*, 2005). The cholinergic projection neurons from the medial septum, which innervate the dentate granule cells, are extensively polysialylated (Foley *et al*, 2003b) and disruption of this pathway by lesions to the septal nuclei or fornix leads to impairments in rodent tasks of learning and memory (Everitt and Robbins, 1997). Moreover, rats trained in an odor discrimination paradigm exhibit a significant increase in *c-fos* expression in the CA1 and CA3 regions of the hippocampus but not in the dentate (Hess *et al*, 1995), suggesting that modulation in the PSA status of the mossy fiber afferents may induce *c-fos* expression in the trisynaptic pathway of the hippocampus.

These studies are the first to show the procognitive actions of an H₃ receptor antagonist to be associated with a profound activation of NCAM PSA state, a neuroplastic mechanism intimately associated with the synaptic remodeling that accompanies memory and learning. Improved expression of NCAM PSA, whether through the direct action of a PSA peptidomimetic (Florian *et al*, 2006) or indirectly by a receptor-mediated mechanism, not only exerts a profound effect on cognitive competence but is also associated with neuroprotective actions (Foley *et al*, 2005; Murphy *et al*, 2006; Duveau *et al*, 2007) that have the potential to provide a disease-modifying action in conditions that harbor fundamental deficits in NCAM PSA-mediated neuroplasticity, such as schizophrenia and Alzheimer's disease (Barbeau *et al*, 1995; Mikkonen *et al*, 1999).

DISCLOSURE/CONFLICT OF INTEREST

This work was supported by funding from GlaxoSmithKline. The authors NP and ADM are current employees of GlaxoSmithKline

REFERENCES

- Arrang JM (2007). Histamine and schizophrenia. *Int Rev Neurobiol* 78: 247–287.
- Arrang JM, Garbarg M, Lancelot JC, Lecomte JM, Pollard H, Robba M *et al* (1987). Highly potent and selective ligands for histamine H₃-receptors. *Nature* 327: 117–123.
- Arrang JM, Garbarg M, Lancelot JC, Lecomte JM, Pollard H, Robba M *et al* (1988). Potential interest in powerful and specific ligands for the histamine H₃ receptor. *Allerg Immunol (Paris)* 20: 327–329, 331.
- Barbeau D, Liang JJ, Robitaille Y, Quirion R, Srivastava LK (1995). Decreased expression of the embryonic form of the neural cell adhesion molecule in schizophrenic brains. *Proc Natl Acad Sci USA* 92: 2785–2789.
- Becker CG, Artola A, Gerardy-Schahn R, Becker T, Welzl H, Schachner M (1996). The polysialic acid modification of the neural cell adhesion molecule is involved in spatial learning and hippocampal long-term potentiation. *J Neurosci Res* 45: 143–152.
- Benali A, Weiler E, Benali Y, Dinse HR, Eysel UT (2008). Excitation and inhibition jointly regulate cortical reorganization in adult rats. *J Neurosci* 28: 12284–12293.
- Berman KF, Weinberger DR (1990). The prefrontal cortex in schizophrenia and other neuropsychiatric diseases: *in vivo* physiological correlates of cognitive deficits. *Prog Brain Res* 85: 521–536.
- Bonfanti L, Olive S, Poulain DA, Theodosis DT (1992). Mapping of the distribution of polysialylated neural cell adhesion molecule

- throughout the central nervous system of the adult rat: an immunohistochemical study. *Neuroscience* **49**: 419–436.
- Bonfanti L (2006). PSA-NCAM in mammalian structural plasticity and neurogenesis. *Prog Neurobiol* **80**: 129–164.
- Bouzioukh F, Tell F, Jean A, Rougon G (2001). NMDA receptor and nitric oxide synthase activation regulate polysialylated neural cell adhesion molecule expression in adult brainstem synapses. *J Neurosci* **21**: 4721–4730.
- Brezun JM, Daszuta A (2000). Serotonin may stimulate granule cell proliferation in the adult hippocampus, as observed in rats grafted with foetal raphe neurons. *Eur J Neurosci* **12**: 391–396.
- Brown RE, Stevens DR, Haas HL (2001). The physiology of brain histamine. *Prog Neurobiol* **63**: 637–672.
- Bruel-Jungerman E, Davis S, Rampon C, Laroche S (2006). Long-term potentiation enhances neurogenesis in the adult dentate gyrus. *J Neurosci* **26**: 5888–5893.
- Catheline G, Touquet B, Lombard M-C, Poulain DA, Theodosis DT (2006). A study of the role of neuro-glial remodelling in the oxytocin system at lactation. *Neuroscience* **137**: 309–316.
- Celanire S, Wijtmans M, Talaga P, Leurs R, de Esch IJ (2005). Keynote review: histamine H₃ receptor antagonists reach out for the clinic. *Drug Discov Today* **10**: 1613–1627.
- Conboy L, Foley AG, O'Boyle NM, Lawlor M, Gallagher HC, Murphy KJ et al (2009). Curcumin-induced degradation of PKC δ is associated with enhanced dentate NCAM PSA expression and spatial learning in adult and aged Wistar rats. *Biochem Pharmacol* **77**: 1254–1265.
- Dai H, Fu Q, Shen Y, Hu W, Zhang Z, Timmerman H et al (2007). The histamine H₃ receptor antagonist clobenpropit enhances GABA release to protect against NMDA-induced excitotoxicity through the cAMP/protein kinase A pathway in cultured cortical neurons. *Eur J Pharmacol* **563**: 117–123.
- Dawson L, Nguyen HQ, Li P (2001). The 5HT₆ receptor antagonist SB-271046 selectively enhances excitatory neurotransmission in the rat frontal cortex and hippocampus. *Neuropsychopharmacology* **25**: 662–668.
- Doyle E, Nolan PM, Bell R, Regan CM (1992). Hippocampal NCAM180 transiently increases sialylation during the acquisition and consolidation of a passive avoidance response in the adult rat. *J Neurosci Res* **31**: 513–523.
- Doyle E, Regan CM (1993). Cholinergic and dopaminergic agents which inhibit a passive avoidance response attenuate the paradigm-specific increases in NCAM sialylation state. *J Neural Transm* **92**: 33–49.
- Dunnett SB (1985). Comparative effects of cholinergic drugs and lesions of nucleus basalis or fimbria fornix on delayed matching in rats. *Psychopharmacology* **87**: 357–363.
- Dunnett SB (1993). Operant delayed matching and non-matching to position in rats. In: Sahgal A (ed). *Behavioural Neuroscience: A Practical Approach*, Oxford University Press, UK. Vol. I. pp 123–136.
- Dupret D, Fabre A, Dobrossy MD, Panatier A, Rodriguez JJ, Lamarque S et al (2007). Spatial learning depends on both the addition and removal of new hippocampal neurons. *PLoS Biol* **5**: e214.
- Duveau V, Arthaud S, Rougier A, Le Gal le Salle G (2007). Polysialylation of NCAM is upregulated by hyperthermia and participates in heat shock preconditioning-induced neuroprotection. *Neurobiol Disease* **26**: 385–395.
- Ehninger D, Kempermann G (2003). Regional effects of wheel running and environmental enrichment on cell genesis and microglia proliferation in adult murine neocortex. *Cereb Cortex* **13**: 845–851.
- Eriksson KS, Sergeeva O, Brown RE, Haas H (2001). Orexin/hypocretin excites the histaminergic neurons of the tuberomammillary nucleus. *J Neurosci* **21**: 9273–9279.
- Esbenshade TA, Krueger KM, Miller TR, Kang CH, Denny LI, Witte DG et al (2003). Two novel and selective nonimidazole histamine H₃ receptor antagonists A-304121 and A-317920. I. *In vitro* pharmacological effects. *J Pharmacol Exp Ther* **305**: 887–896.
- Esbenshade TA, Fox GB, Krueger KM, Baranowski JL, Miller TR, Kang CH et al (2004). Pharmacological and behavioral properties of A-349821, a selective and potent human histamine H₃ receptor antagonist. *Biochem Pharmacol* **68**: 933–945.
- Esbenshade TA, Fox GB, Krueger KM, Miller TR, Kang CH, Denny LI et al (2005). Pharmacological properties of ABT-239 [4-(2-{2-[(2R)-2-methylpyrrolidinyl]ethyl}-benzofuran-5-yl)benzotrile]. I. Potent and selective histamine H₃ receptor antagonist with drug-like properties. *J Pharmacol Exp Ther* **313**: 165–175.
- Everitt BJ, Robbins TW (1997). Central cholinergic systems and cognition. *Annu Rev Psychol* **48**: 649–684.
- Eyre MD, Richter-Levin G, Avital A, Stewart MG (2003). Morphological changes in the hippocampal dentate gyrus synapses following spatial learning in rats are transient. *Eur J Neurosci* **17**: 1973–1980.
- Ferri P, Cecchini T, Ambrogini P, Betti M, Cuppini R, Del Grande P et al (2006). α -Tocopherol affects neuronal plasticity in adult rat dentate gyrus: the possible role of PKC δ . *J Neurobiol* **66**: 793–810.
- Florian C, Foltz J, Norreel J-C, Rougon G, Rouillet P (2006). Post-training intrahippocampal injection of synthetic poly- α -2,8-sialic acid-neural cell adhesion molecule mimetic peptide improves spatial long-term performance in mice. *Learn Mem* **13**: 335–341.
- Foley AG, Hedigan K, Rouillet P, Sara SJ, Murphy KJ, Regan CM (2003a). Consolidation of odor-reward associative memory involves neural cell adhesion molecule polysialylation-mediated synaptic plasticity within the rodent hippocampus. *J Neurosci Res* **74**: 570–576.
- Foley AG, Rønn LCR, Murphy KJ, Regan CM (2003b). Distribution of polysialylated neural cell adhesion molecule in the rat septal nuclei and septohippocampal pathway: transient increase of polysialylated cells in sub-triangular septal zone during memory consolidation. *J Neurosci Res* **74**: 807–817.
- Foley AG, Murphy KJ, Hirst WD, Gallagher HC, Hagan JJ, Upton N et al (2004). The 5-HT₆ receptor antagonist SB-271046 reverses scopolamine-disrupted consolidation of a passive avoidance task and ameliorates spatial task deficits in aged rats. *Neuropsychopharmacol* **29**: 93–100.
- Foley AG, Murphy KJ, Regan CM (2005). Complex environment rearing prevents prenatal hypoxia-induced deficits in hippocampal cellular mechanisms necessary for memory consolidation in the adult Wistar rat. *J Neurosci Res* **82**: 245–254.
- Foley AG, Hirst WD, Gallagher HG, Barry C, Hagan JJ, Upton N et al (2008). The selective 5-HT₆ receptor antagonists SB-271046 and SB-399885 potentiate NCAM-PSA immunolabelling of dentate granule cells, but not neurogenesis, in the hippocampal formation of mature Wistar rats. *Neuropharmacology* **54**: 1166–1174.
- Fox GB, O'Connell AW, Murphy KJ, Regan CM (1995). Memory consolidation induces a transient and time-dependent increase in the frequency of NCAM-polysialylated cells in the adult rat hippocampus. *J Neurochem* **65**: 2796–2799.
- Fox GB, Fichera G, Barry T, O'Connell AW, Murphy KJ, Regan CM (2000). Consolidation of passive avoidance learning is associated with transiently increased polysialylated neuron number in layer II of the rat medial temporal cortex. *J Neurobiol* **45**: 135–141.
- Fox GB, Pan JB, Radek RJ, Lewis AM, Bitner RS, Esbenshade TA et al (2003). Two novel and selective nonimidazole H₃ receptor antagonists A-304121 and A-317920: II. *In vivo* behavioral and neurophysiological characterization. *J Pharmacol Exp Ther* **305**: 897–908.
- Fox GB, Esbenshade TA, Pan JB, Radek RJ, Krueger KM, Yao BB et al (2005). Pharmacological properties of ABT-239 [4-(2-{2-[(2R)-2-methylpyrrolidinyl]ethyl}-benzofuran-5-yl)benzotrile].

- II. Neurophysiological characterization and broad preclinical efficacy in cognition and schizophrenia of a potent and selective histamine H₃ receptor antagonist. *J Pharmacol Exp Ther* 313: 176–190.
- Gage FH (2000). Mammalian neural stem cells. *Science* 287: 1433–1438.
- Gallagher HC, Odumeru OA, Regan CM (2000). Regulation of neural cell adhesion molecule polysialylation state by cell-cell contact and protein kinase C delta. *J Neurosci Res* 61: 636–645.
- Gallagher HC, Murphy KJ, Foley AG, Regan CM (2001). Protein kinase C delta regulates neural cell adhesion molecule polysialylation state in the rat brain. *J Neurochem* 77: 425–434.
- Gascon E, Vutskits L, Kiss JZ (2007). Polysialic acid-neural cell adhesion molecule in brain plasticity: from synapses to integration of new neurons. *Brain Res Rev* 56: 101–118.
- Ge S, Yang C-h, Hsu K-s, Ming G-l, Song H (2007). A critical period for enhanced synaptic plasticity in newly generated neurons of the adult brain. *Neuron* 54: 559–566.
- Giannoni P, Passani MB, Nosi D, Medhurst AD, Chazot P, Shenton F et al (2007). Detection of functional heterogeneity of histaminergic neurons in response to GSK189254, a novel H₃ receptor antagonist. XXXVI Annual Conference of European Histamine Research Society, Florence, May 9–12th 2007.
- Goldberg TE, Weinberger DR, Berman KF, Pliskin NH, Podd MH (1987). Further evidence for dementia of the prefrontal type in schizophrenia? A controlled study of teaching the Wisconsin Card Sorting Test. *Arch Gen Psychiat* 44: 1008–1014.
- Gould E, Beylin A, Tanapat P, Reeves A, Shors TJ (1999). Learning enhances adult neurogenesis in the hippocampal formation. *Nat Neurosci* 2: 260–265.
- Guzowski JF (2002). Insights into immediate-early gene function in hippocampal memory consolidation using antisense oligonucleotide and fluorescent imaging approaches. *Hippocampus* 12: 86–104.
- Hancock AA (2003). H₃ receptor antagonists/inverse agonists as anti-obesity agents. *Curr Opin Investig Drugs* 4: 1190–1197.
- Hancock AA, Esbenshade TA, Krueger KM, Yao BB (2003). Genetic and pharmacological aspect of histamine H₃ receptor heterogeneity. *Life Sci* 73: 3043–3072.
- Hancock AA, Blitner RS, Krueger KM, Otte S, Nikkel AL, Fey TA et al (2006). Distinctions and contradistinctions between antiobesity histamine H₃ receptor (H₃R) antagonists compared to cognition-enhancing H₃ receptor antagonists. *Inflamm Res* 55(Suppl 1): S42–S44.
- Hammond MSL, Sims C, Parameshwaran K, Suppiramaniam V, Schachner M, Dityatev A (2006). Neural cell adhesion molecule-associated polysialic acid inhibits NR2B-containing N-methyl-D-aspartate receptors and prevents glutamate-induced cell death. *J Biol Chem* 281: 34859–34869.
- Hass HL, Sergeeva OA, Selbach O (2008). Histamine in the nervous system. *Physiol Rev* 88: 1183–1241.
- Herremans AH, Hijzen TH, Welborn PF, Olivier B, Slangen JL (1995). Effects of infusion of cholinergic drugs into the prefrontal cortex area on delayed matching to position performance in the rat. *Brain Res* 711: 102–111.
- Hess US, Lynch G, Gall CM (1995). Changes in c-fos mRNA expression in rat brain during odor discrimination learning: differential involvement of hippocampal subfields CA1 and CA3. *J Neurosci* 15: 4786–4795.
- ter Horst JPF, Loscher JS, Pickering M, Regan CM, Murphy KJ (2008). Learning-associated regulation of polysialylated neural cell adhesion molecule expression in the rodent prefrontal cortex is region-, cell type- and paradigm-specific. *Eur J Neurosci* 28: 419–427.
- Hough LB (2001). Genomics meets histamine receptors: new subtypes, new receptors. *Mol Pharmacol* 59: 415–419.
- Hoyle Z, Parducz A, Theodosis DT (2001). The highly sialylated isoform of the neural cell adhesion molecule is required for estradiol-induced morphological plasticity in the adult arcuate nucleus. *Eur J Neurosci* 13: 649–656.
- Ichikawa J, Dai J, O’Laughlin IA, Fowler WL, Meltzer HY (2002). Atypical, but not typical, antipsychotic drugs increase cortical acetylcholine release without an effect in the nucleus accumbens or striatum. *Neuropsychopharmacology* 26: 325–339.
- Irizarry MC, Hyman BT (2001). Alzheimer disease therapeutics. *J Neuropathol Exp Neurol* 60: 923–928.
- Johnson CN, Roland A, Upton N (2004). New symptomatic strategies in Alzheimer’s disease. *Drug Discov Today* 1: 13–19.
- Kaczmarek L, Lapinska-Dzwonek J, Szymczak S (2002). Matrix metalloproteinases in the adult brain physiology: a link between c-Fos, AP-1 and remodeling of neuronal connections? *EMBO J* 21: 6643–6648.
- Kee N, Teixeira CM, Wang AH, Frankland PW (2007). Preferential incorporation of adult-generated granule cells into spatial memory networks in the dentate gyrus. *Nat Neurosci* 10: 355–362.
- Kirkby DL, Jones DNC, Higgins GA (1995). Influence of prefeeding and scopolamine upon performance in a delayed-matching-to-position task. *Behav Brain Res* 67: 221–227.
- Kjaer A, Larsen PJ, Knigge U, Moller M, Warberg J (1994). Histamine stimulates c-fos expression in hypothalamic vasopressin-, oxytocin-, corticotropin-releasing hormone-containing neurons. *Endocrinology* 134: 482–491.
- Knafo S, Barkai E, Herrero AI, Libersat F, Sandi C, Venero C (2005). Olfactory learning-related NCAM expression is state, time, and location specific and is correlated with individual learning capabilities. *Hippocampus* 15: 316–325.
- Le S, Gruner JA, Mathiasen JR, Marino MJ, Schaffhauser H (2008). Correlation between *ex vivo* receptor occupancy and wake-promoting activity of selective H₃ receptor antagonists. *J Pharm Exp Ther* 325: 902–909.
- Leurs R, Bakker RA, Timmerman H, de Esch IJ (2005). The histamine H₃ receptor: from gene cloning to H₃ receptor drugs. *Nat Rev Drug Discov* 4: 107–120.
- Lledo PM, Alonso M, Grubb MS (2006). Adult neurogenesis and functional plasticity in neural circuits. *Nat Rev Neurosci* 7: 179–193.
- Lopez-Fernandez MA, Montaron M-F, Varea E, Rougon G, Venero C, Abrous DN et al (2007). Upregulation of polysialylated neural cell adhesion molecule in the dorsal hippocampus after contextual fear conditioning is involved in long-term memory formation. *J Neurosci* 27: 4552–4561.
- Ma S, Mifflin SW, Cunningham JT, Morilak JT (2008). Chronic intermittent hypoxia sensitizes acute hypothalamic-pituitary-adrenal stress reactivity and Fos induction in the rat locus coeruleus in response to subsequent immobilization stress. *Neuroscience* 154: 1639–1647.
- Malberg J, Eisch AJ, Nestler EJ, Duman RS (2000). Chronic antidepressant treatment increases neurogenesis in adult hippocampus. *J Neurosci* 20: 9104–9110.
- Markram K, Gerardy-Schahn R, Sandi C (2007). Selective learning and memory impairments in mice deficient for polysialylated NCAM in adulthood. *Neuroscience* 144: 788–796.
- Martinez-Mir MI, Pollard H, Moreau J, Arrang JM, Ruat M, Traiffort E et al (1990). Three histamine receptors (H₁, H₂ and H₃) visualised in the brain of human and non-human primates. *Brain Res* 526: 322–327.
- Medhurst AD, Atkins AR, Beresford IJ, Brackenborough K, Michael Briggs MA, Calver AR et al (2007). GSK189254 - a novel H₃ receptor antagonist that binds to histamine H₃ receptors in Alzheimer’s Disease brain and improves cognitive performance in preclinical models. *J Pharmacol Exp Ther* 321: 1032–1045.

- Mikkonen M, Soininen H, Tapiola T, Alafuzoff I, Miettinen R (1999). Hippocampal plasticity in Alzheimer's disease: changes in highly polysialylated NCAM immunoreactivity in the hippocampal formation. *Eur J Neurosci* 11: 1754–1764.
- Merino JJ, Cordero MI, Sandi C (2000). Regulation of hippocampal cell adhesion molecules NCAM and L1 by contextual fear conditioning is dependent upon time and stressor intensity. *Eur J Neurosci* 12: 3283–3290.
- Mirescu C, Peters JD, Gould E (2004). Early life experience alters response of adult neurogenesis to stress. *Nature Neurosci* 7: 841–846.
- Monlezun S, Ouali S, Poulain DA, Theodosis DT (2005). Polysialic acid is required for active phases of morphological plasticity of neurosecretory axons and their glia. *Mol Cell Neurosci* 29: 516–524.
- Muller D, Wang C, Skibo G, Toni N, Cremer H, Calaora V et al (1996). PSA-NCAM is required for activity-induced synaptic plasticity. *Neuron* 17: 413–422.
- Murphy KJ, O'Connell AW, Regan CM (1996). Repetitive and transient increase in hippocampal neural cell adhesion molecule polysialylation state following multi-trial spatial learning. *J Neurochem* 67: 1268–1274.
- Murphy KJ, Foley AG, O'Connell AW, Regan CM (2006). Chronic exposure of rats to cognition enhancing drugs produces a neuroplastic response identical to that obtained by complex environment rearing. *Neuropsychopharmacology* 31: 90–100.
- Nishizaki T, Nomura T, Matuoka T, Kondoh T, Enikolopo G, Sumikawa K et al (2000). The anti-dementia drug nefiracetam facilitates hippocampal synaptic transmission by functionally targeting presynaptic nicotinic ACh receptors. *Brain Res Mol Brain Res* 80: 53–62.
- O'Connell AW, Fox GB, Murphy KJ, Fichera G, Kelly J, Regan CM (1997). Spatial learning activates neural cell adhesion molecule polysialylation in a cortico-hippocampal pathway within the medial temporal lobe. *J Neurochem* 68: 2538–2546.
- O'Malley A, O'Connell C, Regan CM (1998). Ultrastructural analysis reveals avoidance conditioning to induce a transient increase in hippocampal dentate spine density in the 6h post-training period of consolidation. *Neuroscience* 87: 607–613.
- O'Malley A, O'Connell C, Murphy KJ, Regan CM (2000). Transient spine density increases in the mid-molecular layer of the hippocampal dentate gyrus accompany consolidation of a spatial learning task in the rodent. *Neuroscience* 99: 229–232.
- O'Sullivan NC, McGettigan PA, Sheridan GK, Pickering M, Conboy L, O'Connor JJ et al (2007). Temporal change in gene expression in the rat dentate gyrus following passive avoidance learning. *J Neurochem* 101: 1085–1098.
- Owen AM, Sahakian BJ, Semple J, Polkey CE, Robbins TW (1995). Visuo-spatial short-term recognition memory and learning after temporal lobe excisions, frontal lobe excisions, or amygdalo-hippocampectomy in man. *Neuropsychologia* 33: 1–24.
- Paxinos G, Watson C (1986). *The Rat Brain in Stereotaxic Coordinates* 2nd edn. Academic Press: New York.
- Pérez-García C, Morales L, Cano MV, Sancho I, Alguacil LF (1999). Effects of histamine H₃ receptor ligands in experimental models of anxiety and depression. *Psychopharmacology* 142: 215–220.
- Pham K, McEwen BS, LeDoux JE, Nader K (2005). Fear learning transiently impairs hippocampal cell proliferation. *Neuroscience* 130: 17–24.
- Pillot C, Heron A, Cochois V, Tardivel-Lacombe J, Ligneau X, Schwartz JC et al (2002). A detailed mapping of the histamine H₃ receptor and its gene transcripts in rat brain. *Neuroscience* 114: 173–193.
- Pollard H, Moreau J, Arrang JM, Schwartz JC (1993). A detailed autoradiographic mapping of histamine H₃ receptors in rat brain areas. *Neuroscience* 52: 169–189.
- Rogawski MA, Porter RJ (1990). Antiepileptic drugs: pharmacological mechanisms and clinical efficacy with consideration of promising developmental stage compounds. *Pharmacol Rev* 42: 223–286.
- Rougon G, Dubois C, Buckley N, Magnani JL, Zollinger W (1986). A monoclonal antibody against meningococcus group B polysaccharides distinguishes embryonic from adult N-CAM. *J Cell Biol* 103: 2429–2437.
- Roullet P, Mileusnic R, Rose SP, Sara SJ (1997). Neural cell adhesion molecules play a role in rat memory formation in appetitive as well as aversive tasks. *NeuroReport* 8: 1907–1911.
- Rogers DC, Hagan JJ (2001). 5-HT₆ receptor antagonists enhance retention of a water maze task in the rat. *Psychopharmacology* 158: 114–119.
- Rutishauser U (2008). Polysialic acid in the plasticity of the developing and adult vertebrate nervous system. *Nat Rev Neurosci* 9: 26–35.
- Sandi C, Merino JJ, Cordero MI, Krudt ND, Murphy KJ, Regan CM (2003). Modulation of hippocampal NCAM polysialylation and spatial memory consolidation by fear conditioning. *Biol Psychiatry* 54: 599–607.
- Sandi C, Cordero MI, Merino JJ, Krudt ND, Regan CM, Murphy KJ (2004). Neurobiological and endocrine correlates of individual differences in spatial learning ability. *Learn Mem* 11: 244–252.
- Sahgal A (1987). Contrasting effects of vasopressin, desglycinamide-vasopressin and amphetamine on a delayed matching to position task in rats. *Psychopharmacology* 93: 243–249.
- Schlicker E, Kathmann M (1998). Modulation of *in vitro* neurotransmission in the CNS and in the retina via H₃ heteroreceptors. In: Leurs R and Timmerman H (eds). *The Histamine H₃ Receptor: A Target for New Drugs* pp 13–26, Elsevier: Amsterdam, The Netherlands.
- Schlicker E, Malinowska B, Kathmann M, Gothert M (1994). Modulation of neurotransmitter release via histamine H₃ heteroreceptors. *Fundam Clin Pharmacol* 8: 128–137.
- Schmidt-Hieber C, Jonas P, Bischofberger J (2004). Enhanced synaptic plasticity in newly generated granule cells in the adult hippocampus. *Nature* 429: 184–187.
- Seki T (2002). Hippocampal neurogenesis occurs in a microenvironment provided by PSA-NCAM-expressing immature neurons. *J Neurosci Res* 69: 772–783.
- Seymour CM, Foley AG, Murphy KJ, Regan CM (2008). Intraventricular infusions of anti-NCAM PSA impair the process of consolidation of both avoidance conditioning and spatial learning paradigms in Wistar rats. *Neuroscience* 157: 813–820.
- Shors TJ, Miesegaes G, Beylin A, Zhao M, Rydel T, Gould E (2001). Neurogenesis in the adult is involved in the formation of trace memories. *Nature* 410: 372–376.
- Shirazi-Southall S, Rodriguez DE, Nomikos GG (2002). Effects of typical and atypical antipsychotics and receptor selective compounds on acetylcholine efflux in the hippocampus of the rat. *Neuropsychopharmacology* 26: 583–594.
- Sloan HL, Good M, Dunnett SB (2006). Double dissociation between hippocampal and prefrontal lesions on an operant delayed matching task and a water maze reference memory task. *Behav Brain Res* 171: 116–126.
- Stark H (2003). Recent advances in histamine H₃/H₄ receptor ligands. *Expert Opin Ther Pat* 13: 851–865.
- Southam E, Cilia J, Gartlon JE, Woolley ML, Lacroix LP, Jennings CA et al (2009). Preclinical investigations into the antipsychotic potential of the novel histamine H₃ receptor antagonist GSK207040. *Psychopharmacol* 201: 483–494.
- Steinberg SF (2004). Distinctive activation mechanisms and functions for protein kinase Cdelta. *Biochem J* 384: 449–459.
- Suzdak PD, Frederiksen K, Andersen KE, Sørensen PO, Knutsen LJS, Nielsen EB (1992). NNC-711, a novel potent and selective γ -aminobutyric acid uptake inhibitor: pharmacological characterization. *Eur J Pharmacol* 223: 189–198.
- Theodosis DT, Bonhomme R, Vitiello S, Rougon G, Poulain DA (1999). Cell surface expression of polysialic acid on

- NCAM is a prerequisite for activity-dependent morphological neuronal and glial plasticity. *J Neurosci* **19**: 10228–10236.
- Theodosis DT, Traillin A, Poulain DA (2006). Remodelling of astrocytes, a prerequisite for synapse turnover in the adult brain? Insights from the oxytocin system of the hypothalamus. *Am J Physiol Regul Comp Physiol* **290**: R1175–R1182.
- Toni N, Laplagne DA, Zhao C, Lombardi G, Ribak CE, Gage FH et al (2008). Neurons born in the adult dentate gyrus form functional synapses with target cells. *Nature Neurosci* **11**: 901–907.
- Torregrossa P, Buhl L, Bancila M, Durbec P, Schafer C, Schachner M et al (2004). Selection of poly- α 2,8-sialic acid mimotopes from a random phage peptide library and analysis of their bioactivity. *J Biol Chem* **279**: 30707–30714.
- Venero C, Herrero AI, Touyarot K, Cambon K, López-Fernández MA, Berezin V et al (2006). Hippocampal up-regulation of NCAM expression and polysialylation plays a key role on spatial memory. *Eur J Neurosci* **23**: 1585–1595.
- Vizuete ML, Dimitriadou V, Traiffort E, Griffon N, Heron A, Schwartz JC (1995). Endogenous histamine induces *c-fos* expression within paraventricular and supraoptic nuclei. *Neuroreport* **6**: 1041–1044.
- Witkin JM, Nelson DL (2004). Selective histamine H₃ receptor antagonists for treatment of cognitive deficiencies and other disorders of the central nervous system. *Pharmacol Ther* **103**: 1–20.
- Zhao C, Teng EM, Summers Jr RG, Ming GL, Gage FH (2006). Distinct morphological stages of dentate granule neuron maturation in the adult mouse hippocampus. *J Neurosci* **26**: 3–11.
- Zingg JM, Azzi A (2004). Non-antioxidant activities of vitamin E. *Curr Med Chem* **11**: 1113–1133.