

DATI and COMT Effects on Delay Discounting and Trait Impulsivity in Male Adolescents with Attention Deficit/Hyperactivity Disorder and Healthy Controls

Yannis Paloyelis^{*1}, Philip Asherson¹, Mitul A Mehta², Stephen V Faraone³ and Jonna Kuntsi¹

¹MRC SDGP Centre, Institute of Psychiatry, King's College London, London, UK; ²Centre for Neuroimaging Sciences, Institute of Psychiatry, King's College London, London, UK; ³Department of Psychiatry and of Neuroscience and Physiology, SUNY Upstate Medical University, Syracuse, NY, USA

Choice impulsivity has been linked to dopamine function and is consistently observed in attention deficit/hyperactivity disorder (ADHD) as a preference for smaller-immediate over larger-delayed rewards using choice-delay paradigms. More sophisticated delay discounting paradigms have yielded inconsistent results. Context and sample characteristics may have contributed to these variations. In this study we examine the effect of type (real vs hypothetical) and magnitude of reward as well as of variation in dopamine genes on choice impulsivity. We selected 36 male adolescents with ADHD-combined subtype (ADHD-CT) and 32 controls (mean age = 15.42, SD = 2.05) to form four roughly equally sized subgroups on the basis of *DATI*₁₀₁₆ haplotype dosage (2 copies and <2 copies). Participants, who were also genotyped for the *COMT*_{Val158Met} and *DRD4*_{48bp-VNTR} polymorphisms, performed a hypothetical and a real-time discounting task and provided self-ratings of trait impulsivity. The ADHD-CT group discounted rewards more steeply than controls only in the hypothetical task, with delay, but not reward magnitude, influencing choices. They also rated themselves as more impulsive compared with controls. *DATI*₁₀₁₆ dosage and the *COMT*_{Val158Met} genotype predicted trait impulsivity and discounting rates in the hypothetical task, but not in the real-time task. Our results directly link variation in genes putatively influencing dopamine signaling in the prefrontal cortex (*COMT*_{Val158Met}) and the striatum (*DATI*₁₀₁₆) with discounting rates in a hypothetical task (but not a real-time task) and self-ratings of trait impulsivity in ADHD-CT and healthy controls. The lack of magnitude effects in the hypothetical task suggests that discounting in this task may be influenced by different processes in ADHD-CT than in healthy controls.

Neuropsychopharmacology (2010) 35, 2414–2426; doi:10.1038/npp.2010.124; published online 25 August 2010

Keywords: ADHD; delay discounting; impulsivity; dopamine; DAT1; COMT

INTRODUCTION

Impulsive behavior is a key characteristic of several psychiatric conditions, including attention deficit/hyperactivity disorder (ADHD) (Moeller *et al*, 2001). One aspect of impulsivity is the preference for immediate gratification, even when waiting longer might lead to higher absolute gains (Evenden, 1999). The decrease in the subjective value of a reinforcer with time is called delay discounting and is observed in many species (Ainslie, 1975; Rachlin and Green, 1972). In humans, delay discounting is usually assessed through choices between hypothetical gains that systematically vary in magnitude or in the delay one needs to wait to receive them (Richards *et al*, 1999). The pattern of

responses, determined by the rate at which an individual discounts potential gains as a function of time, is mathematically best described by a hyperbolic-like curve (Green and Myerson, 2004; Reynolds, 2006b). A steeper curve indicates a higher discounting rate, reflecting greater impulsivity or reduced self-control (Logue, 1988). Populations with impulse control problems consistently demonstrate higher discounting rates compared with healthy controls (see Green and Myerson, 2004; Reynolds, 2006b for reviews), and prospective studies suggest that high discounting rates may indicate vulnerability for the development of such problems across species (eg, substance abuse/dependence; Anker *et al*, 2009; Audrain-McGovern *et al*, 2009; Dandy and Gatch, 2009; Diergaarde *et al*, 2008; Perry *et al*, 2005; Perry *et al*, 2008; Poulos *et al*, 1995; Wilhelm and Mitchell, 2009).

Different lines of evidence have linked delay discounting to dopamine and the function of dopamine-modulated frontostriatal circuits (Adriani *et al*, 2009; Cardinal *et al*, 2001; Kable and Glimcher, 2009; Kobayashi and Schultz, 2008;

*Correspondence: Y Paloyelis, MRC SDGP Centre, Institute of Psychiatry, King's College London, De Crespigny Park, Denmark Hill, SE5 8AF, UK, Tel: +44 20 7848 5403, Fax: +44 20 7848 0866, E-mail: yannis.paloyelis@kcl.ac.uk

Received 4 May 2010; revised 5 July 2010; accepted 17 July 2010

Lee *et al*, 2009; Pothuizen *et al*, 2005). Dysfunctions in the neurobiological systems involved, from the cellular to the network level, are also postulated to give rise to changes in reinforcement sensitivity in ADHD (see, eg, Frank *et al*, 2007; Sagvolden *et al*, 2005; Tripp and Wickens, 2008; and see Luman *et al*, 2010 for a review). Pharmacological, genetic, and neuroimaging studies converge on the idea that ADHD is associated with deficiencies in dopamine signaling and abnormalities in dopamine-modulated frontostriatal circuits (Bush, 2010; Gizer *et al*, 2009; Krause *et al*, 2000; Paloyelis *et al*, 2007; Spencer *et al*, 2007; Volkow *et al*, 2009; Volkow *et al*, 2007).

As predicted by most of these neurobiological models (Luman *et al*, 2010), ADHD patients consistently prefer smaller-immediate over larger-delayed rewards using the choice-delay paradigm (Willcutt *et al*, 2008; see Paloyelis *et al*, 2009 for a review). In the choice-delay paradigm, participants make repeated choices between two *standard* amounts available at *fixed* delays. This paradigm is, however, faced with certain methodological issues: reduced sensitivity to detect individual differences in impulsivity, the confounding of time on task with choice preference, and concerns over the validity of using tertiary reinforcers to elicit impulsive behavior (Kuntsi *et al*, 2006b; Paloyelis *et al*, 2009).

Delay discounting paradigms overcome many of these problems, but have been used in few studies with ADHD patients, yielding somewhat inconsistent findings. In a study using a hypothetical discounting task, adolescents with ADHD and comorbid oppositional-defiant disorder made more impulsive choices than controls when the maximum hypothetical gain per trial was \$100, but not when it was \$1000 (Barkley *et al*, 2001). Using tasks presenting choices between real rewards and delays, Scheres *et al* (2006) did not find any differences, but in a later study where they examined ADHD subtypes separately they found that the ADHD-combined subtype (ADHD-CT) group, but not the inattentive subtype group, discounted rewards more steeply than controls (Scheres *et al*, 2010).

Experimentally measured discounting rates using hypothetical tasks show good temporal stability (Audrain-McGovern *et al*, 2009; Beck and Triplett, 2009; Kirby, 2009; Ohmura *et al*, 2006), similar to personality traits (Costa and McCrae, 1992), suggesting that they reflect a relatively stable individual difference. Real-time tasks, where participants experience the rewards and delays associated with each decision they make, are thought to be sensitive to state changes in impulsivity (McDonald *et al*, 2003; Reynolds *et al*, 2006; Reynolds and Schiffbauer, 2004). Evidence from acute pharmacological interventions with psychomotor stimulants and suppressants supports the greater sensitivity of real-time tasks to manipulations likely to induce changes in state impulsivity: discounting rates are affected when real-time task are used, but not in hypothetical tasks (Acheson and de Wit, 2008; Acheson *et al*, 2006; De Wit, 2009; McDonald *et al*, 2003; Richards *et al*, 1999). For example, methylphenidate reduced discounting rates in a real-time task but not in a hypothetical discounting task in ADHD patients (Shiels *et al*, 2009).

To our knowledge, no studies to date have examined the effects of variation in genes involved in dopamine signaling

on impulsive decision making in adolescents, whereas a single study has examined the effects of *COMT*_{Val158Met} in an adult sample of abstinent alcoholics and healthy participants (Boettiger *et al*, 2007). The present study was therefore designed to address these issues, focusing on the following three aims:

- (1) To compare impulsive decision-making paradigms using a hypothetical and a real-time discounting task. We predicted that the ADHD-CT group would show higher discounting rates on both tasks and higher trait impulsivity scores on a self-report measure.
- (2) To test if the impact of reward magnitude on the rate of discounting of hypothetical rewards differs depending on diagnosis. If delay exerts a greater influence in guiding impulsive choice in ADHD-CT compared with controls, as some theories would predict (Sonuga-Barke, 2005), we would expect that the effect of reward magnitude on discounting rates in the hypothetical task would be attenuated in the ADHD-CT group compared with controls (expecting a significant interaction between diagnosis and reward magnitude).
- (3) To investigate whether common variants of the genes for the dopamine transporter (*DAT1*_{10/6} haplotype) (Asherson *et al*, 2007; Brookes *et al*, 2006b) and the catechol-O-methyl transferase (*COMT*_{Val158Met}) enzyme (Lachman *et al*, 1996), which are responsible for the bulk of dopamine degradation in the striatum and the prefrontal cortex, respectively (Akil *et al*, 2003; Chen *et al*, 2004; Lewis *et al*, 2001; Mazei *et al*, 2002; Morón *et al*, 2002), and a common variant of the dopamine D₄-receptor gene (*DRD4*_{48bp-VNTR}) (Asghari *et al*, 1995) that has consistently been associated with ADHD (Brookes *et al*, 2006a; Gizer *et al*, 2009), can predict performance in measures of different aspects of impulsivity.

MATERIALS AND METHODS

Sample

In all, 68 boys of Caucasian origin (ADHD-CT = 36; control = 32) aged 11–20 years ($M = 15.42$, $SD = 2.05$) were recruited from a sample that had taken part in a previous study (Andreou *et al*, 2007; Kuntsi *et al*, in press; Wood *et al*, 2009) (see Table 1 for clinical and demographic data). The clinical group was part of the London branch of the International Multi-Centre ADHD Genetics (IMAGE) project (Chen *et al*, 2008; Kuntsi *et al*, 2006a; see Supplementary Material for details). Participants were selected based on *DAT1*_{10/6} haplotype dosage (2 copies and <2 copies) (Asherson *et al*, 2007; Brookes *et al*, 2006b), forming four roughly equally sized subgroups. Participants receiving stimulant treatment for ADHD (72%) discontinued their medication for at least 48 h before testing (see Supplementary Material, Table S1 for details on medication history). Parents completed the long form of the revised Conners' Rating Scale (Conners *et al*, 1998) at the time of testing. In all, 30 participants (83%) had current total DSM-IV ADHD T-scores in the diagnostic range (with 67% scoring in the diagnostic range on both the inattention and

Table 1 Clinical, Experimental, and Demographic Data for the ADHD-CT and Control Participants, Presented by *DAT1*_{10/6} Haplotype

	Mean (SD)				Significance test	p
	ADHD-CT <i>DAT1</i> _{10/6} (2 copies) (n = 20)	ADHD-CT <i>DAT1</i> _{10/6} (<2 copies) (n = 16)	Control <i>DAT1</i> _{10/6} (2 copies) (n = 16)	Control <i>DAT1</i> _{10/6} (<2 copies) (n = 16)		
Age (years)	15.05 (2.35)	15.65 (2.33)	15.35 (1.4)	15.76 (2.02)	F _{group} (1, 64) = 0.16 F _{haplotype} (1, 64) = 0.97 F _{interaction} (1, 64) = 0.03	0.69 0.32 0.85
IQ	107.8 (13.68)	102.38 (12.97)	111.06 (9.9)	114.12 (12.19)	F _{group} (1, 64) = 6.21 F _{haplotype} (1, 64) = 0.15 F _{interaction} (1, 64) = 1.98	0.015 0.70 0.16
Parents' SES	41.42 (10.74)	38.07 (15.53)	47.16 (11.62)	48.06 (12.68)	F _{group} (1, 60) = 6.18 F _{haplotype} (1, 60) = 0.15 F _{interaction} (1, 60) = 0.45	0.016 0.70 0.50
Conners' DSM-IV ADHD inattention ratings (parent)	67.05 (9.89)	72.25 (7.72)	46.53 (4.61)	47.75 (6.43)	F _{group} (1, 63) = 144.00 F _{haplotype} (1, 63) = 2.93 F _{interaction} (1, 63) = 1.13	0.000 0.092 0.29
Conners' DSM-IV ADHD hyperactivity/impulsivity ratings (parent)	78.1 (13.6)	81.94 (12.17)	47.4 (7.19)	47.81 (5.84)	F _{group} (1, 63) = 157.20 F _{haplotype} (1, 63) = 0.68 F _{interaction} (1, 63) = 0.44	0.000 0.41 0.51
Conners' DSM-IV ADHD total ratings (parent)	73.95 (12.07)	79.44 (9.35)	46.53 (5.42)	47.56 (5.93)	F _{group} (1, 63) = 182.61 F _{haplotype} (1, 63) = 2.21 F _{interaction} (1, 63) = 1.03	0.000 0.14 0.31
Barratt's Impulsiveness Scale, version 11, adolescents	72.7 (9.56)	84.19 (10.66)	66.12 (10.16)	68.94 (8.02)	(see Tables 3 and 4)	
Hypothetical delay discounting task (AUC)	0.5 (0.27)	0.36 (0.21)	0.48 (0.23)	0.72 (0.28)	(see Tables 3 and 4)	
Real-time delay discounting task (AUC)	0.73 (0.25)	0.74 (0.23)	0.81 (0.19)	0.79 (0.13)	(see Tables 3 and 4)	

Abbreviations: ADHD-CT, attention deficit/hyperactivity disorder-combined subtype; SES, socioeconomic status; AUC, area under the curve.

hyperactivity/impulsivity subscales). Of the six participants with current total parent T-scores of <63 (the clinical cutoff in IMAGE; Marco *et al*, 2009), three were still receiving stimulant treatment. To include as many participants with ADHD as possible, we did not exclude participants with current parent ratings below the diagnostic range, but we repeated analyses excluding them. No comorbid disorder was associated with any of the three dopamine gene variants (see Supplementary Material, Tables S2–S4). Current levels of substance use were assessed using a self-report questionnaire (see Supplementary Material, Table S5). Analyses were repeated excluding four ADHD-CT participants with frequent or occasional use of drugs and/or heavy cigarette smoking (>5 cigarettes per day).

Control participants were aged matched to the clinical group. To exclude potential undiagnosed ADHD cases, participants with initial parent or teacher Conners' T-scores of >63 on any DSM-IV subscale were excluded. The *DRD4*_{48bp-VNTR} genotype could not be determined for one control participant, who was not included in some of the analyses.

Genotyping

Standard genotyping procedures were used (described in the Supplementary Material). *DAT1*_{10/6} status (2 copies and

<2 copies) was determined on the basis of zygosity for the constituent *DAT1* polymorphisms (see Supplementary Material, Table S6). Frequencies for the *DRD4*_{48bp-VNTR} at exon 3 and the *COMT*_{Val158Met} polymorphisms are presented in Supplementary Material (Table S7); no significant associations were observed with disease status. For statistical analyses we created binary groups for each genotype (*COMT*_{Val158Met}: methionine homozygotes vs valine carriers; *DRD4*_{48bp-VNTR}: 7R-carriers vs rest).

Behavioral Measures of Impulsivity

Hypothetical delay discounting task. A computerized, adjusting-amount algorithm asked participants to make successive choices between a standard monetary amount available after a delay and a smaller adjusting amount available immediately (see Richards *et al*, 1999 for details on the adjusting-amount procedure). The subjective value at which a participant was indifferent between the larger-delayed and the smaller-immediate reward (*indifference point*) was estimated for eight delay intervals (1, 2, 7, 14, 30, 60, 120, and 180 days) using three standard amounts (£5, £15, and £30). Discounting curves were plotted for each participant and for each monetary magnitude using normalized indifference points (proportions of the standard amount) and delays

(proportions of the maximum delay). Questions were presented in a random order. To increase motivation and accuracy, participants were told at the end of the task that the choice from one question, selected at random by themselves, would be honored. To keep the general level of compensation at acceptable levels for adolescents, we followed a credible procedure ensuring that the 'randomly' selected question always came from the £5 category.

Real-time delay discounting task. We used the UK version of the experiential discounting task (for a detailed description, see Supplementary Material) (Reynolds *et al*, 2006; Reynolds and Schiffbauer, 2004), in which participants experience the delays and receive the rewards associated with each decision they make. An indifference point was calculated for each of four choice blocks where participants made successive choices between a certain and immediate adjusting amount (initially £0.10) and a probabilistic (35%) £0.20 available at different delays (1, 7, 14, or 28 s depending on choice block). Because of the probabilistic nature of the larger amount (consistent across choice blocks), indifference points for each choice block were divided by the indifference point for the no-delay (1 s) choice block to control for individual differences in responses to probabilistic outcomes (Reynolds and Schiffbauer, 2004). The indifference points for the remaining blocks now reflected the relative change in discounting for each choice block compared with the no-delay block. This data adjustment and the systematic manipulation of delays render this task a measure of delay discounting, rather than risk discounting (Reynolds *et al*, 2006). Individual discounting curves were plotted after normalizing indifference points and delays (see description above). The four choice blocks were administered in random order after the participant had completed a practice session with the 7-s choice block.

Estimating discounting rates. In both tasks, discounting rates were assessed using the widely used (Scheres *et al*, 2010; Shiels *et al*, 2009) area-under-the-curve (AUC) method, which presents theoretical and practical advantages over alternatives (Beck and Triplett, 2009; Myerson *et al*, 2001). AUC values range from 0 to 1, with lower values indicating a higher rate of discounting and higher levels of impulsivity.

Self-Report Measures of Trait Impulsivity

Barratt's Impulsiveness Scale for Adolescents, version 11 (BIS-11A). Adapted from the adult version (Patton *et al*, 1995; Stanford *et al*, 2009), it has been increasingly used with younger samples in many languages (Cosi *et al*, 2008; Fossati *et al*, 2002; Leshem and Glicksohn, 2007; von Diemen *et al*, 2007). With adult samples BIS-11 can differentiate between non-planning, attentional, and motor aspects of impulsivity (Patton *et al*, 1995; Stanford *et al*, 2009), yet a consistent factor structure is not reproduced with younger groups (Cosi *et al*, 2008; Fossati *et al*, 2002; Leshem and Glicksohn, 2007; von Diemen *et al*, 2007). Thus, the total scale score (ranging from 30 to 120) has been recommended (Fossati *et al*, 2002) and used (Melanko *et al*, 2009) as the most appropriate measure of trait impulsivity with adolescents. Higher scores reflect higher trait impulsivity levels.

Other Measures

ADHD rating scales. ADHD symptoms were assessed using the 18 DSM-IV items from the long form of the revised Conners' Parent Rating Scales (Conners *et al*, 1998).

General intelligence. The vocabulary, similarities, picture completion, and block design subtests of the Wechsler Intelligence scale for children III (Wechsler, 1991) or the Wechsler Intelligence Scale for Adults (Wechsler, 1997) were used to obtain an estimate of the child's IQ at the time of the initial assessment.

Socioeconomic status (SES). The SES status of participants' families was estimated based on information about both parents' educational and occupational background using the Barratt Simplified Measure of Social Status (Barratt, 2006), which updated Hollingshead's Four-Factor Index of Social Status (Hollingshead, 1975) with a current occupation list (Baldwin and Dadds, 2007). Higher scores indicate higher socioeconomic levels.

Procedure

The discounting tasks (in counterbalanced order) and the BIS-11A questionnaire were administered under laboratory conditions in the afternoon session during a day-long visit to our research center as part of a larger study. Parent ratings were obtained during the same visit. Any cash earned during the tasks was exchanged for vouchers of equivalent value for the participant's preferred store. Ethics approval was obtained from the South London and Maudsley NHS Foundation Trust and informed consent was obtained from all participants.

Statistical Analyses

Age was covaried in all analyses; *p*-values after covarying IQ and SES are reported separately. A three-step approach was utilized to address the goals of this study.

First, separate MANCOVA/ANCOVA tests were conducted for each measure of impulsivity to test for an ADHD-CT deficit independently from genetic effects (obtaining results directly comparable to previous studies). The MANCOVA was used for the hypothetical discounting task, with the AUC measures for each reward magnitude used as the dependent variables and diagnosis (ADHD-CT vs control) as the between-subjects factor. Given the high correlations (Pearson's *r* from 0.80 to 0.94) in the discounting rates across magnitudes in the hypothetical task, we estimated a single overall AUC measure by averaging the three indifference points at each delay interval for each participant. This measure was used in an ANCOVA to estimate the effect size for the diagnosis effect and in subsequent genetic analyses.

Second, to test whether reward magnitude differentially affected the rate of discounting (AUC; reflecting the decrease in the subjective value of a reward with delay) of hypothetical rewards for the ADHD-CT and control groups, we used a mixed ANCOVA, with reward magnitude as the within-subjects factor (£5, £15, and £30) and diagnosis as the between-subjects factor.

Table 2 Pearson's Correlation Coefficients Between Impulsivity Measures and ADHD Symptom Ratings

	Control group			ADHD-CT group			Pooled within-group correlations		
	BIS-11A	HDT	EDT	BIS-11A	HDT	EDT	BIS-11A	HDT	EDT
Hypothetical discounting task (HDT)	−0.24			−0.29			−0.26*		
Real-time discounting task (EDT)	−0.20	−0.12		0.27	−0.13		0.11	−0.12	
DSM-IV ADHD inattention ratings ^a	0.42*	−0.23	−0.34	0.32	−0.19	−0.22	0.35**	−0.20	−0.25*
DSM-IV ADHD hyperactivity impulsivity ratings ^a	0.44*	−0.08	−0.26	0.27	−0.21	−0.24	0.31*	−0.16	−0.24
DSM-IV ADHD total ratings ^a	0.51**	−0.20	−0.39*	0.36*	−0.20	−0.24	0.39***	−0.19	−0.27*

Abbreviations: ADHD, attention deficit/hyperactivity disorder; BIS-11A, Barratt's Impulsiveness Scale (version 11, adolescents).

^aParent ratings on the revised version of the Conners' Rating Scales.

* $p < 0.05$; ** $p < 0.01$; *** $p < 0.001$.

See Supplementary Material for details on the calculation of pooled within-group correlations.

Third, we tested for the main effects of *DAT1*_{10/6}, *COMT*_{val158met} and the *DRD4*_{48bp-VNTR}, and their interactions with diagnosis, using separate ANCOVAs for each measure of impulsivity. Family-wise error rate was contained at the $\alpha = 0.05$ level using a Bonferroni adjustment of the critical p -value for each ANCOVA to $\alpha = 0.05/3 = 0.017$. Similarly, a critical p -value of $\alpha = 0.017/4 = 0.004$ was used when simple effects analyses for significant or trend-level (< 0.10) interactions were conducted. Owing to the relatively small sample and cell sizes, p -values obtained from parametric tests were confirmed using nonparametric procedures. We performed 10 000 random permutations on the age-regressed residuals for each measure and estimated p -values and 95% confidence intervals on the basis of Monte Carlo simulations.

All analyses were repeated without including the six ADHD participants with current T-scores of < 63 on the parent Conners' total ADHD scale. In all cases, p -values and effect sizes either improved or remained approximately the same (see Supplementary Material). Similarly, excluding the four ADHD participants with frequent or occasional current use of cannabis (including two with frequent/occasional use of cocaine and all heavy smokers (> 5 cigarettes per day)) did not change the observed pattern of results (see Supplementary Material). Pearson's correlation coefficients among ADHD ratings and the impulsivity measures are provided in Table 2.

RESULTS

Adolescents with ADHD-CT Displayed Higher Levels of Impulsivity

The ADHD-CT group displayed significantly steeper discounting rates (AUC) compared with controls in the hypothetical but not the real-time discounting task (see Table 3 and Figure 1). They also reported significantly higher levels of trait impulsivity. These effects remained significant after covarying IQ and SES.

Delay but not Reward Magnitude Affected Discounting Rates (AUC) in the Hypothetical Discounting Task in the ADHD-CT Group

Discounting rates in the hypothetical task were submitted to a mixed ANCOVA with monetary magnitude (£5, £15, and

£30) as the repeated factor and diagnosis as the between-subjects factor. The rate of discounting decreased as the magnitude of the monetary amount increased (magnitude effect: $F_{(2,132)} = 7.62$, $p = 0.001$; covarying IQ/SES: $p < 0.001$), but only in the control group (magnitude \times diagnosis: $F_{(2,132)} = 5.39$, $p = 0.009$; covarying IQ/SES: $p = 0.022$; Figure 2; p -values were adjusted using the Greenhouse-Geisser epsilon).

*DAT1*_{10/6} and *COMT*_{val158Met} Polymorphisms Predicted Discounting Rates (AUC) in the Hypothetical Task and Trait Impulsivity Ratings

Hypothetical discounting task. *DAT1*_{10/6} status interacted with diagnosis (Table 4 and Figure 3a); this effect remained significant after covarying trait impulsivity scores ($p_{\text{corrected}} < 0.05$; $p_{\text{uncorrected}} = 0.005$; $R^2 = 0.086$). Adolescents with ADHD-CT showed higher discounting rates than controls only among carriers of < 2 copies of *DAT1*_{10/6} ($p_{\text{corrected}} < 0.05$; $p_{\text{uncorrected}} < 0.001$); there was no group difference for *DAT1*_{10/6} homozygotes ($p_{\text{uncorrected}} = 0.61$). Furthermore, *DAT1*_{10/6} homozygotes showed significantly higher discounting rates compared with those with < 2 copies ($p_{\text{corrected}} < 0.05$; $p_{\text{uncorrected}} = 0.002$) among control participants. Numerically, the effect was in the opposite direction for the ADHD-CT group, but was not significant ($p_{\text{uncorrected}} = 0.12$).

The *COMT*_{val158met} polymorphism predicted discounting rates independent of diagnosis; covarying trait impulsivity ratings had no impact ($p_{\text{corrected}} < 0.05$; $p_{\text{uncorrected}} = 0.013$; $R^2 = 0.068$). Met-allele homozygotes showed higher discounting rates than Val-allele carriers ($p_{\text{corrected}} < 0.05$; $p_{\text{uncorrected}} = 0.014$).

Real-time discounting task. No significant genetic effects were observed (Table 4).

Self-reported trait impulsivity ratings. *DAT1*_{10/6} status predicted trait impulsivity ratings independent of diagnosis (Table 4 and Figure 3b). *DAT1*_{10/6} homozygotes reported lower impulsivity levels than carriers of < 2 copies, an effect that remained highly significant after covarying discounting rates in the hypothetical task ($p_{\text{corrected}} < 0.05$; $p_{\text{uncorrected}} = 0.004$; $R^2 = 0.095$). A trend for a diagnosis \times *DAT1*_{10/6} ($p_{\text{uncorrected}} < 0.10$) suggested that this effect might be significant only in the ADHD-CT group ($p_{\text{corrected}}$

Table 3 (M)ANCOVAs Examining the Effect of Diagnosis on Measures of Impulsivity (Age Is Covaried)

Measure	Statistical test	Pillai's trace	F	df	p	R ²	η_p^2	Covarying IQ	Covarying IQ/SES
Hypothetical delay discounting task	MANCOVA	0.20	5.21	3, 63	0.003	—	—	0.011	0.024
	ANCOVA	—	6.24	1, 65	0.015	0.081	0.088	—	—
Barratt's Impulsiveness Scale (version 11)	ANCOVA	—	16.05	1, 65	<0.001	0.198	0.198	0.001	0.003
Real-time delay discounting task	ANCOVA	—	1.69	1, 65	>0.10	0.025	0.025	>0.10	>0.10

Diagnosis: attention deficit/hyperactivity disorder-combined subtype and controls.




Figure 1 Discounting curves for the hypothetical (overall discounting rate, group median) and real-time (group mean) delay discounting tasks for the attention deficit/hyperactivity disorder-combined subtype (ADHD-CT) and control groups. The graphs plot group indifference points—the subjective value (percentage of larger-delayed amount) at which participants were indifferent between the larger-delayed and the smaller-immediate reward—against delay intervals. In the real-time task, indifference points are adjusted for individual differences in response to probabilistic outcomes.

<0.05; $p_{\text{uncorrected}} = 0.002$; control group: $p_{\text{uncorrected}} > 0.10$). It further indicated that adolescents with ADHD-CT reported higher levels of impulsivity compared with controls if they carried <2 *DAT1*_{10/6} copies ($p_{\text{corrected}} < 0.05$; $p_{\text{uncorrected}} < 0.001$) but not among *DAT1*_{10/6} homozygotes ($p_{\text{uncorrected}} = 0.13$).

There were no effects for the *DRD4*_{48bp-VNTR} polymorphism on any on the impulsivity measures (see Table 4).

DISCUSSION

This study used a relatively homogeneous sample of male adolescents with ADHD-CT and typically developing controls, and reports three novel findings. First, when comparing performance on a hypothetical and a real-time discounting task, we found that the ADHD-CT group showed steeper discounting rates (smaller AUC, denoting higher levels of impulsivity) in the hypothetical task, but not when rewards/delays were real. The ADHD-CT group also reported higher trait impulsivity self-ratings than controls.

Second, discounting rates in the hypothetical task were influenced by delay but not reward magnitude for the ADHD-CT group. Third, *DAT1*_{10/6} haplotype dosage and the *COMT*_{Val158Met} genotype predicted discounting rates in the hypothetical discounting task (but not the real-time task) and trait impulsivity ratings.

The contrast between discounting rates in the hypothetical and real-time tasks in terms of diagnostic group and gene effects is novel. It cannot be attributed to validity issues with the real-time task, as both groups showed discounting in this study, supporting evidence from earlier studies (Fields *et al*, 2009; Krishnan-Sarin *et al*, 2007; Melanko *et al*, 2009). These data therefore suggest that discounting rates in hypothetical and real-time tasks may reflect different aspects of impulsivity. As discussed in the Introduction, one possible distinction is that performance in the hypothetical task putatively reflects a relatively stable individual difference (an aspect of trait impulsivity), whereas performance in the real-time task may be more sensitive to the transient effects of factors affecting state impulsivity, such as acute pharmacological interventions




Figure 2 Effect of monetary magnitude of the delayed amount on discounting rates (measured using the area under the curve) for adolescents with attention deficit/hyperactivity disorder-combined subtype (ADHD-CT) and control participants in the hypothetical delay discounting task.

(Shiels *et al*, 2009) or task characteristics. Further support for this distinction is provided by the lack of significant correlations between discounting rates in the real-time task and trait impulsivity scores, or discounting rates in the hypothetical task, which is consistent with the literature (Fields *et al*, 2009; Krishnan-Sarin *et al*, 2007; Melanko *et al*, 2009; Reynolds *et al*, 2008; Reynolds *et al*, 2006) with few exceptions (Meda *et al*, 2009; Reynolds, 2006a).

Although impulsive decision making in discounting paradigms is consistent with the pattern of neurobiological deficits observed in ADHD and predicted by most neurobiological models of the disorder (see Luman *et al*, 2010 for a review), such models do not distinguish between performance in real-time and hypothetical paradigms, as most of them imply context-invariant deficits. One possible explanation for the lack of case-control differences in the real-time discounting task could be that decision making in this task (where participants experience the delays and rewards associated with each choice they make) impacts on an individual's current state of arousal considerably more than in the hypothetical task, which consequently might induce changes in state impulsivity (to which real-time tasks are assumed to be more sensitive). As shown using reaction time performance paradigms, the performance of ADHD patients tends to approach or equal control levels under task conditions that increase stimulation/arousal levels (Andreou *et al*, 2007; Johnson *et al*, 2007a; Johnson *et al*, 2007b; Konrad *et al*, 2000; Kuntsi *et al*, 2009; McInerney and Kerns, 2003; O'Connell *et al*, 2008; Slusarek *et al*, 2001; Uebel *et al*, 2010). Therefore, it is possible that discrepancies among studies using real-time tasks (eg, this study and Scheres *et al*, 2010) might be explained by altered arousal-regulation processes in ADHD. This could occur if case-control differences in measures of state impulsivity

depend on the extent to which specific task characteristics alter arousal levels, because of differences in factors such as the monotony of the task, salience of the reward, and length and expectancy of the delay. This explanatory framework leads to specific predictions that can be investigated by examining the relationship between indices of arousal-regulation deficits and performance in a range of real-time discounting paradigms.

Another novel finding in this study was the lack of reward magnitude effects in the hypothetical task for the ADHD-CT group, suggesting that their choices were influenced only by delay. Discounting rates in the control group were inversely related to reward magnitude, consistent with the literature (Chapman and Winquist, 1998; Kirby and Marakovi, 1996; Myerson and Green, 1995; Smith and Hantula, 2008). This discrepancy indicates that different processes might drive discounting in ADHD-CT and controls. One possible explanation could be that ADHD-CT participants were particularly sensitive to delays, as would be predicted by delay aversion theory (Sonuga-Barke, 2005), thus overriding the effect of reward magnitude, at least when choices do not affect current state and with the range of values used in this study.

The predictive value of variants for the *DAT1* and *COMT* genes was restricted to measures of relatively stable aspects of impulsivity. Previous studies have linked the *DAT1* 9-repeat allele (one of the two polymorphisms constituting the haplotype studied here) with increased ventral striatal reactivity during reward anticipation (Forbes *et al*, 2009), which in turn had been linked with steeper hypothetical discounting rates (Hariri *et al*, 2006). In this study we extend this work, showing that *DAT1*_{10/6} dosage directly predicted hypothetical discounting rates and trait impulsivity in ADHD-CT and controls. Previous studies had failed to show a reliable effect of *DAT1* genotype on executive functions/response inhibition (Rommelse *et al*, 2008), although a study focusing on the same *DAT1*_{10/6} haplotype (Bellgrove *et al*, 2009) reported a significant interaction of diagnosis with spatial attention measures explaining 4.9–5.8% of the variance. In our study, *DAT1*_{10/6} interacted with diagnosis in the hypothetical task, explaining 11.7% of the variance. Although the *DAT1*_{10/6} interaction with diagnosis was only a trend for trait impulsivity ratings, it accounted for 4.7% of the variance, which is comparable to the effect sizes observed with the same haplotype in the study by Bellgrove *et al* (2009).

Post hoc analyses of the interaction effect (controlling for multiple testing) revealed remarkably similar effects of *DAT1*_{10/6} dosage across measures, providing a within-sample replication of the effect. The differential effects of *DAT1*_{10/6} dosage depending on diagnostic status (and the reverse) are not entirely surprising. Dopamine effects on neurocognitive functions have been shown to reflect an inverted-U shape (Goldman-Rakic *et al*, 2000). Therefore, assuming that there is some optimal level of dopamine modulating the function of frontal-striatal circuits underlying impulsive behavior, any factor disturbing this balance in either direction would be likely to impair behavior (Egan *et al*, 2001; Goldberg *et al*, 2003; Williams-Gray *et al*, 2007).

Steeper discounting rates in the hypothetical task were also associated with homozygosity for the *COMT*_{Val158Met} Met-allele (which substantially reduces dopamine catabo-

Table 4 ANCOVAs Examining the Role of *DAT1*_{10/6} haplotype, *COMT*_{Val158Met} and *DRD4*_{48bp-VNTR} Status on Measures of Impulsivity

Measure	Effect	Parametric tests						Permutation tests ^a		Covarying IQ	Covarying IQ/SES
		F	df	P _{corr.}	P _{uncorr.}	R ²	η ² _p	p	CI 95%	P	p
Hypothetical delay discounting task	<i>DAT1</i> _{10/6}	1.34	1,58	NS	>0.10	0.014	0.023	>0.10	—	>0.10	>0.10
	Diagnosis × <i>DAT1</i> _{10/6}	10.98	1,58	<0.05	0.002	0.117	0.159	0.002	0.001, 0.003	0.002	0.004
	<i>COMT</i> _{Val158Met}	6.45	1,58	<0.05	0.014	0.069	0.100	0.014	0.011, 0.016	0.043	0.031
	Diagnosis × <i>COMT</i>	0.17	1,58	NS	>0.10	0.002	0.003	>0.10	—	>0.10	>0.10
	<i>DRD4</i> _{48bp-VNTR}	0.15	1,58	NS	>0.10	0.002	0.005	>0.10	—	>0.10	>0.10
	Diagnosis × <i>DRD4</i> _{48bp-VNTR}	0.51	1,58	NS	>0.10	0.005	0.009	>0.10	—	>0.10	>0.10
Barratt's Impulsiveness Scale (version 11)	<i>DAT1</i> _{10/6}	7.16	1,58	<0.05	0.010	0.108	0.110	0.008	0.007, 0.010	0.010	0.052
	Diagnosis × <i>DAT1</i> _{10/6}	2.83	1,58	NS	0.098	0.032	0.047	0.093	0.089, 0.099	>0.10	>0.10
	<i>COMT</i> _{Val158Met}	0.00	1,58	NS	>0.10	0.000	0.000	>0.10	—	>0.10	>0.10
	Diagnosis × <i>COMT</i>	0.05	1,58	NS	>0.10	0.000	0.000	>0.10	—	>0.10	>0.10
	<i>DRD4</i> _{48bp-VNTR}	0.08	1,58	NS	>0.10	0.000	0.000	>0.10	—	>0.10	>0.10
	Diagnosis × <i>DRD4</i> _{48bp-VNTR}	0.73	1,58	NS	>0.10	0.000	0.000	>0.10	—	>0.10	>0.10
Real-time delay discounting task	<i>DAT1</i> _{10/6}	0.00	1,58	NS	>0.10	0.001	0.001	>0.10	—	>0.10	>0.10
	Diagnosis × <i>DAT1</i> _{10/6}	0.01	1,58	NS	>0.10	0.003	0.004	>0.10	—	>0.10	>0.10
	<i>COMT</i> _{Val158Met}	0.02	1,58	NS	>0.10	0.006	0.006	>0.10	—	>0.10	>0.10
	Diagnosis × <i>COMT</i>	0.00	1,58	NS	>0.10	0.000	0.000	>0.10	—	>0.10	>0.10
	<i>DRD4</i> _{48bp-VNTR}	0.03	1,58	NS	>0.10	0.011	0.011	>0.10	—	>0.10	>0.10
	Diagnosis × <i>DRD4</i> _{48bp-VNTR}	0.01	1,58	NS	>0.10	0.002	0.002	>0.10	—	>0.10	>0.10

^aIn all, 10 000 random permutations on the age-regressed residuals for each measure were performed and *p*-values and 95% confidence intervals (95% CIs) were estimated on the basis of Monte Carlo simulations.

Diagnosis: attention deficit/hyperactivity disorder-combined subtype and controls.




Figure 3 (a) Mean overall discounting rates (measured using the area under the curve (AUC)) in the hypothetical discounting task as a function of diagnosis (attention deficit/hyperactivity disorder-combined subtype (ADHD-CT) and controls) and *DAT1*_{10/6} haplotype status (2 copies and <2 copies). Significance values from simple effects analyses are shown. (b) Self-reported scores on Barratt's Impulsiveness Scale (BIS-11A) as a function of diagnosis (ADHD-CT and controls) and *DAT1*_{10/6} status (2 copies and <2 copies). Significance values from simple effects analyses are shown.

lism in the prefrontal cortex) (Chen *et al*, 2004; Lachman *et al*, 1996; Lotta *et al*, 1995), independent of diagnosis ($R^2 = 6.9\%$). This finding is consistent with evidence showing that the dopamine-modulated prefrontal cortex

has a key role in choice impulsivity (Kable and Glimcher, 2007; Kable and Glimcher, 2009; Kheramin *et al*, 2004; McClure *et al*, 2007; McClure *et al*, 2004; Winstanley *et al*, 2006). The role of *COMT*_{Val158Met} in impulsivity has received

scant attention to date, with the Met-allele being associated with reduced immediate-reward bias (Boettiger *et al*, 2007) and increased novelty seeking (Golimbet *et al*, 2007). Variants of *COMT*_{Val158Met} and *MAOA* (which perform most of the catecholamine catabolism in the prefrontal cortex) (Chen *et al*, 2004; Gogos *et al*, 1998; Huotari *et al*, 2002; Matsumoto *et al*, 2003; Tunbridge *et al*, 2004; Volavka *et al*, 2004) with reduced enzymatic activity have been associated with aggressive impulsivity and suicide (Contini *et al*, 2006; Huang *et al*, 2004; Volavka *et al*, 2004). Meta-analytic evidence has not associated ADHD with *COMT*_{Val158Met} (Gizer *et al*, 2009), although some evidence suggests that gender could be a moderating factor with Met being the risk allele in boys (Biederman *et al*, 2008; Qian *et al*, 2003). This is consistent with evidence that *COMT*_{Val158Met} is differentially expressed across genders (Dempster *et al*, 2006; Gogos *et al*, 1998). A recent study confirmed the association of the Met-allele with increased risk for ADHD in children in a predominantly male (84%) sample, and also reported that Met was associated with increased symptom severity (Pálmason *et al*, 2010). Moreover, Met has been associated with ADHD inattention and hyperactivity impulsivity symptoms in adult community samples (Ettinger *et al*, 2006; Gothelf *et al*, 2007; Michaelovsky *et al*, 2008; Reuter *et al*, 2006). The *DRD4*_{48bp-VNTR} was not associated with any impulsivity measure, in line with a recent meta-analytic review (which suggested that another marker on the gene might be related to impulsivity; Munafó *et al*, 2008).

Although our data confirmed our hypotheses that variation in *DAT1*_{10/6} and *COMT*_{Val158Met} predicts behavioral impulsivity, the precise mechanisms mediating these effects, in terms of dopamine signaling or the neural processes involved, remain unknown. One reason is the lack of sufficient evidence regarding the precise functional effects of these genetic variants. For example, although the *DAT1*_{3'UTR} 48 base-pair VNTR polymorphism (part of the *DAT1*_{10/6} haplotype) has been associated with alterations in gene expression, evidence regarding the impact of specific variants is inconsistent (for a review, see van de Giessen *et al*, 2009). Second, even when the effects of genetic variation in terms of gene expression and dopamine catabolism are better understood (eg, *COMT*_{Val158Met}), predicting the precise role of a variant on behavioral impulsivity may still be elusive, given that it is likely to depend on the dynamic interaction among many factors and reflect the outcome of long-term adaptation processes. For example, the *DAT1* 10-repeat allele is considered to increase risk for ADHD in children and adolescents, yet in adults (and thus, by definition, in persisting forms of the disorder) ADHD has been associated with the 9-repeat allele (Franke *et al*, 2008; Franke *et al*, 2010). Our study, alongside recent work (Bellgrove *et al*, 2009), has documented that variation in *DAT1*_{10/6} modulates behavioral outcomes, highlighting the need for a better understanding of the impact of this haplotype on gene expression, dopamine signaling, and neural mechanisms involved. Some promising initial evidence has indicated that the *DAT1*_{3'UTR} polymorphism may be modulating the responsiveness of the striatum during a response inhibition and a reward-processing paradigm (Durstun *et al*, 2008; Forbes *et al*, 2009).

Our genetic findings finally suggest that the rate of experimental discounting and trait impulsivity ratings reflect distinct aspects of impulsivity. Covarying trait impulsivity ratings or discounting rates when examining the genetic effects on either measure left effect sizes virtually unchanged and results remained significant. This finding provides no support for the mediation of gene effects on discounting by impulsive behavior (or the reverse), but rather that the two measures of relatively stable impulsivity predispositions (hypothetical discounting and self-report rating scale) represent pleiotropic effects (multiple outcomes) of the genetic influences.

In summary, the pattern of findings in this study distinguishes between aspects of impulsivity by demonstrating specific effects of diagnosis and variation in dopamine genes (*DAT1*_{10/6} and *COMT*_{Val158Met}) on behavioral and laboratory measures assumed to reflect relatively stable aspects of impulsivity (discounting in the hypothetical task and trait impulsivity self-ratings) but not on discounting rates in a real-time task, which may be more sensitive to factors impacting on state impulsivity levels. However, our study was not designed to distinguish between trait and state aspects of impulsivity, or test the various models predicting impulsive behavior in ADHD; these goals should be addressed in future research. Our data suggest that existing ADHD models need to distinguish between trait and state aspects of impulsivity, and that arousal regulation should be further investigated as one potential mechanism contributing to real-time impulsive decision making in ADHD. Future studies should also employ a wider range of real-time paradigms. The sample size is a limitation of this study; the genetic findings should be considered as preliminary until confirmed within a larger sample, which should also examine gene-by-gene interactions and assess the impact of potential confounds pertaining to the ADHD-CT group, such as possible gene interactions with the effects of long-term stimulant treatment.

ACKNOWLEDGEMENTS

This work was supported in part by UK Medical Research Council Grant G03001896 to J Kuntsi and NIH grants R01MH62873 and R01MH081803 to SV Faraone. We thank all who make this research possible: our participants and their families, who give their time and support so unstintingly, and our research team, Kelly Harris and Chloe Booth. We also thank Dr Brady Reynolds for his permission to use the Experiential Discounting Task, and his advice on analyzing the data obtained from this task.

DISCLOSURE

PA has given sponsored talks and/or been an advisor for Shire, Janssen-Cilag, Eli-Lilly, Flynn Pharma, and Pfizer, regarding the diagnosis and treatment of ADHD; all funds have been donated to the University Research Fund for studies of ADHD. MAM receives research funding from Eli Lilly and is a scientific advisor to Cambridge Cognition. SVF has, in the past year, received consulting fees and has been on Advisory Boards for Eli Lilly, Ortho-McNeil, and Shire

Development and has received research support from Eli Lilly, Pfizer, Shire, and the National Institutes of Health. In previous years, SVF has received consulting fees or has been on Advisory Boards or has been a speaker for the following sources: Shire, McNeil, Janssen, Novartis, Pfizer, and Eli Lilly. In previous years, he has received research support from Eli Lilly, Shire, Pfizer, and the National Institutes of Health. JK has received a speaker's fee from Eli Lilly that has been used for educational and research activities.

REFERENCES

- Acheson A, de Wit H (2008). Bupropion improves attention but does not affect impulsive behavior in healthy young adults. *Exp Clin Psychopharmacol* **16**: 113–123.
- Acheson A, Reynolds B, Richards JB, De Wit H (2006). Diazepam impairs behavioral inhibition but not delay discounting or risk taking in healthy adults. *Exp Clin Psychopharmacol* **14**: 190–193.
- Adriani W, Boyer F, Gioiosa L, Macri S, Dreyer JL, Laviola G (2009). Increased impulsive behavior and risk proneness following lentivirus-mediated dopamine transporter over-expression in rats' nucleus accumbens. *Neuroscience* **159**: 47–58.
- Ainslie G (1975). Specious reward: a behavioral theory of impulsiveness and impulse control. *Psychol Bull* **82**: 463–496.
- Akil M, Kolachana BS, Rothmond DA, Hyde TM, Weinberger DR, Kleinman JE (2003). Catechol-O-methyltransferase genotype and dopamine regulation in the human brain. *J Neurosci* **23**: 2008–2013.
- Andreou P, Neale BM, Chen W, Christiansen H, Gabriels I, Heise A et al (2007). Reaction time performance in ADHD: improvement under fast-incentive condition and familial effects. *Psychol Med* **37**: 1703–1715.
- Anker JJ, Perry JL, Gliddon LA, Carroll ME (2009). Impulsivity predicts the escalation of cocaine self-administration in rats. *Pharmacol Biochem Behav* **93**: 343–348.
- Asghari V, Sanyal S, Buchwaldt S, Paterson A, Jovanovic V, Van Tol HH (1995). Modulation of intracellular cyclic AMP levels by different human dopamine D4 receptor variants. *J Neurochem* **65**: 1157–1165.
- Asherson P, Brookes K, Franke B, Chen W, Gill M, Ebbstein RP et al (2007). Confirmation that a specific haplotype of the dopamine transporter gene is associated with combined-type ADHD. *Am J Psychiatry* **164**: 674–677.
- Audrain-McGovern J, Rodriguez D, Epstein LH, Cuevas J, Rodgers K, Wileyto EP (2009). Does delay discounting play an etiological role in smoking or is it a consequence of smoking? *Drug Alcohol Depend* **103**: 99–106.
- Baldwin JS, Dadds MR (2007). Reliability and validity of parent and child versions of the multidimensional anxiety scale for children in community samples. *J Am Acad Child Adolesc Psychiatry* **46**: 252–260.
- Barkley RA, Edwards G, Laneri M, Fletcher K, Metevia L (2001). Executive functioning, temporal discounting, and sense of time in adolescents with attention deficit hyperactivity disorder (ADHD) and oppositional defiant disorder (ODD). *J Abnorm Child Psychol* **29**: 541–556.
- Barratt W (2006). The Barratt Simplified Measure of Social Status. (Copy available online at: http://wbarratt.indstate.edu/social-class/Barratt_Simplified_Measure_of_Social_Status.pdf).
- Beck RC, Triplett MF (2009). Test-retest reliability of a group-administered paper-pencil measure of delay discounting. *Exp Clin Psychopharmacol* **17**: 345–355.
- Bellgrove M, Johnson K, Barry E, Mulligan A, Hawi Z, Gill M et al (2009). Dopaminergic haplotype as a predictor of spatial inattention in children with attention-deficit/hyperactivity disorder. *Arch Gen Psychiatry* **66**: 1135–1142.
- Biederman J, Kim JW, Doyle AE, Mick E, Fagerness J, Smoller JW et al (2008). Sexually dimorphic effects of four genes (COMT, SLC6A2, MAOA, SLC6A4) in genetic associations of ADHD: a preliminary study. *Am J Med Genet B Neuropsychiatr Genet* **147B**: 1511–1518.
- Boettiger CA, Mitchell JM, Tavares VC, Robertson M, Joslyn G, D'Esposito M et al (2007). Immediate reward bias in humans: fronto-parietal networks and a role for the catechol-O-methyltransferase 158(Val/Val) genotype. *J Neurosci* **27**: 14383–14391.
- Brookes K, Xu X, Chen W, Zhou K, Neale B, Lowe N et al (2006a). The analysis of 51 genes in DSM-IV combined type attention deficit hyperactivity disorder: association signals in DRD4, DAT1 and 16 other genes. *Mol Psychiatry* **11**: 934–953.
- Brookes KJ, Mill J, Guindalini C, Curran S, Xu X, Knight J et al (2006b). A common haplotype of the dopamine transporter gene associated with attention-deficit/hyperactivity disorder and interacting with maternal use of alcohol during pregnancy. *Arch Gen Psychiatry* **63**: 74–81.
- Bush G (2010). Attention-deficit/hyperactivity disorder and attention networks. *Neuropsychopharmacology* **35**: 278–300.
- Cardinal RN, Pennicott DR, Sugathapala CL, Robbins TW, Everitt BJ (2001). Impulsive choice induced in rats by lesions of the nucleus accumbens core. *Science* **292**: 2499–2501.
- Chapman GB, Winquist JR (1998). The magnitude effect: temporal discount rates and restaurant tips. *Psychon B Rev* **5**: 119–123.
- Chen J, Lipska BK, Halim N, Ma QD, Matsumoto M, Melhem S et al (2004). Functional analysis of genetic variation in catechol-O-methyltransferase (COMT): effects on mRNA, protein, and enzyme activity in postmortem human brain. *Am J Hum Genet* **75**: 807–821.
- Chen W, Zhou K, Sham P, Franke B, Kuntsi J, Campbell D et al (2008). DSM-IV combined type ADHD shows familial association with sibling trait scores: a sampling strategy for QTL linkage. *Am J Med Genet B Neuropsychiatr Genet* **147B**: 1450–1460.
- Conners CK, Sitarenios G, Parker JD, Epstein JN (1998). The revised Conners' Parent Rating Scale (CPRS-R): factor structure, reliability, and criterion validity. *J Abnorm Child Psychol* **26**: 257–268.
- Contini V, Marques FZC, Garcia CED, Hutz MH, Bau CHD (2006). MAOA-uVNTR polymorphism in a Brazilian sample: further support for the association with impulsive behaviors and alcohol dependence. *Am J Med Genet B Neuropsychiatr Genet* **141B**: 305–308.
- Cosi S, Vigil-Colet A, Canals J, Lorenzo-Seva U (2008). Psychometric properties of the Spanish adaptation of the Barratt Impulsiveness Scale-11-A for children. *Psychol Rep* **103**: 336–346.
- Costa PT, McCrae RR (1992). *Revised NEO Personality Inventory (NEO-PI-R) and NEO Five-Factor Inventory (NEO-FFI) Professional Manual*. Psychological Assessment Resources: Odessa, FL.
- Dandy KL, Gatch MB (2009). The effects of chronic cocaine exposure on impulsivity in rats. *Behav Pharmacol* **20**: 400–405.
- De Wit H (2009). Impulsivity as a determinant and consequence of drug use: a review of underlying processes. *Addict Biol* **14**: 22–31.
- Dempster EL, Mill J, Craig IW, Collier DA (2006). The quantification of COMT mRNA in post mortem cerebellum tissue: diagnosis, genotype, methylation and expression. *BMC Med Genet* **7**: 10.
- Diergaarde L, Pattij T, Poortvliet I, Hogenboom F, de Vries W, Schoffelmeer AN et al (2008). Impulsive choice and impulsive action predict vulnerability to distinct stages of nicotine seeking in rats. *Biol Psychiatry* **63**: 301–308.
- Durston S, Fossella JA, Mulder MJ, Casey BJ, Ziermans TB, Vessaz MN et al (2008). Dopamine transporter genotype conveys familial risk of attention-deficit/hyperactivity disorder through striatal activation. *J Am Acad Child Adolesc Psychiatry* **47**: 61–67.
- Egan MF, Goldberg TE, Kolachana BS, Callicott JH, Mazzanti CM, Straub RE et al (2001). Effect of COMT Val108/158 Met genotype

- on frontal lobe function and risk for schizophrenia. *Proc Natl Acad Sci USA* **98**: 6917–6922.
- Ettinger U, Joobar R, DE Guzman R, O'driscoll GA (2006). Schizotypy, attention deficit hyperactivity disorder, and dopamine genes. *Psychiatry Clin Neurosci* **60**: 764–767.
- Evenden JL (1999). Varieties of impulsivity. *Psychopharmacology (Berl)* **146**: 348–361.
- Fields S, Collins C, Leraas K, Reynolds B (2009). Dimensions of impulsive behavior in adolescent smokers and nonsmokers. *Exp Clin Psychopharmacol* **17**: 302–311.
- Forbes EE, Brown SM, Kimak M, Ferrell RE, Manuck SB, Hariri AR (2009). Genetic variation in components of dopamine neurotransmission impacts ventral striatal reactivity associated with impulsivity. *Mol Psychiatry* **14**: 60–70.
- Fossati A, Barratt ES, Acquarini E, Di Ceglie A (2002). Psychometric properties of an adolescent version of the Barratt Impulsiveness Scale-11 for a sample of Italian high school students. *Percept Mot Skills* **95**: 621–635.
- Frank MJ, Santamaria A, O'Reilly RC, Willcutt E (2007). Testing computational models of dopamine and noradrenaline dysfunction in attention deficit/hyperactivity disorder. *Neuropsychopharmacology* **32**: 1583–1599.
- Franke B, Hoogman M, Arias Vasquez A, Heister JGaM, Savelkoul PJ, Naber M et al (2008). Association of the dopamine transporter (SLC6A3/DAT1) gene 9-6 haplotype with adult ADHD. *Am J Med Genet B Neuropsychiatr Genet* **147B**: 1576–1579.
- Franke B, Vasquez AA, Johansson S, Hoogman M, Romanos J, Boreatti-Hümmer A et al (2010). Multicenter analysis of the SLC6A3/DAT1 VNTR haplotype in persistent ADHD suggests differential involvement of the gene in childhood and persistent ADHD. *Neuropsychopharmacology* **35**: 656–664.
- Gizer IR, Ficks C, Waldman ID (2009). Candidate gene studies of ADHD: a meta-analytic review. *Hum Genet* **126**: 51–90.
- Gogos JA, Morgan M, Luine V, Santha M, Ogawa S, Pfaff D et al (1998). Catechol-O-methyltransferase-deficient mice exhibit sexually dimorphic changes in catecholamine levels and behavior. *Proc Natl Acad Sci USA* **95**: 9991–9996.
- Goldberg TE, Egan MF, Gscheidle T, Coppola R, Weickert T, Kolachana BS et al (2003). Executive subprocesses in working memory: relationship to catechol-O-methyltransferase Val158Met genotype and schizophrenia. *Arch Gen Psychiatry* **60**: 889–896.
- Goldman-Rakic PS, Muly III EC, Williams GV (2000). D1 receptors in prefrontal cells and circuits. *Brain Res Rev* **31**: 295–301.
- Golimbet VE, Alfimova MV, Gritsenko IK, Ebstein RP (2007). Relationship between dopamine system genes and extraversion and novelty seeking. *Neurosci Behav Physiol* **37**: 601–606.
- Gothelf D, Michaelovsky E, Frisch A, Zohar AH, Presburger G, Burg M et al (2007). Association of the low-activity COMT 158Met allele with ADHD and OCD in subjects with velocardiofacial syndrome. *Int J Neuropsychopharmacol* **10**: 301–308.
- Green L, Myerson J (2004). A discounting framework for choice with delayed and probabilistic rewards. *Psychol Bull* **130**: 769–792.
- Hariri AR, Brown SM, Williamson DE, Flory JD, de Wit H, Manuck SB (2006). Preference for immediate over delayed rewards is associated with magnitude of ventral striatal activity. *J Neurosci* **26**: 13213–13217.
- Hollingshead AA (1975). *Four-Factor Index of Social Status*. Yale University: New Haven, CT.
- Huang YY, Cate SP, Battistuzzi C, Oquendo MA, Brent D, Mann JJ (2004). An association between a functional polymorphism in the monoamine oxidase a gene promoter, impulsive traits and early abuse experiences. *Neuropsychopharmacology* **29**: 1498–1505.
- Huotari M, Gogos JA, Karayiorgou M, Koponen O, Forsberg M, Raasmaja A et al (2002). Brain catecholamine metabolism in catechol-O-methyltransferase (COMT)-deficient mice. *Eur J Neurosci* **15**: 246–256.
- Johnson KA, Kelly SP, Bellgrove MA, Barry E, Cox M, Gill M et al (2007a). Response variability in attention deficit hyperactivity disorder: evidence for neuropsychological heterogeneity. *Neuropsychologia* **45**: 630–638.
- Johnson KA, Robertson IH, Kelly SP, Silk TJ, Barry E, Daibhis A et al (2007b). Dissociation in performance of children with ADHD and high-functioning autism on a task of sustained attention. *Neuropsychologia* **45**: 2234–2245.
- Kable JW, Glimcher PW (2007). The neural correlates of subjective value during intertemporal choice. *Nat Neurosci* **10**: 1625–1633.
- Kable JW, Glimcher PW (2009). The neurobiology of decision: consensus and controversy. *Neuron* **63**: 733–745.
- Kheramin S, Body S, Ho MY, Velazquez-Martinez DN, Bradshaw CM, Szabadi E et al (2004). Effects of orbital prefrontal cortex dopamine depletion on inter-temporal choice: a quantitative analysis. *Psychopharmacology (Berl)* **175**: 206–214.
- Kirby KN (2009). One-year temporal stability of delay-discount rates. *Psychon Bull Rev* **16**: 457–462.
- Kirby KN, Marakovi NN (1996). Delay-discounting probabilistic rewards: rates decrease as amounts increase. *Psycho Bull Rev* **3**: 100–104.
- Kobayashi S, Schultz W (2008). Influence of reward delays on responses of dopamine neurons. *J Neurosci* **28**: 7837–7846.
- Konrad K, Gauggel S, Manz A, Scholl M (2000). Lack of inhibition: a motivational deficit in children with attention deficit/hyperactivity disorder and children with traumatic brain injury. *Child Neuropsychol* **6**: 286–296.
- Krause KH, Dresel SH, Krause J, Kung HF, Tatsch K (2000). Increased striatal dopamine transporter in adult patients with attention deficit hyperactivity disorder: effects of methylphenidate as measured by single photon emission computed tomography. *Neurosci Lett* **285**: 107–110.
- Krishnan-Sarin S, Reynolds B, Duhig AM, Smith A, Liss T, McFetridge A et al (2007). Behavioral impulsivity predicts treatment outcome in a smoking cessation program for adolescent smokers. *Drug Alcohol Depend* **88**: 79–82.
- Kuntsi J, Neale BM, Chen W, Faraone SV, Asherson P (2006a). The IMAGE project: methodological issues for the molecular genetic analysis of ADHD. *Behav Brain Funct* **2**: 27.
- Kuntsi J, Rogers H, Swinard G, Borger N, van der Meere J, Rijdsdijk F et al (2006b). Reaction time, inhibition, working memory and 'delay aversion' performance: genetic influences and their interpretation. *Psychol Med* **36**: 1613–1624.
- Kuntsi J, Wood AC, Rijdsdijk F, Johnson KA, Andreou P, Albrecht B et al (in press). Separation of cognitive impairments in attention deficit hyperactivity disorder into two familial factors. *Arch Gen Psychiatry*.
- Kuntsi J, Wood AC, Rijdsdijk F, Johnson KA, Andreou P, Albrecht B (2009). Why cognitive performance in ADHD may not reveal true potential: findings from a large population-based sample. *J Int Neuropsychol Soc* **15**: 570–579.
- Lachman HM, Papolos DF, Saito T, Yu YM, Szumlanski CL, Weinshilboum RM (1996). Human catechol-O-methyltransferase pharmacogenetics: description of a functional polymorphism and its potential application to neuropsychiatric disorders. *Pharmacogenetics* **6**: 243–250.
- Lee B, London ED, Poldrack RA, Farahi J, Nacca A, Monterosso JR et al (2009). Striatal dopamine d2/d3 receptor availability is reduced in methamphetamine dependence and is linked to impulsivity. *J Neurosci* **29**: 14734–14740.
- Leshem R, Glicksohn J (2007). The construct of impulsivity revisited. *Pers Individ Differ* **43**: 681–691.
- Lewis DA, Melchitzky DS, Sesack SR, Whitehead RE, Auh S, Sampson A (2001). Dopamine transporter immunoreactivity in monkey cerebral cortex: regional, laminar, and ultrastructural localization. *J Comp Neurol* **432**: 119–136.
- Logue AW (1988). Research on self-control: an integrating framework. *Behav Brain Sci* **11**: 665–709.

- Lotta T, Vidgren J, Tilgmann C, Ulanen I, Melen K, Julkunen I et al (1995). Kinetics of human soluble and membrane-bound catechol O-methyltransferase: a revised mechanism and description of the thermolabile variant of the enzyme. *Biochemistry (Mosc)* **34**: 4202–4210.
- Luman M, Tripp G, Scheres A (2010). Identifying the neurobiology of altered reinforcement sensitivity in ADHD: a review and research agenda. *Neurosci Biobehav Rev* **34**: 744–754.
- Marco R, Miranda A, Schlotz W, Melia A, Mulligan A, Muller U et al (2009). Delay and reward choice in ADHD: an experimental test of the role of delay aversion. *Neuropsychology* **23**: 367–380.
- Matsumoto M, Weickert CS, Beltaifa S, Kolachana B, Chen J, Hyde TM et al (2003). Catechol O-methyltransferase (COMT) mRNA expression in the dorsolateral prefrontal cortex of patients with schizophrenia. *Neuropsychopharmacology* **28**: 1521–1530.
- Mazei MS, Pluto CP, Kirkbride B, Pehek EA (2002). Effects of catecholamine uptake blockers in the caudate-putamen and subregions of the medial prefrontal cortex of the rat. *Brain Res* **936**: 58–67.
- McClure SM, Ericson KM, Laibson DI, Loewenstein G, Cohen JD (2007). Time discounting for primary rewards. *J Neurosci* **27**: 5796–5804.
- McClure SM, Laibson DI, Loewenstein G, Cohen JD (2004). Separate neural systems value immediate and delayed monetary rewards. *Science* **306**: 503–507.
- McDonald J, Schleifer L, Richards JB, de Wit H (2003). Effects of THC on behavioral measures of impulsivity in humans. **28**: 1356–1365.
- McInerney RJ, Kerns KA (2003). Time reproduction in children with ADHD: motivation matters. *Child Neuropsychol* **9**: 91–108.
- Meda SA, Stevens MC, Potenza MN, Pittman B, Gueorguieva R, Andrews MM et al (2009). Investigating the behavioral and self-report constructs of impulsivity domains using principal component analysis. *Behav Pharmacol* **20**: 390–399.
- Melanko S, Leraas K, Collins C, Fields S, Reynolds B (2009). Characteristics of psychopathy in adolescent nonsmokers and smokers: relations to delay discounting and self reported impulsivity. *Exp Clin Psychopharmacol* **17**: 258–265.
- Michaelovsky E, Gothelf D, Korostishevsky M, Frisch A, Burg M, Carmel M et al (2008). Association between a common haplotype in the COMT gene region and psychiatric disorders in individuals with 22q11.2DS. *Int J Neuropsychopharmacol* **11**: 351–363.
- Moeller FG, Barratt ES, Dougherty DM, Schmitz JM, Swann AC (2001). Psychiatric aspects of impulsivity. *Am J Psychiatry* **158**: 1783–1793.
- Morón JA, Brockington A, Wise RA, Rocha BA, Hope BT (2002). Dopamine uptake through the norepinephrine transporter in brain regions with low levels of the dopamine transporter: evidence from knock-out mouse lines. *J Neurosci* **22**: 389–395.
- Munafò MR, Yalcin B, Willis-Owen SA, Flint J (2008). Association of the dopamine D4 receptor (DRD4) gene and approach-related personality traits: meta-analysis and new data. *Biol Psychiatry* **63**: 197–206.
- Myerson J, Green L (1995). Discounting of delayed rewards—models of individual choice. *J Exp Anal Behav* **64**: 263–276.
- Myerson J, Green L, Warusawitharana M (2001). Area under the curve as a measure of discounting. *J Exp Anal Behav* **76**: 235–243.
- O'Connell RG, Bellgrove MA, Dockree PM, Lau A, Fitzgerald M, Robertson IH (2008). Self-alert training: volitional modulation of autonomic arousal improves sustained attention. *Neuropsychologia* **46**: 1379–1390.
- Ohmura Y, Takahashi T, Kitamura N, Wehr P (2006). Three-month stability of delay and probability discounting measures. *Exp Clin Psychopharmacol* **14**: 318–328.
- Pálmason H, Moser D, Sigmund J, Vogler C, Hänig S, Schneider A et al (2010). Attention-deficit/hyperactivity disorder phenotype is influenced by a functional catechol-O-methyltransferase variant. *J Neural Transm* **117**: 259–267.
- Paloyelis Y, Asherson P, Kuntsi J (2009). Are ADHD symptoms associated with delay aversion or choice impulsivity? A general population study. *J Am Acad Child Adolesc Psychiatry* **48**: 837–846.
- Paloyelis Y, Mehta MA, Kuntsi J, Asherson P (2007). Functional MRI in ADHD: a systematic literature review. *Expert Rev Neurother* **7**: 1337–1356.
- Patton JH, Stanford MS, Barratt ES (1995). Factor structure of the Barratt impulsiveness scale. *J Clin Psychol* **51**: 768–774.
- Perry JL, Larson EB, German JP, Madden GJ, Carroll ME (2005). Impulsivity (delay discounting) as a predictor of acquisition of IV cocaine self-administration in female rats. *Psychopharmacology (Berl)* **178**: 193–201.
- Perry JL, Nelson SE, Carroll ME (2008). Impulsive choice as a predictor of acquisition of IV cocaine self-administration and reinstatement of cocaine-seeking behavior in male and female rats. *Exp Clin Psychopharmacol* **16**: 165–177.
- Pothuizen HH, Jongen-Relo AL, Feldon J, Yee BK (2005). Double dissociation of the effects of selective nucleus accumbens core and shell lesions on impulsive-choice behaviour and salience learning in rats. *Eur J Neurosci* **22**: 2605–2616.
- Poulos CX, Le AD, Parker JL (1995). Impulsivity predicts individual susceptibility to high levels of alcohol self-administration. *Behav Pharmacol* **6**: 810–814.
- Qian Q, Wang Y, Zhou R, Li J, Wang B, Glatt S et al (2003). Family-based and case-control association studies of catechol-O-methyltransferase in attention deficit hyperactivity disorder suggest genetic sexual dimorphism. *Am J Med Genet B Neuropsychiatr Genet* **118B**: 103–109.
- Rachlin H, Green L (1972). Commitment, choice and self-control. *J Exp Anal Behav* **17**: 15–22.
- Reuter M, Kirsch P, Hennig J (2006). Inferring candidate genes for attention deficit hyperactivity disorder (ADHD) assessed by the World Health Organization Adult ADHD Self-Report Scale (ASRS). *J Neural Transm (Vienna)* **113**: 929–938.
- Reynolds B (2006a). The experiential discounting task is sensitive to cigarette-smoking status and correlates with a measure of delay discounting. *Behav Pharmacol* **17**: 133–142.
- Reynolds B (2006b). A review of delay-discounting research with humans: relations to drug use and gambling. *Behav Pharmacol* **17**: 651–667.
- Reynolds B, Penfold RB, Patak M (2008). Dimensions of impulsive behavior in adolescents: laboratory behavioral assessments. *Exp Clin Psychopharmacol* **16**: 124–131.
- Reynolds B, Richards JB, de Wit H (2006). Acute-alcohol effects on the experiential discounting task (EDT) and a question-based measure of delay discounting. *Pharmacol Biochem Behav* **83**: 194–202.
- Reynolds B, Schiffbauer R (2004). Measuring state changes in human delay discounting: an experiential discounting task. *Behav Processes* **67**: 343–356.
- Richards JB, Zhang L, Mitchell SH, de Wit H (1999). Delay or probability discounting in a model of impulsive behavior: effect of alcohol. *J Exp Anal Behav* **71**: 121–143.
- Rommelse NNJ, Altink ME, Arias-Vásquez A, Buschgens CJM, Fliers E, Faraone SV et al (2008). A review and analysis of the relationship between neuropsychological measures and DAT1 in ADHD. *Am J Med Genet B Neuropsychiatr Genet* **147B**: 1536–1546.
- Sagvolden T, Johansen EB, Aase H, Russell VA (2005). A dynamic developmental theory of attention-deficit/hyperactivity disorder (ADHD) predominantly hyperactive/impulsive and combined subtypes. *Behav Brain Sci* **28**: 397–419.
- Scheres A, Dijkstra M, Ainslie E, Balkan J, Reynolds B, Sonuga-Barke E et al (2006). Temporal and probabilistic discounting of rewards in children and adolescents: effects of age and ADHD symptoms. *Neuropsychologia* **44**: 2092–2103.

- Scheres A, Tontsch C, Thoeny AL, Kaczurkin A (2010). Temporal reward discounting in attention-deficit/hyperactivity disorder: the contribution of symptom domains, reward magnitude, and session length. *Biol Psychiatry* 67: 641–648.
- Shiels K, Hawk LW, Reynolds B, Mazzullo RJ, Rhodes JD, Pelham WE et al (2009). Effects of methylphenidate on discounting of delayed rewards in attention deficit/hyperactivity disorder. *Exp Clin Psychopharmacol* 17: 291–301.
- Slusarek M, Velling S, Bunk D, Eggers C (2001). Motivational effects on inhibitory control in children with ADHD. *J Am Acad Child Adolesc Psychiatry* 40: 355–363.
- Smith CL, Hantula DA (2008). Methodological considerations in the study of delay discounting in intertemporal choice: a comparison of tasks and modes. *Behav Res Methods* 40: 940–953.
- Sonuga-Barke EJS (2005). Causal models of attention-deficit/hyperactivity disorder: from common simple deficits to multiple developmental pathways. *Biol Psychiatry* 57: 1231–1238.
- Spencer TJ, Biederman J, Madras BK, Dougherty DD, Bonab AA, Livni E et al (2007). Further evidence of dopamine transporter dysregulation in ADHD: a controlled PET imaging study using altopane. *Biol Psychiatry* 62: 1059–1061.
- Stanford MS, Mathias CW, Dougherty DM, Lake SL, Anderson NE, Patton JH (2009). Fifty years of the Barratt Impulsiveness Scale: an update and review. *Pers Individ Differ* 47: 385–395.
- Tripp G, Wickens JR (2008). Research review: dopamine transfer deficit: a neurobiological theory of altered reinforcement mechanisms in ADHD. *J Child Psychol Psychiatry* 49: 691–704.
- Tunbridge EM, Bannerman DM, Sharp T, Harrison PJ (2004). Catechol-o-methyltransferase inhibition improves set-shifting performance and elevates stimulated dopamine release in the rat prefrontal cortex. *J Neurosci* 24: 5331–5335.
- Uebel H, Albrecht B, Asherson P, Borger NA, Butler L, Chen W et al (2010). Performance variability, impulsivity errors and the impact of incentives as gender-independent endophenotypes for ADHD. *J Child Psychol Psychiatry* 51: 210–218.
- van de Giessen EM, de Win MM, Tanck MW, van den Brink W, Baas F, Booi J (2009). Striatal dopamine transporter availability associated with polymorphisms in the dopamine transporter gene SLC6A3. *J Nucl Med* 50: 45–52.
- Volavka J, Bilder R, Nolan K (2004). Catecholamines and aggression: the role of COMT and MAO polymorphisms. *Ann NY Acad Sci* 1036: 393–398.
- Volkow ND, Wang GJ, Kollins SH, Wigal TL, Newcorn JH, Telang F et al (2009). Evaluating dopamine reward pathway in ADHD: clinical implications. *JAMA* 302: 1084–1091.
- Volkow ND, Wang GJ, Newcorn J, Telang F, Solanto MV, Fowler JS et al (2007). Depressed dopamine activity in caudate and preliminary evidence of limbic involvement in adults with attention-deficit/hyperactivity disorder. *Arch Gen Psychiatry* 64: 932–940.
- von Diemen L, Szobot CM, Kessler F, Pechansky F (2007). Adaptation and construct validation of the Barratt Impulsiveness Scale (BIS 11) to Brazilian Portuguese for use in adolescents. *Rev Bras Psiquiatr* 29: 153–156.
- Wechsler D (1991). *Wechsler Intelligence Scale for Children*, 3rd edn. The Psychological Corporation: London.
- Wechsler D (1997). *Wechsler Intelligence Scale for Adults*. The Psychological Corporation: London.
- Wilhelm CJ, Mitchell SH (2009). Strain differences in delay discounting using inbred rats. *Genes Brain Behav* 8: 426–434.
- Willcutt E, Sonuga-Barke E, Nigg J, Sergeant J (2008). Recent developments in neuropsychological models of childhood psychiatric disorders. In: Banaschewski T, Rhode LA (eds). *Biological Child Psychiatry. Recent Trends and Developments*, vol. 24. Karger: Basel. pp 195–226.
- Williams-Gray CH, Hampshire A, Robbins TW, Owen AM, Barker RA (2007). Catechol O-methyltransferase Val158Met genotype influences frontoparietal activity during planning in patients with Parkinson's disease. *J Neurosci* 27: 4832–4838.
- Winstanley CA, Theobald DEH, Dalley JW, Cardinal RN, Robbins TW (2006). Double dissociation between serotonergic and dopaminergic modulation of medial prefrontal and orbitofrontal cortex during a test of impulsive choice. *Cereb Cortex* 16: 106–114.
- Wood AC, Asherson P, Rijdsdijk F, Kuntsi J (2009). Is overactivity a core feature in ADHD? Familial and receiver operating characteristic curve analysis of mechanically assessed activity level. *J Am Acad Child Adolesc Psychiatry* 10: 1023–1030.

Supplementary Information accompanies the paper on the Neuropsychopharmacology website (<http://www.nature.com/npp>)