

## Commentary

## Is there Evidence for Neurotoxicity in the Prodromal and Early Stages of Schizophrenia?

Adrienne C Lahti\*<sup>1</sup> and Meredith A Reid<sup>1,2</sup><sup>1</sup>Department of Psychiatry and Behavioral Neurobiology, The University of Alabama at Birmingham, Birmingham, AL, USA;<sup>2</sup>Department of Biomedical Engineering, The University of Alabama at Birmingham, Birmingham, AL, USA

Neuropsychopharmacology (2011) 36, 1779–1780; doi:10.1038/npp.2011.83

Retrospective and prospective studies in schizophrenia have identified a period of time, termed the prodromal period, that precedes the first episode of frank psychosis by a variable length of time (days–years) and is characterized by subthreshold psychotic symptoms. Importantly, both the prodromal phase and the early stages of the schizophrenia illness are associated with significant loss in social and intellectual abilities (Yung *et al*, 2005), as well as gray matter volume (Cahn *et al*, 2006; Pantelis *et al*, 2003). In addition, patients' response to antipsychotic medication decreases from first to subsequent episodes (Agid *et al*, 2011). It has been hypothesized that a neurobiological process might underlie this early progressive deterioration. In order to gain insight into this period, researchers have identified and followed subjects at 'ultra high risk', either because of a strong family history of schizophrenia or because of the presence of subthreshold symptoms. In addition, studies of first-episode subjects have become more common, in part because they lack the confounding effect of antipsychotic medication treatment.

In this issue of *Neuropsychopharmacology*, de la Fuente-Sandoval *et al* (2011) use <sup>1</sup>H magnetic resonance spectroscopy (MRS) to obtain glutamate measurements in the caudate nucleus of antipsychotic-naïve prodromal subjects, antipsychotic-naïve first-episode psychosis subjects, and matched healthy control subjects. These investigators report that the high-risk and first-episode groups showed higher levels of glutamate than controls, without differences between ultra high risk and first episode. These findings are noteworthy for several reasons. Few studies have compared ultra high-risk and first-episode populations cross-sectionally. Because only 15–35% of high-risk subjects transition to psychosis, the interpretation of purely cross-sectional studies is somewhat ambiguous. Yet, in this study, the identification of the same pattern of high

glutamate levels in the high-risk and first-episode populations provides stronger support for the relevance of this abnormality. Clearly, prospective studies of ultra high-risk populations are needed to address whether the degree of this early abnormality is predictive of the illness. Nevertheless, this study suggests that high levels of glutamate measured in the precommissural dorsal-caudate nucleus, a region with prominent projections throughout the cortical mantle, could induce cortical neuronal toxicity leading to a progressive functional and intellectual deterioration. Because there is preliminary evidence that stable medicated subjects either do not show a difference or evidence a decrease in glutamate levels compared with normal subjects (de la Fuente-Sandoval *et al*, 2011; Reid *et al*, 2010), it is tempting to hypothesize that a time-limited neurotoxic process may characterize the prodromal period and the early stages of the illness. Although more research is needed in this area, this suggests the possibility of curtailing functional deterioration by intervening therapeutically during the early stages. Importantly, this study provides strong clues as to the nature of these interventions, as well as specific targets, that is, glutamate level, to monitor treatment intervention. This study also points towards potential prediction algorithms to identify those who might, or might not, go on to develop the illness.

These data need to be considered in light of a recent study that reported increased striatal <sup>18</sup>F-dopa uptake, which reflects increased presynaptic dopaminergic function, in another group of ultra –high-risk subjects (Howes *et al*, 2009). These data parallel findings of increased presynaptic dopamine function in psychotic and first-episode subjects. Do glutamatergic and dopaminergic dysfunctions coexist in the prodromal period, and is one the consequence of the other? In the caudate, both glutamatergic and dopaminergic afferents synapse on the same dendritic spines and shafts of medium-sized GABA-ergic projection neurons, providing an ideal anatomical arrangement for interactions between these neurotransmitters (Sesack, 2010). Alternatively, because of the use of different operational definitions of the at-risk population, the studies of Howes *et al* (2009) and de la Fuente-Sandoval *et al* (2011) may have enrolled

\*Correspondence: Dr AC Lahti, Department of Psychiatry and Behavioral Neurobiology, The University of Alabama at Birmingham, 1530 3<sup>rd</sup> AVE S, Birmingham, AL 35294-0017, USA, Tel: +1 205 996 6776, Fax: +1 205 975 4879, E-mail: alahti@uab.edu

Received 19 April 2011; accepted 19 April 2011

slightly different populations. Nevertheless, these studies emphasize the need for prospective multimodal brain imaging of the prodromal period. The stakes are high. Any breakthrough in the neurobiology of the prodromal period could dramatically impact the life of subjects with schizophrenia.

Finally, the study of de la Fuente-Sandoval *et al* (2011) re-emphasizes the role of MRS technology for the non-invasive investigation of the brain's neurochemistry. It needs to be remembered that there are a number of limitations to this technique, such as factors that can confound measurements, including but not limited to age, nicotine and substances of abuse, and psychotropic medications (Licata and Renshaw, 2010). Thus, more than ever, caution will need to be exercised when matching populations. Nonetheless, with the introduction of more powerful magnets (that is, 7T), MRS will acquire greater precision and become an even more important tool to characterize, *in vivo*, key elements underlying the neuropathology of psychiatric conditions, especially with regard to glutamate, and to assess novel treatment strategies.

## DISCLOSURE

ACL receives research funds from the National Institute of Health grants R01 MH081014, R01 NCI141663, R01 AG021927, and an investigator-initiated grant from Pfizer. MAR reports no biomedical financial interests or potential conflicts of interest.

## REFERENCES

- Agid O, Arenovich T, Sajeew G, Zipursky RB, Kapur S, Foussias G *et al* (2011). An algorithm-based approach to first-episode schizophrenia: response rates over 3 prospective antipsychotic trials with a retrospective data analysis. *J Clin Psychiatry*; e-pub ahead of print 8 March 2011 (PMID: 21457676).
- Cahn W, van Haren NEM, Hulshoff Pol HE, Schnack HG, Caspers E, Laponder DAJ *et al* (2006). Brain volume changes in the first year of illness and 5-year outcome of schizophrenia. *Br J Psychiatry* 189: 381–382.
- de la Fuente-Sandoval C, Leon-Ortiz P, Favila R, Stephano S, Mamo D, Ramirez-Bermudez J *et al* (2011). Higher levels of glutamate in the associative-striatum of subjects with prodromal symptoms of schizophrenia and patients with first-episode psychosis. *Neuropsychopharmacology* 36: 1781–1791.
- Howes OD, Montgomery AJ, Asselin MC, Murray RM, Valli I, Tabraham P *et al* (2009). Elevated striatal dopamine function linked to prodromal signs of schizophrenia. *Arch Gen Psychiatry* 66: 13–20.
- Licata SC, Renshaw PF (2010). Neurochemistry of drug action: insights from proton magnetic resonance spectroscopic imaging and their relevance to addiction. *Ann N Y Acad Sci* 1187: 148–171.
- Pantelis C, Velakoulis D, McGorry P, Wood S, Suckling J, Phillips L *et al* (2003). Neuroanatomical abnormalities before and after onset of psychosis: a cross-sectional and longitudinal MRI comparison. *Lancet* 361: 281–288.
- Reid MA, Stoeckel LE, White DM, Avsar KB, Bolding MS, Akella NS *et al* (2010). Assessments of function and biochemistry of the anterior cingulate cortex in schizophrenia. *Biol Psychiatry* 68: 625–633.
- Sesack SR (2010). Functional implications of dopamine D2 receptor localization in relation to glutamate neurons. In: Iversen LL, Iversen SD, Dunnett SB, Bjorklund A (eds). *Dopamine Handbook*. Oxford University Press: New York. pp 22–37.
- Yung AR, Yuen HP, McGorry PD, Phillips LJ, Kelly D, Dell'Olio M *et al* (2005). Mapping the onset of psychosis: the Comprehensive Assessment of At-Risk Mental States. *Aust N Z J Psychiatry* 39: 964–971.