Exercise and Sodium Butyrate Transform a Subthreshold Learning Event into Long-Term Memory via a Brain-Derived Neurotrophic factor-Dependent Mechanism

Karlie A Intlekofer¹, Nicole C Berchtold^{*1}, Melissa Malvaez², Anthony J Carlos¹, Susan C McQuown², Michael J Cunningham³, Marcelo A Wood² and Carl W Cotman¹

¹Institute for Memory Impairments and Neurological Disorders, University of California Irvine, Irvine, CA, USA; ²Department of Neurobiology and Behavior, Center for the Neurobiology of Learning and Memory, University of California, Irvine, CA, USA; ³University of Vermont College of Medicine, Burlington, VT, USA

We demonstrate that exercise enables hippocampal-dependent learning in conditions that are normally subthreshold for encoding and memory formation, and depends on hippocampal induction of brain-derived neurotrophic factor (BDNF) as a key mechanism. Using a weak training paradigm in an object location memory (OLM) task, we show that sedentary mice are unable to discriminate 24 h later between familiar and novel object locations. In contrast, 3 weeks of prior voluntary exercise enables strong discrimination in the spatial memory task. Cognitive benefits of exercise match those attained with post-training sodium butyrate (NaB), a histone deacetylase (HDAC) inhibitor previously shown to enable subthreshold learning. We demonstrate that the enabling effects of exercise and NaB on subthreshold OLM learning are dependent on hippocampal BDNF upregulation, and are blocked by hippocampal infusion of BDNF short-interfering RNA. Exercise and NaB increased *bdnf* transcripts I and IV, and the increases were associated with BDNF promoter acetylation on H4K8 but not H4K12. These data provide support for the concept that exercise engages epigenetic control mechanisms and serves as a natural stimulus that operates in part like NaB and potentially other HDAC inhibitors, placing the brain into a state of readiness for plasticity.

Neuropsychopharmacology (2013) **38**, 2027–2034; doi:10.1038/npp.2013.104; published online 15 May 2013

Keywords: object location memory; hippocampus; siRNA; subthreshold; plasticity

INTRODUCTION

Exercise is well established to improve brain health and function in both humans and other animals (Cotman *et al*, 2007; Middleton *et al*, 2010; Ahlskog *et al*, 2011; Voss *et al*, 2011). Animal studies reveal improved performance in numerous hippocampal-dependent tasks following several days to weeks of exercise participation (Cotman and Berchtold, 2002; Cotman *et al*, 2007; Bekinschtein *et al*, 2011). An idea that has not been tested is that exercise may enable hippocampal-dependent learning in conditions that are normally subthreshold for encoding and memory formation. In particular, a weak training paradigm that is normally insufficient (subthreshold) for encoding may become sufficient for long-term memory formation in an animal that has engaged in physical activity.

Exercise induces multiple mechanisms in the hippocampus that support the possibility that exercise may improve the acquisition and consolidation of normally subthreshold training. For example, exercise increases hippocampal expression of several plasticity-related growth factors including brain-derived neurotrophic factor (BDNF; Neeper *et al*, 1996; Cotman and Berchtold, 2002), thought to be a central mechanism mediating beneficial effects of exercise on hippocampal-dependent cognition (Cotman *et al*, 2007). BDNF, which is essential for certain forms of learning and memory, enhances synaptic plasticity by triggering functional and structural changes in neurons and synapses (Lu *et al*, 2008; Gottmann *et al*, 2009), such as facilitating the induction of long-term potentiation (LTP) (Rex *et al*, 2007), a synaptic correlate of learning and memory. Previously we have suggested that BDNF induction with exercise serves to place the brain into a state of readiness for plasticity (Cotman *et al*, 2007), an idea we test in this study using a subthreshold learning paradigm.

The induction of BDNF with exercise may involve epigenetic mechanisms, by which dynamic and reversible changes to chromatin structure modify the accessibility of a gene to the molecular machinery regulating transcription. One such epigenetic mechanism involves regulation of a

*Correspondence: Dr NC Berchtold, Institute for Memory Impairments and Neurological Disorders, University of California Irvine, 1206 Gillespie Building, Irvine, CA 92697-4260, USA, Tel: +1 949 824 6071, Fax: +1 949 824 2071, E-mail: nberchto@uci.edu
Received 5 March 2013; revised 12 April 2013; accepted 14 April 2013; accepted article preview online 24 April 2013

with 21-day running wheel access, receiving vehicle (Ex/Veh). Tissue limitations required that acetylation assessment was conducted with half of the samples each for H4K8 and H4K12 acetylation assays (eg, $n=4$ per group). Eight mice that were not exposed to handling or injections were used as negative controls in chromatin immunoprecipitation (ChIP) assays to establish baseline acetylation levels. For the exercise arm of the siRNA study, three treatment groups were used ($n=12$ per group): Sed/Control siRNA, Ex/Control siRNA, and Ex/*bdnf* siRNA. For the NaB arm of the siRNA study, the two treatment groups ($n=10$ per group) were: (1) Sed/NaB/Control siRNA and (2) Sed/NaB/*bdnf* siRNA. For detailed methods see the Supplementary Information available at the *Neuropsychopharmacology* website.

Quantitative PCR and Acetylation Assay

Gene expression for total *bdnf* and specific *bdnf* transcripts was assessed by quantitative PCR following reverse transcription (RT-qPCR). Total *bdnf* primers were designed to the common 3' coding exon. Primers for *bdnf* transcripts I, IV, and VI correspond to each unique exon sequence (Aid *et al* 2007; Supplementary Figure S1). RT-qPCR primer sets (Supplementary Table S1) were designed using the Roche Universal Probe Library Assay Design Center and obtained from Integrated DNA Technologies (Coralville, IA). RT-qPCR reactions were run in a Stratagene MX3005P thermocycler at 95 °C for 3 min, followed by 45 cycles of 95 °C for 10 s, and 58 °C for 15 s. Each RT-qPCR run included all samples run in triplicate and a standard curve. Data were analyzed by the $2^{-\Delta\Delta C_t}$ method and expressed as fold change over control after normalizing with input samples, as described previously (Sahar *et al*, 2007). ChIP assay was followed by RT-qPCR for specific BDNF promoters to assess histone acetylation (H4K8 and H4K12) at *bdnf* promoter regions (see Supplementary Table 2 for RT-qPCR primer sequences used in conjunction with ChIP). Chromatin extraction from hippocampal tissue was performed using the 'EZ-Magna ChIP-A Chromatin Immunoprecipitation Kit' (Millipore, Billerica, MA), following the manufacturer's protocol. The following antibodies were used for immunoprecipitation: anti-acetyl histone H4 lysine 8 (H4K8Ac; Millipore), anti-acetyl histone H4 lysine 12 (H4K12Ac; Abcam, Cambridge, MA), anti-acetylated H3 (positive control; Millipore), and non-immune rabbit IgG (negative control; Millipore).

Statistics

Treatment effects were detected using one-way analysis of variance, followed by *post hoc* Bonferroni's *t*-tests to delineate between-group differences.

For additional details, see Supplementary Materials and Methods at the *Neuropsychopharmacology* website.

RESULTS

Exercise Enables Learning in a Training Paradigm that is Normally Subthreshold for Long-Term Memory

Previously it has been demonstrated that a 3-min training session in a location-dependent OLM task is normally

Figure 2 In a subthreshold object location memory (OLM) paradigm, exercise (Ex) improves learning equivalent to sedentary (Sed) mice injected with post-acquisition sodium butyrate (NaB), a histone deacetylase (HDAC) inhibitor. (a) Twenty-four hours after a 3-min acquisition trial, vehicle-treated Sed animals do not show preference for the novel location over the familiar location, as measured by the discrimination index ratio (DI). Following 3 weeks voluntary exercise, Ex animals show marked preference for the novel location to a similar extent as post-acquisition injection of NaB. (b) Quantitative reverse transcription PCR (RT-qPCR) revealed that hippocampal brain-derived neurotrophic factor (*bdnf*) mRNA levels are increased by exercise and NaB, in parallel with the cognitive enhancement. (c) Effects of NaB on cognition persist 7 days after the acquisition trial, whereas the exercise effects on spatial memory in the subthreshold paradigm eventually decay after daily physical activity ceases. (d) RT-qPCR revealed that *bdnf* mRNA expression patterns at the 7-day timepoint paralleled the cognitive data across the treatment groups, remaining elevated after NaB treatment but not with exercise. Data are expressed as means $\pm$ SEM. For the 24-h timepoint: ** $p < 0.01$ vs sedentary. For the 7-day timepoint: $\sigma p < 0.05$; $\sigma\sigma\sigma p < 0.001$ vs sedentary.

insufficient, or subthreshold, to allow for short- or long-term memory formation in sedentary animals (Stefanko *et al*, 2009; McQuown *et al*, 2011). We investigated the possibility that exercise may enable learning in subthreshold conditions. Mice were provided free access to running wheels for 3 weeks, followed by subthreshold OLM training (3 min), and 24 h later, assessment of long-term memory. In parallel, we compare the efficacy of 3 weeks exercise with post-training NaB treatment, an HDAC inhibitor previously been shown to enable consolidation of the subthreshold event (see Figure 1a for Experimental Design).

The 24-h retention test revealed a significant effect of treatment on OLM performance ($F_{(2,21)} = 10.14$, $p < 0.001$). Confirming that the training paradigm is normally subthreshold for learning, sedentary mice were unable to discriminate between familiar and novel object locations in the 24-h retention test (Figure 2a). In contrast, exercised animals showed significant discrimination and long-term memory, with exercise similarly effective as post-training

injection of NaB (Figure 2a). These data reveal that exercise enables learning in conditions that are normally subthreshold for memory formation, and that exercise facilitates encoding to a similar extent as NaB, a memory-enhancing drug.

Exercise- and NaB-induced Gains in OLM Performance are Associated with *bdnf* Transcription

We next investigated the possibility that a common molecular mechanism may be engaged by exercise and NaB that enables encoding of normally subthreshold learning conditions. Specifically, we evaluated whether exercise and NaB induce hippocampal transcription of *bdnf*, a plasticity gene central to hippocampal-dependent learning. Hippocampal *bdnf* expression was evaluated 1 h after the 24-hour OLM retention test by RT-qPCR. There was a significant effect of treatment on *bdnf* gene expression ($F_{(2,21)} = 11.99$, $p < 0.001$), with both exercise and NaB significantly increasing *bdnf* mRNA over sedentary levels (1.60-fold and 1.37-fold, respectively; Figure 2b).

As transcription of *bdnf* is regulated by several exons, of which *bdnf* I, IV, and VI are the most highly expressed transcripts in the hippocampus, expression levels of *bdnf* I, IV, and VI transcripts were also evaluated. We found a significant effect of treatment on *bdnf* I ($F_{(2,21)} = 4.80$, $p < 0.05$) and *bdnf* IV ($F_{(2,21)} = 4.02$, $p < 0.05$), but not *bdnf* VI. *Post hoc* tests revealed that exercise and NaB treatments induced *bdnf* I (Supplementary Figure S2A) and *bdnf* IV (Supplementary Figure S2B), whereas neither treatment increased *bdnf* VI gene expression (Supplementary Figure S2C). These data indicate that hippocampal induction of *bdnf* mRNA (specifically *bdnf* I and *bdnf* IV) is associated with the successful encoding and consolidation, following exercise or NaB, of a normally subthreshold task.

Behavioral and Molecular Effects of NaB (but not Exercise) Remain Present in a 7-Day Retention Test

It has previously been demonstrated that post-training NaB leads to long-term memory formation for novel objects that persists at least 7 days, lasting beyond the point at which normal memory fails (Stefanko et al, 2009). However, it is unknown whether NaB induces a similar persistence of memory for the hippocampal-dependent OLM task, and the relationship to BDNF has not been explored. Here, we determine whether the behavioral and molecular effects of NaB and exercise are long lasting, by evaluating OLM and *bdnf* expression 7 days after the acquisition trial (see Figure 1a for Experimental Design).

OLM retention testing at 7 days after the acquisition trial revealed persistent long-term memory with NaB treatment but not exercise (Figure 2c), suggesting that the effects of exercise on spatial memory in the subthreshold paradigm eventually decay after daily physical activity ceases. Interestingly, the pattern of *bdnf* mRNA expression across treatment groups paralleled the cognitive data. Specifically, memory testing 7 days after the acquisition trial revealed that exercised animals had no enduring cognitive enhancement (Figure 2c), and no significant elevation of total *bdnf* mRNA (Figure 2d) or any *bdnf* transcript examined (Supplementary Figure S2 D–F). In contrast, animals treated

with post-acquisition NaB showed persistent long-term memory (Figure 2c), along with elevated levels of total *bdnf* (Figure 2d) at the 7 day timepoint. The elevation of total *bdnf* was associated with increased levels of *bdnf* IV (Supplementary Figure S2E) but not *bdnf* I (Supplementary Figure S2D) or *bdnf* VI (Supplementary Figure S2F). These findings indicate that exercise effects on hippocampal function and *bdnf* gene expression patterns are reversible, whereas effects of HDAC inhibition appear to be relatively stable over time.

Exercise and NaB Hyperacetylate H4K8-Associated *bdnf* I and *bdnf* IV Promoters

As NaB is an HDAC inhibitor, we next investigated whether the increased transcription of *bdnf* and its exons by NaB is associated with specific promoter acetylation. In parallel, we investigated whether exercise increases *bdnf* promoter acetylation, and compared the patterns of *bdnf* promoter acetylation associated with the NaB and exercise treatments. We specifically examined acetylation of H4K8Ac and H4K12Ac of *bdnf* promoters I, IV, and VI, based on literature that acetylation of H4K8 and H4K12 are important in memory consolidation processes (Peleg et al, 2010; McQuown et al, 2011).

A significant effect of treatment was found for H4K8Ac at *bdnf* I ($F_{(3,12)} = 6.98$, $p < 0.05$) and *bdnf* IV ($F_{(3,12)} = 5.624$, $p < 0.05$) but not *bdnf* VI (Figure 3). *Post hoc* tests revealed that exercise and NaB each increased H4K8Ac at *bdnf* I ($p < 0.05$; Figure 3a) and *bdnf* IV ($p < 0.05$; Figure 3b) with no change in *bdnf* VI (Figure 3c), paralleling the respective mRNA patterns for the related transcripts (Supplementary Figure S2 A–C). These results suggest that increased *bdnf* mRNA levels with exercise and NaB are likely regulated through H4K8Ac of BDNF. At the H4K12 acetylation site, there was a significant effect of treatment for *bdnf* I ($F_{(3,12)} = 4.476$, $p < 0.05$), *bdnf* IV ($F_{(3,12)} = 5.593$, $p < 0.05$) and *bdnf* VI promoters ($F_{(3,12)} = 3.83$, $p < 0.05$); however, *post hoc* tests revealed no significant increase in H4K12Ac levels in the exercise- or NaB-treated groups relative to the sedentary vehicle group at any *bdnf* promoter (Figure 3d–f). Interestingly, OLM testing on its own had a small effect on acetylation patterns relative to baseline acetylation levels (eg, sedentary-veh vs baseline), upregulating promoter H4K8Ac for *bdnf* I (Figure 3a) and IV (Figure 3b) as well as H4K12Ac of *bdnf* IV (Figure 3e), suggesting that these histone modifications are highly sensitive to environmental stimulation. Finally, although acetylation of H4K12 has been shown to be key for normal memory consolidation in the aged mouse brain (Peleg et al, 2010), our results suggest that H4K12 acetylation is not involved in benefits of exercise and NaB on OLM learning in young mice.

Taken together, these data demonstrate that exercise, like NaB, affects chromatin modification on the BDNF gene. Exercise and NaB increase acetylation at H4K8 but not H4K12, revealing highly specific and parallel epigenetic effects of exercise and NaB on *bdnf* promoters. Importantly, these results indicate that exercise may permit greater access of transcriptional machinery by establishing a more accessible chromatin state, similar to the mechanisms of NaB.

Figure 3 Exercise (Ex) and sodium butyrate (NaB) increase acetylation of histone H4, specifically acetylation at H4K8 of promoters for (a) brain-derived neurotrophic factor (*bdnf*) I and (b) *bdnf* IV, but not (c) *bdnf* VI. Acetylation at H4K12 was not significantly increased by Ex or NaB at either promoters (d) *bdnf* I, (e) *bdnf* IV, or (f) *bdnf* VI. Comparing sedentary (Sed) animals with or without object location memory (OLM) testing, behavioral testing had a small effect on H4K8 acetylation on *bdnf* promoters I and IV, and H4K12 acetylation in *bdnf* promoter IV. Data are expressed as means $\pm$ SEM, * p < 0.05 vs Sed + Veh (OLM tested), # p < 0.05, vs baseline acetylation level in sedentary unhandled animals (indicated by dotted line).

Figure 4 Preventing increased brain-derived neurotrophic factor (*bdnf*) mRNA expression with intrahippocampal short-interfering RNA (siRNA) blocks the cognitive benefits of exercise (Ex) and sodium butyrate (NaB) in a subthreshold object location memory (OLM) learning task. (a) Exercise-dependent improvements in the subthreshold OLM paradigm were abolished by intrahippocampal infusion of *bdnf* siRNA, but not by control siRNA. Intrahippocampal *bdnf* siRNA blocked induction of (b) *bdnf* mRNA and (c) BDNF protein expression by exercise and decreased levels of mRNA and protein below baseline. (d) Cognitive improvements in the subthreshold OLM paradigm with NaB treatment were blocked by intrahippocampal infusion of *bdnf* siRNA, but not control siRNA. Data are expressed as means $\pm$ SEM. * p < 0.05; ** p < 0.01; *** p < 0.001.

BDNF Induction is Required for Cognitive Enhancement by Exercise and NaB

Finally, we investigated whether induction of hippocampal BDNF transcription is necessary for exercise and NaB to transform a subthreshold learning event into robust long-term memory. siRNA targeting BDNF was infused bilaterally into the hippocampus 48 h before the subthreshold OLM task to selectively block elevation of hippocampal *bdnf* mRNA by exercise or NaB, and memory was assessed 24 h after the acquisition trial.

There was a significant effect of treatment for exercise and siRNA on OLM ($F_{(2,29)} = 4.80$, $p < 0.01$). Consistent with our previous findings, exercised animals treated with control siRNA showed significant discrimination and learning at the 24-h memory test after subthreshold training (Figure 4a). In contrast, BDNF siRNA prevented the cognitive enhancement by exercise, such that the exercise/BDNF siRNA group showed no significant preference for

the object in the novel location, similar to sedentary controls (Figure 4a). In parallel with the effect on cognitive performance, BDNF siRNA prevented the induction of *bdnf* gene expression (Figure 4b) and BDNF protein (Figure 4c) in response to exercise, while in the presence of control siRNA, exercise increased *bdnf* gene expression 1.6-fold over sedentary/control siRNA levels (Figure 4b). These data demonstrate that induction of *bdnf* mRNA is required for exercise to transform a normally subthreshold learning event into long-term memory.

Similarly, our data reveal that induction of *bdnf* mRNA is necessary for the cognitive facilitation by NaB. Specifically, whereas NaB-treated mice infused with control siRNA exhibited significant preference for the object in the novel location, NaB-treated mice infused with BDNF siRNA exhibited no preference (Figure 4d), demonstrating that BDNF is necessary for HDAC inhibition via NaB to transform a subthreshold learning event into long-term memory. Interestingly, BDNF siRNA reduced levels of hippocampal

bdnf mRNA and protein below baseline expression to 61% of sedentary controls (Figure 4b and c), predominantly because of reductions below baseline of *bdnf* transcript IV (Supplementary Figure S3B) and *bdnf* VI (Supplementary Figure S3C), whereas no effect on *bdnf* I was observed (Supplementary Figure S3A). The drop below baseline in the presence of BDNF siRNA is consistent with the fact that *bdnf* mRNA undergoes turnover in the hippocampus, even in sedentary animals.

Taken together, these data demonstrate that induction of *bdnf* mRNA with exercise and NaB is necessary to transform a subthreshold learning event into robust long-term memory, such that exercise and NaB facilitate encoding in a hippocampal *bdnf*-dependent manner.

DISCUSSION

In this study, we report the novel finding that exercise enables the acquisition of subthreshold experiences, such that a hippocampal-dependent learning event that would not normally be remembered is encoded into long-term memory. Remarkably, exercise-induced improvements in the subthreshold learning paradigm were similar to those attained by treatment with NaB (Stefanko *et al*, 2009), a memory-enhancing drug. Our data demonstrate that the enabling effect of exercise and NaB on subthreshold OLM learning is dependent on BDNF upregulation, and is prevented by hippocampal infusion of BDNF siRNA.

The finding that exercise, like NaB, enables the acquisition of subthreshold experiences adds to growing evidence that exercise alters the sensitivity of hippocampal neural networks and enhances learning (Cotman and Berchtold, 2007). In addition, these data provide a behavioral readout consistent with electrophysiology data that a weak LTP induction paradigm that is normally subthreshold becomes a sufficient induction stimulus for LTP after exercise (Farmer *et al*, 2004) or exposure to NaB (Vecsey *et al*, 2007). Finally, the demonstration that OLM is facilitated following exercise reveals that voluntary physical activity improves a non-reinforced form of hippocampal-dependent learning, complementing previous findings that exercise improves learning in negatively motivated hippocampal-dependent tasks (such as the Morris water maze, which relies on an intention to escape a swimming stressor; van Praag *et al*, 1999; Vaynman *et al*, 2004; Berchtold *et al*, 2010; Cassilhas *et al*, 2012). OLM, being a non-reinforced form of learning, may more closely approximate the type of learning most prevalent in everyday life for humans, where learning accompanies daily activities in the absence of explicit instruction or positive/negative reinforcement.

Our finding that the enabling effect of exercise and NaB on encoding is blocked in the presence of BDNF siRNA is consistent with several lines of evidence demonstrating a key role for BDNF in various hippocampus-dependent spatial memory tasks (Ma *et al*, 1998; Mu *et al*, 1999; Mizuno *et al*, 2000; Gorski *et al*, 2003; Vaynman *et al*, 2004). The data underscore the requirement for hippocampal BDNF signaling in spatial memory, and support the concept that physical activity elevates BDNF to levels that enhance hippocampal function. In addition, the finding that exercise increases hippocampal levels of *bdnf* I and IV (but not VI)

builds on previous evidence that physical activity selectively increases expression of these transcripts (Oliff *et al*, 1998; Russo-Neustadt *et al*, 2000). Finally, our data reveal that NaB targets the same *bdnf* transcripts as exercise, upregulating *bdnf* I and IV, but not *bdnf* VI. These data suggest that hippocampal expression of *bdnf* I and IV may be triggered by common signaling pathways as part of the adaptive response to exercise or NaB, and serve to facilitate learning.

In parallel with assessing performance benefits of exercise and NaB at 24 h after OLM training, a timepoint that informs whether learning was encoded and transferred into long-term memory, we extended our analysis to 7 days after the acquisition trial to assess the stability of the memory enhancement. The enhancing effects of NaB on memory endured at the 7-day timepoint, consistent with previous findings (Stefanko *et al*, 2009; McQuown *et al*, 2011), and the effects on spatial memory were paralleled by sustained increases in total *bdnf* and *bdnf* transcripts I and IV. In contrast, the enhancement of OLM by exercise was not maintained after 7 days, and the loss of cognitive enhancement was paralleled by return of *bdnf* mRNA to baseline levels. These findings indicate that the effects of exercise on hippocampal function and *bdnf* gene expression patterns are reversible, while effects of HDAC inhibition appear to be relatively stable over time. It is important to note that a critical aspect of plasticity is cognitive flexibility, to allow for the rapid adaptation to changing rules and for previous associations to be updated.

Finally, the data reveal that exercise and NaB establish a permissive chromatin state that allows signaling stimulated by subthreshold learning to induce BDNF expression, resulting in robust encoding. The permissive chromatin states induced by exercise and NaB have epigenetic signatures associated with increased H4K8Ac of promoters specific for *bdnf* I and *bdnf* IV (but not *bdnf* VI), and the respective mRNA transcripts are increased in parallel with the acetylation patterns. Increased H4K8 acetylation enhances hippocampal gene expression required to transform a subthreshold OLM training event into long-term memory (previously shown for NaB treatment; McQuown *et al*, 2011), indicating that H4K8 acetylation may also be an important epigenetic mechanism by which exercise enhances hippocampal function. One mechanism underlying H4K8 acetylation with exercise likely involves enhanced CREB activity (Shen *et al*, 2001), as exercise increases CREB, and CREB activity in turn increases H4K8 acetylation (Wang *et al*, 2010). Importantly, activated CREB binds *bdnf* I and IV promoters (Ou and Gean, 2007) and recruits coactivators with HAT activity required for spatial learning (Alarcon *et al*, 2004; Korzus *et al*, 2004; Wood *et al*, 2005). Another mechanism underlying H4K8 acetylation with exercise may be via decreased HDAC expression by exercise, as has recently been reported (Gomez-Pinilla *et al*, 2011; Abel and Rissman, 2012). Interestingly, while acetylation at H4K12 has been shown to be key for memory consolidation in the aged mouse brain (Peleg *et al*, 2010), neither exercise nor NaB significantly increased acetylation of H4K12 on *bdnf* promoters, suggesting that H4K12 acetylation is not involved in benefits of exercise and NaB on subthreshold OLM learning in young adult animals. Although we have focused on H4 acetylation of BDNF in

this study, a potential role of H3 acetylation may also be involved in BDNF induction and the enabling effects of exercise and NaB on OLM memory, as several learning and memory studies have implicated H3K14 acetylation in the consolidation mechanisms involving gene regulation (Levenson *et al*, 2004; Chwang *et al*, 2006; Barrett *et al*, 2011; Stafford *et al*, 2012). Ultimately, deciphering the roles of specific acetylation sites is critical for identifying novel therapeutic approaches for cognitive dysfunction, especially considering that histone hypoacetylation is a feature of several neurodegenerative diseases where learning and memory are compromised (Rouaux *et al*, 2003; Urdinguio *et al*, 2009).

In summary, these data provide strong support for the concept that exercise engages epigenetic control mechanisms and serves as a natural stimulus that operates in part like NaB and potentially other HDAC inhibitors. The permissive chromatin state induced by exercise enhances the effect of signaling events, driving gene expression required for long-term memory formation, and enabling learning to occur in conditions that are normally subthreshold for encoding into long-term memory. These findings add to accruing evidence that physical activity is a potent strategy to enhance hippocampal function and cognitive performance.

ACKNOWLEDGEMENTS

Funding was provided by the following: NIA Grants AG00538 and AG34667 (CWC), T32AG9629 (KAI), NIMH Grant MH081004 (MAW), and NRSA predoctoral fellowship F31DA29368 (MM). We thank Arpine Kirakos for technical assistance with ChIP assays and Dina Matheos for helpful RT-qPCR advice.

DISCLOSURE

The authors declare no conflict of interest.

REFERENCES

- Abel JL, Rissman EF (2012). Running-induced epigenetic and gene expression changes in the adolescent brain. *Int J Dev Neurosci*: e-pub ahead of print. doi:pii:S0736-5748(12)00585-0.
- Ahlskog JE, Geda YE, Graff-Radford NR, Petersen RC (2011). Physical exercise as a preventive or disease-modifying treatment of dementia and brain aging. *Mayo Clin Proc* 86: 876–884.
- Alarcon JM, Malleret G, Touzani K, Vronskaya S, Ishii S, Kandel ER *et al* (2004). Chromatin acetylation, memory, and LTP are impaired in CBP +/− mice: a model for the cognitive deficit in Rubinstein-Taybi syndrome and its amelioration. *Neuron* 42: 947–959.
- Barrett RM, Malvaez M, Kramar E, Matheos DP, Arrizon A, Cabrera SM *et al* (2011). Hippocampal focal knockout of CBP affects specific histone modifications, long-term potentiation, and long-term memory. *Neuropsychopharmacology* 36: 1545–1556.
- Bekinschtein P, Oomen CA, Saksida LM, Bussey TJ (2011). Effects of environmental enrichment and voluntary exercise on neurogenesis, learning and memory, and pattern separation: BDNF as a critical variable? *Semin Cell Dev Biol* 22: 536–542.
- Berchtold NC, Castello N, Cotman CW (2010). Exercise and time-dependent benefits to learning and memory. *Neuroscience* 167: 588–597.
- Bredy TW, Wu H, Crego C, Zellhoefer J, Sun YE, Barad M (2007). Histone modifications around individual BDNF gene promoters in prefrontal cortex are associated with extinction of conditioned fear. *Learn Mem* 14: 268–276.
- Calfa G, Chapleau CA, Campbell S, Inoue T, Morse SJ, Lubin FD *et al* (2012). HDAC activity is required for BDNF to increase quantal neurotransmitter release and dendritic spine density in CA1 pyramidal neurons. *Hippocampus* 22: 1493–1500.
- Cassilhas RC, Lee KS, Fernandes J, Oliveira MG, Tufik S, Meeusen R *et al* (2012). Spatial memory is improved by aerobic and resistance exercise through divergent molecular mechanisms. *Neuroscience* 202: 309–317.
- Chwang WB, O’Riordan KJ, Levenson JM, Sweatt JD (2006). ERK/MAPK regulates hippocampal histone phosphorylation following contextual fear conditioning. *Learn Mem* 13: 322–328.
- Cotman CW, Berchtold NC (2002). Exercise: a behavioral intervention to enhance brain health and plasticity. *Trends Neurosci* 25: 292–298.
- Cotman CW, Berchtold NC (2007). Physical activity and the maintenance of cognition: learning from animal models. *Alzheimers Dement* 3: S30–S37.
- Cotman CW, Berchtold NC, Christie LA (2007). Exercise builds brain health: an interplay of central and peripheral factors. *Trends Neurosci* 30: 467–471.
- Farmer J, Zhao X, van Praag H, Wodtke K, Gage FH, Christie BR (2004). Effects of voluntary exercise on synaptic plasticity and gene expression in the dentate gyrus of adult male Sprague-Dawley rats in vivo. *Neuroscience* 124: 71–79.
- Gomez-Pinilla F, Zhuang Y, Feng J, Ying Z, Fan G (2011). Exercise impacts brain-derived neurotrophic factor plasticity by engaging mechanisms of epigenetic regulation. *Eur J Neurosci* 33: 383–390.
- Gorski JA, Balogh SA, Wehner JM, Jones KR (2003). Learning deficits in forebrain-restricted brain-derived neurotrophic factor mutant mice. *Neuroscience* 121: 341–354.
- Gottmann K, Mittmann T, Lessmann V (2009). BDNF signaling in the formation, maturation and plasticity of glutamatergic and GABAergic synapses. *Exp Brain Res* 199: 203–234.
- Korzus E, Rosenfeld MG, Mayford M (2004). CBP histone acetyltransferase activity is a critical component of memory consolidation. *Neuron* 42: 961–972.
- Kouzarides T (2007). Chromatin modifications and their function. *Cell* 128: 693–705.
- Levenson JM, O’Riordan KJ, Brown KD, Trinh MA, Molfese DL, Sweatt JD (2004). Regulation of histone acetylation during memory formation in the hippocampus. *J Biol Chem* 279: 40545–40559.
- Lu Y, Christian K, Lu B (2008). BDNF: a key regulator for protein synthesis-dependent LTP and long-term memory? *Neurobiol Learn Mem* 89: 312–323.
- Ma YL, Wang HL, Wu HC, Wei CL, Lee EH (1998). Brain-derived neurotrophic factor antisense oligonucleotide impairs memory retention and inhibits long-term potentiation in rats. *Neuroscience* 82: 957–967.
- McQuown SC, Barrett RM, Matheos DP, Post RJ, Rogge GA, Alenghat T *et al* (2011). HDAC3 is a critical negative regulator of long-term memory formation. *J Neurosci* 31: 764–774.
- Middleton LE, Barnes DE, Lui LY, Yaffe K (2010). Physical activity over the life course and its association with cognitive performance and impairment in old age. *J Am Geriatr Soc* 58: 1322–1326.
- Mizuno M, Yamada K, Olariu A, Nawa H, Nabeshima T (2000). Involvement of brain-derived neurotrophic factor in spatial memory formation and maintenance in a radial arm maze test in rats. *J Neurosci* 20: 7116–7121.
- Mu JS, Li WP, Yao ZB, Zhou XF (1999). Deprivation of endogenous brain-derived neurotrophic factor results in impairment of spatial learning and memory in adult rats. *Brain research* 835: 259–265.

- Neeper SA, Gomez PF, Choi J, Cotman CW (1996). Physical activity increases mRNA for brain-derived neurotrophic factor and nerve growth factor in rat brain. *Brain Res* 726: 49–56.
- Oliff HS, Berchtold NC, Isackson P, Cotman CW (1998). Exercise-induced regulation of brain-derived neurotrophic factor (BDNF) transcripts in the rat hippocampus. *Brain Res Mol Brain Res* 61: 147–153.
- Ou LC, Gean PW (2007). Transcriptional regulation of brain-derived neurotrophic factor in the amygdala during consolidation of fear memory. *Mol Pharmacol* 72: 350–358.
- Peleg S, Sananbenesi F, Zovoilis A, Burkhardt S, Bahari-Javan S, Agis-Balboa RC *et al* (2010). Altered histone acetylation is associated with age-dependent memory impairment in mice. *Science* 328: 753–756.
- Rex CS, Lin CY, Kramar EA, Chen LY, Gall CM, Lynch G (2007). Brain-derived neurotrophic factor promotes long-term potentiation-related cytoskeletal changes in adult hippocampus. *J Neurosci* 27: 3017–3029.
- Rouaux C, Jokic N, Mbebi C, Boutillier S, Loeffler JP, Boutillier AL (2003). Critical loss of CBP/p300 histone acetylase activity by caspase-6 during neurodegeneration. *EMBO J* 22: 6537–6549.
- Russo-Neustadt AA, Beard RC, Huang YM, Cotman CW (2000). Physical activity and antidepressant treatment potentiate the expression of specific brain-derived neurotrophic factor transcripts in the rat hippocampus. *Neuroscience* 101: 305–312.
- Sahar S, Reddy MA, Wong C, Meng L, Wang M, Natarajan R (2007). Cooperation of SRC-1 and p300 with NF-kappaB and CREB in angiotensin II-induced IL-6 expression in vascular smooth muscle cells. *Arterioscler Thromb Vasc Biol* 27: 1528–1534.
- Shen H, Tong L, Balazs R, Cotman CW (2001). Physical activity elicits sustained activation of the cyclic AMP response element-binding protein and mitogen-activated protein kinase in the rat hippocampus. *Neuroscience* 107: 219–229.
- Stafford JM, Raybuck JD, Ryabinin AE, Lattal KM (2012). Increasing histone acetylation in the hippocampus-infralimbic network enhances fear extinction. *Biol Psychiatry* 72: 25–33.
- Stefanko DP, Barrett RM, Ly AR, Reolon GK, Wood MA (2009). Modulation of long-term memory for object recognition via HDAC inhibition. *Proc Natl Acad Sci USA* 106: 9447–9452.
- Urdinguio RG, Sanchez-Mut JV, Esteller M (2009). Epigenetic mechanisms in neurological diseases: genes, syndromes, and therapies. *Lancet Neurol* 8: 1056–1072.
- van Praag H, Christie BR, Sejnowski TJ, Gage FH (1999). Running enhances neurogenesis, learning, and long-term potentiation in mice. *Proc Natl Acad Sci USA* 96: 13427–13431.
- Vaynman S, Ying Z, Gomez-Pinilla F (2004). Hippocampal BDNF mediates the efficacy of exercise on synaptic plasticity and cognition. *Eur J Neurosci* 20: 2580–2590.
- Vecsey CG, Hawk JD, Lattal KM, Stein JM, Fabian SA, Attner MA *et al* (2007). Histone deacetylase inhibitors enhance memory and synaptic plasticity via CREB:CBP-dependent transcriptional activation. *J Neurosci* 27: 6128–6140.
- Voss MW, Nagamatsu LS, Liu-Ambrose T, Kramer AF (2011). Exercise, brain, and cognition across the lifespan. *J Appl Physiol* 111: 1505–1513.
- Wang D, Xia X, Weiss RE, Refetoff S, Yen PM (2010). Distinct and histone-specific modifications mediate positive versus negative transcriptional regulation of TSHalpha promoter. *PLoS ONE* 5: e9853.
- Wood MA, Kaplan MP, Park A, Blanchard EJ, Oliveira AM, Lombardi TL *et al* (2005). Transgenic mice expressing a truncated form of CREB-binding protein (CBP) exhibit deficits in hippocampal synaptic plasticity and memory storage. *Learn Mem* 12: 111–119.
- Zeng Y, Tan M, Kohyama J, Sneddon M, Watson JB, Sun YE *et al* (2011). Epigenetic enhancement of BDNF signaling rescues synaptic plasticity in aging. *J Neurosci* 31: 17800–17810.

Supplementary Information accompanies the paper on the Neuropsychopharmacology website (<http://www.nature.com/npp>)