Memory Allocation

Memories are thought to be represented in the brain as enduring physical changes in ensembles of neurons, the 'engram'. How are neurons chosen (or allocated) to become part of the engram? Is this process random? Evidence suggests it is not. We (Han *et al*, 2009) and others (Zhou *et al*, 2009) examined how fear/threat memories are encoded in the brain. In this paradigm, rodents learn to associate a tone with a shock and this association critically depends on the lateral amygdala (LA) (Repa *et al*, 2001). Manipulating levels of the transcription factor CREB in LA neurons influenced the likelihood of a neuron becoming part of a fear/threat memory engram. To determine 'engram membership' we quantified expression of the activity-regulated gene *arc* after encoding and/or retrieval. When CREB was virally overexpressed in random LA neurons, those infected neurons were more likely to be allocated to the engram than their non-infected neighbors. Conversely, when CREB function was virally suppressed, infected neurons were excluded from the engram (Han *et al*, 2009). These findings suggested that neurons with relatively higher CREB are more likely to 'capture' the memory. Definitive evidence for this emerged from subsequent experiments: genetic ablation or suppression of the CREB overexpressing cells (and not an equivalently sized random population of LA neurons) was sufficient to erase the memory (Han *et al*, 2009; Zhou *et al*, 2009).

CREB is involved in many biological processes. Which process is responsible for the increased likelihood of allocation? One candidate is neuronal excitability. Overexpression of CREB increases a neuron's intrinsic excitability (Dong *et al*, 2006; Zhou *et al*, 2009). Might changes in neuronal excitability allow LA neurons with high CREB function to win the neuronal competition and become allocated to the engram? To test this, we manipulated excitability in LA neurons by targeting K^+ channels

and using genetic mediators of excitability (DREADDs, optogenetics). Remarkably, we found that increasing neuronal excitability via different methods mimicked the effects of CREB overexpression: Fear/threat memories were funnelled into these more excitable neurons. Conversely, blocking CREB-induced increases in neuronal excitability (by co-expressing Kir2.1, an inwardly rectifying K^+ channel, which reduces neuronal excitability) prevented their preferential allocation (Yiu *et al*, 2014). Our finding that neurons are recruited to an engram based on neuronal excitability was predicted by a recent biophysical modeling study (Kim *et al*, 2013).

Why allocate? Recalling a particular event might conjure up memories of closely related episodes. This phenomenon may reflect some underlying structure of the way in which our memories are organized, with memories that are related either in content or in time encoded by overlapping ensembles of neurons. Within this associative network, fluctuations in CREB/excitability determine whether memories are linked or, alternately, segregated. Is it possible to alter the structure or function of this fundamental associative network by hijacking the allocation process? For example, manipulating CREB levels in different neuronal ensembles might artificially link otherwise unrelated memories or, conversely, uncouple memories that would normally be allocated to overlapping populations of neurons. Understanding the rules of allocation might provide insights into a range of psychiatric conditions that are characterized by inappropriate associations such as schizophrenia. We wonder, therefore, to what extent different psychopathologies can be thought of as disorders of mis-allocation.

ACKNOWLEDGEMENTS

This work was supported by grants from the Canadian Institutes of Health Research (CIHR; MOP-74650), Brain and Behavior Foundation (NARSAD), Natural Sciences and

Engineering Research Council of Canada (NSERC), EJLB Foundation and the Alzheimer's Society of Canada to SAJ and CIHR (MOP-86762), and NSERC to PWF.

Paul W Frankland^{1,2,3,4} and Sheena A Josselyn^{1,2,3,4}

¹Program in Neurosciences & Mental Health, Hospital for Sick Children, Toronto, ON, Canada; ²Department of Psychology, University of Toronto, Toronto, ON, Canada; ³Institute of Medical Sciences, University of Toronto, Toronto, ON, Canada; ⁴Department of Physiology, University of Toronto, Toronto, ON, Canada

E-mail: sheena.josselyn@sickkids.ca

FUNDING AND DISCLOSURE

The authors declare no conflict of interest.

Dong Y, Green T, Saal D, Marie H, Neve R, Nestler EJ *et al*. (2006). CREB modulates excitability of nucleus accumbens neurons. *Nat Neurosci* **9**: 475–477.

Han JH, Kushner SA, Yiu AP, Hsiang HL, Buch T, Waisman A *et al*. (2009). Selective erasure of a fear memory. *Science* **323**: 1492–1496.

Kim D, Pare D, Nair SS (2013). Assignment of model amygdala neurons to the fear memory trace depends on competitive synaptic interactions. *J Neurosci* **33**: 14354–14358.

Repa JC, Muller J, Apergis J, Desrochers TM, Zhou Y, LeDoux JE (2001). Two different lateral amygdala cell populations contribute to the initiation and storage of memory. *Nat Neurosci* **4**: 724–731.

Yiu AP, Mercaldo V, Yan C, Richards B, Rashid AJ, Hsiang HL *et al*. (2014). Neurons are recruited to a memory trace based on relative neuronal excitability immediately before training. *Neuron* **83**: 722–735.

Zhou Y, Won J, Karlsson MG, Zhou M, Rogerson T, Balaji J *et al*. (2009). CREB regulates excitability and the allocation of memory to subsets of neurons in the amygdala. *Nature Neurosci* **12**: 1438–1443.

Neuropsychopharmacology Reviews (2015) **40**, 243; doi:10.1038/npp.2014.234

Oxytocin, Social Cognition and Psychiatry

Oxytocin (OT) has an ancient role in modulating sensing and responding to social stimuli, from nematodes to man. OT regulates not only mammalian labor and nursing, but also maternal behavior. There is increasing evidence that OT influences human parenting and mediates the impact of parenting

on infant socio-emotional development (Rilling and Young, 2014). Preclinical studies suggest that OT increases the salience and rewarding value of social stimuli. OT acts in the rodent amygdala to enhance the salience of social olfactory cues, thereby facilitating social recognition, in the striatum to mediate social reward and in the hippocampus to enhance signal to noise neurotransmission. These fundamental processes likely contribute to more complex OT-mediated behaviors, including social bonding.

The effects of OT on social information processing in rodents make it an enticing pharmacological target for enhancing social cognition. However, two issues introduce skepticism for translating the compelling preclinical observations into effective pharmacotherapies to improve social functioning in psychiatric disorders, including autism and schizophrenia: (1) rodents use olfaction as the primary social perception modality, while primates rely more on visual and auditory social perception; (2) little is known regarding the pharmacokinetics of current OT administration methods or the impact of chronic OT treatment. Recent studies from our laboratory address these issues.

A common polymorphism in the human OT receptor (OXTR) gene predicts face recognition skills in families with a child with autism. This effect was present in all family members in two independent populations, yet there was no evidence of an association with autism diagnosis (Skuse *et al*, 2014). This study supports a role for the OT system in human visual social information processing analogous to its role in olfactory processing in rodents.

Nonhuman primates are useful for exploring the mechanisms of intranasal OT (IN-OT) administration. We showed that OT administered nasally by a pediatric nebulizer modestly elevates OT in the cerebrospinal fluid of anesthetized macaques (Modi *et al*, 2014). Importantly, intranasal OT also robustly elevated plasma OT for an extended period of time. Thus IN-OT may increase brain OT signaling, but

peripheral mechanisms should be considered.

Comparative studies of brain OXTR distribution in primates reveal the potential mechanisms by which OT modulates social information processing (Freeman *et al*, 2014a, b). In all primate species examined, OXTRs are concentrated in cholinergic regions involved in visual and auditory processing, including the nucleus basalis of Meynert, which coordinates neural activity in the amygdala and cortex, thereby modulating attention to visual cues.

IN-OT may enhance some aspects of social cognition through the mechanisms described above, but the efficacy may be limited by brain penetration. Stimulating endogenous central OT release pharmacologically is a viable alternative for increasing OT neural signaling. Melanocortin receptor agonists stimulate OT release from hypothalamic slices, potentiate OT release in the ventral striatum, and enhance OT-dependent behavior in prairie voles (unpublished data). Neonatal melanocortin receptor activation acutely activates OT neurons, and daily treatment for the first week of life enhances adult social bonding in prairie voles (Barrett *et al*, 2014). Thus, the OT system remains an attractive target for clinically enhancing social cognition, and alternative pharmacological strategies for enhancing OT neurotransmission should be explored.

ACKNOWLEDGEMENTS

I would like to acknowledge NIH grants R01MH096983 and 1P50MH100023 and an Autism Speaks grant #7745. Additional funding was provided by NIH OD P51OD11132 to YNPRC.

Larry J Young¹

¹Silvio O. Conte Center for Oxytocin and Social Cognition, Center for Translational Social Neuroscience, Department of Psychiatry and Behavioral Sciences, Yerkes National Primate Research Center, Emory University, Atlanta, GA, USA
E-mail: lyou03@emory.edu

FUNDING AND DISCLOSURE

LJY has applied for a patent (US20120108510—Methods of improving behavioral therapies) for combining melanocortin agonists with behavioral therapies to enhance social cognition in psychiatric disorders.

Barrett CE, Modi ME, Zhang BC, Walum H, Inoue K, Young LJ (2014). Neonatal melanocortin receptor agonist treatment reduces play fighting and promotes adult attachment in prairie voles in a sex-dependent manner. *Neuropharmacology* **85C**: 357–366.

Freeman SM, Inoue K, Smith AL, Goodman MM, Young LJ (2014a). The neuroanatomical distribution of oxytocin receptor binding and mRNA in the male rhesus macaque (*Macaca mulatta*). *Psychoneuroendocrinology* **45**: 128–141.

Freeman SM, Walum H, Inoue K, Smith AL, Goodman MM, Bales KL *et al*. (2014b). Neuroanatomical distribution of oxytocin and vasopressin 1a receptors in the socially monogamous coppery titi monkey (*Callicebus cupreus*). *Neuroscience* **273**: 12–23.

Modi ME, Connor-Stroud F, Landgraf R, Young LJ, Parr LA (2014). Aerosolized oxytocin increases cerebrospinal fluid oxytocin in rhesus macaques. *Psychoneuroendocrinology* **45**: 49–57.

Rilling JK, Young LJ (2014). The biology of mammalian parenting and its impact on offspring development. *Science* **345**: 771–776.

Skuse DH, Lori A, Cubells JF, Lee I, Conneely KN, Puura K *et al*. (2014). Common polymorphism in the oxytocin receptor gene (OXTR) is associated with human social recognition skills. *Proceedings of the national academy of sciences of the U* **111**: 1987–1992.

Neuropsychopharmacology Reviews (2015) **40**, 243–244; doi:10.1038/npp.2014.186

FKBP5 Allele-Specific Epigenetic Modification in Gene by Environment Interaction

The likelihood to develop stress-related psychiatric disorders in response to childhood trauma exposure may be moderated by the individual's genetic predisposition (Manuck and McCaffery, 2014). One of the genetic variants reported to alter the risk for psychiatric disorders following childhood trauma is a functional variant in *FKBP5*, a gene encoding a co-chaperone of the glucocorticoid receptor (GR). *FKBP5* is strongly induced following stress exposure via binding of activated GR to a number of intronic and promoter GR response elements (GREs). The protein itself then binds to the GR complex, reduces the affinity of GR to cortisol and decreases translocation of the GR to the nucleus, providing an ultrashort negative feedback for GR activation on the genomic and protein level (Zannas and Binder, 2014). We