

The Placenta as a Mediator of Stress Effects on Neurodevelopmental Reprogramming

Stefanie L Bronson¹ and Tracy L Bale^{*1}

¹Department of Biomedical Sciences, University of Pennsylvania School of Veterinary Medicine, Philadelphia, PA, USA

Adversity experienced during gestation is a predictor of lifetime neuropsychiatric disease susceptibility. Specifically, maternal stress during pregnancy predisposes offspring to sex-biased neurodevelopmental disorders, including schizophrenia, attention deficit/hyperactivity disorder, and autism spectrum disorders. Animal models have demonstrated disease-relevant endophenotypes in prenatally stressed offspring and have provided unique insight into potential programmatic mechanisms. The placenta has a critical role in the deleterious and sex-specific effects of maternal stress and other fetal exposures on the developing brain. Stress-induced perturbations of the maternal milieu are conveyed to the embryo via the placenta, the maternal–fetal intermediary responsible for maintaining intrauterine homeostasis. Disruption of vital placental functions can have a significant impact on fetal development, including the brain, outcomes that are largely sex-specific. Here we review the novel involvement of the placenta in the transmission of the maternal adverse environment and effects on the developing brain. *Neuropsychopharmacology Reviews* (2016) **41**, 207–218; doi:10.1038/npp.2015.231; published online 26 August 2015

INTRODUCTION

Prenatal adversity is a risk factor for lifetime neuropsychiatric disease susceptibility. During gestation, rapid growth and plasticity render the brain sensitive to the effects of environmental factors that can confer adaptive advantages or lasting vulnerability. As the fetal origins of disease hypothesis was pioneered four decades ago by Barker (1997) and Dörner (1973), substantial advances have been achieved in understanding the neurodevelopmental consequences of intrauterine challenges, including maternal psychosocial stress, infection, and metabolic dysfunction (Bock *et al*, 2014, 2015; Brown and Derkits, 2010; Markham and Koenig, 2011; Weinstock, 2008). Maternal stress during pregnancy, in particular, has been identified in autism spectrum disorders (ASDs), attention deficit/hyperactivity disorder (ADHD), and schizophrenia risk (Beverdorf *et al*, 2005; Khashan *et al*, 2008; Kinney *et al*, 2008a; Li *et al*, 2010; Ronald *et al*, 2010). These neurodevelopmental disorders exhibit a strong sex bias, where boys are more likely to develop ASD, ADHD, and earlier-onset schizophrenia (Baio, 2012; Bloom *et al*, 2011; Froehlich *et al*, 2007; Zhang *et al*, 2012). Consistent with the male predominance of these disorders, recent epidemiological and preclinical studies

suggest that males are uniquely vulnerable to prenatal adversity (Clifton, 2010; Davis and Pfaff, 2014). Although studies have revealed sex-specific changes in the developing brain (Bale, 2011; Kapoor and Matthews, 2005, 2008; Kapoor *et al*, 2009; Weinstock, 2011), the specific mechanisms by which perturbations in the maternal environment promote sex-biased fetal reprogramming remain unclear.

The programmatic effects of prenatal stress likely involve a complex interaction between the fetal genetic background, sex, and gestational age at the time of exposure (Bale, 2011). As the placenta resides at the interface between mother and fetus, it is uniquely positioned to modulate interactions within an adverse intrauterine environment. The placenta actively maintains intrauterine homeostasis through vital functions including exchange of nutrients, oxygen, and waste, immunoprotection of the semi-allogenic fetus, buffering from deleterious maternal factors, and secretion of hormones and growth factors into both maternal and fetal compartments (Jansson and Powell, 2007). Impairment of placental organogenesis and function can broadly have an impact on fetal development, conferring lasting effects on the brain (Myatt, 2006). The placenta comprises specialized cells derived from the embryo and thereby expresses the fetal genetic sex (Rossant and Cross, 2001). In uncomplicated pregnancies, sex differences in placental size and gene expression are present throughout gestation (Buckberry *et al*, 2014; Gabory *et al*, 2013; Mao *et al*, 2010; O'Connell *et al*, 2013; Sood *et al*, 2006). Such basal placental sex differences likely facilitate sex-specific responses to both

*Correspondence: Dr TL Bale, Department of Biomedical Sciences, University of Pennsylvania School of Veterinary Medicine, 380 S University Avenue, Hill 410F, Philadelphia, PA 19104, USA, Tel: +1 215 898 1369, Fax: +1 215 373 5187, E-mail: tbale@vet.upenn.edu
Received 29 April 2015; revised 10 July 2015; accepted 30 July 2015; accepted article preview online 7 August 2015

normal and pathologic environments. Supporting this, sex differences in placental inflammatory responses, vascular remodeling, and placental size and efficiency have been reported in human pregnancies complicated by asthma, preeclampsia, and malnutrition (Clifton, 2010; Gabory *et al*, 2013). Such complications are also associated with altered neurodevelopment (Bale *et al*, 2010; Walker *et al*, 2015). In addition, prospective studies have found abnormal placental histology associated with an autism diagnosis (Anderson *et al*, 2007; Walker *et al*, 2013). Together, these data support the importance of the placenta, and changes in this tissue in response to perturbations across gestation, in the etiology of neurodevelopmental disorders.

In this review we will discuss the programmatic role of the placenta in the transmission of maternal adversity to the developing brain. We will broadly review the effects of maternal stress on offspring neurodevelopment and long-term changes in stress responsivity and behavior, discussing the placenta as a key orchestrator of this sex-specific programming.

LONG-TERM IMPACT OF MATERNAL STRESS

Prenatal Stress and Neuropsychiatric Vulnerability

Epidemiological evidence implicates maternal stress during pregnancy as a factor in poor offspring neuropsychiatric outcome across the lifespan. Children prenatally exposed to maternal stressors such as anxiety, depression, bereavement, and natural or manmade disasters are more likely to present with neurodevelopmental disorders and subclinical psychosocial problems (Beverdors *et al*, 2005; King and Laplante, 2005; Kinney *et al*, 2008a, 2008b; Li *et al*, 2010; Rodriguez and Bohlin, 2005). For example, a higher prevalence of autism was associated with *in utero* exposure to severe tropical storms and ADHD risk was significantly increased following prenatal maternal bereavement (Kinney *et al*, 2008a; Li *et al*, 2010). In addition, children whose mothers experienced high stress levels during the 1998 Quebec Ice Storm exhibited reduced cognitive and language abilities and increased autistic-like traits (Laplante *et al*, 2008; Walder *et al*, 2014). Studies suggest susceptibility is greatest in the first half of gestation, a period of development where males and females appear differentially affected (Gerardin *et al*, 2011; Huttunen and Niskanen, 1978; Khashan *et al*, 2008; van Os and Selten, 1998; Walder *et al*, 2014). Although males present more frequently with childhood behavioral disorders and intellectual impairment associated with prenatal stress, females acquire more subtle later-onset anxiety and affective disorders (Davis and Pfaff, 2014). Such studies predict important sex differences in underlying mechanisms originating early in gestation.

Animal models of prenatal stress have demonstrated disease-relevant endophenotypes that recapitulate aspects of these clinical findings. In nonhuman primates, chronic

maternal stress leads to deficits in attention, object permanence, and motor function in offspring, in particular when the stress was experienced early in gestation (Schneider and Coe, 1993; Schneider, 1992; Schneider *et al*, 1999). In rodent models, prenatal stress induces hypothalamic–pituitary–adrenal (HPA) stress axis dysregulation, heightened behavioral stress reactivity, depression-like behaviors, and cognitive deficits (Bock *et al*, 2015; Brunton and Russell, 2010; Cottrell and Seckl, 2009; Darnaudéry and Maccari, 2008; Kapoor and Matthews, 2005; Kapoor *et al*, 2009; Lemaire *et al*, 2000; Mueller and Bale, 2007, 2008; Weinstock, 2008). Similar to human studies, the gestational timing of the stress and fetal sex are key determinants in offspring outcome, where studies in both mice and guinea pigs found that prenatal stress produced HPA axis dysregulation and cognitive effects only in male offspring exposed during early or mid-gestation (Kapoor and Matthews, 2005, 2008; Kapoor *et al*, 2009; Mueller and Bale, 2007, 2008).

Reprogramming in the Prenatally Stressed Brain

Although the long-term effects of maternal stress in humans on disease outcomes have been studied, earlier assessment of the perinatal brain has been limited to noninvasive measures of growth and macrostructure (Beverdors *et al*, 2005; Khashan *et al*, 2008; Kinney *et al*, 2008b; Li *et al*, 2010; Ronald *et al*, 2010). These studies reported delayed overall brain growth, limbic system-specific volumetric changes, and white matter abnormalities (Li *et al*, 2012; Lou *et al*, 1994; Qiu *et al*, 2013, 2015; Rifkin-Graboi *et al*, 2013). Such early observations appear to persist into childhood and adolescence, are correlated with affective problems, and may contribute to symptoms in patients with schizophrenia and autism (Buss *et al*, 2010, 2012; Davis *et al*, 2013; Du *et al*, 2013; Sarkar *et al*, 2014; Zikopoulos and Barbas, 2010).

Results from animal models of maternal stress support these data and indicate that volumetric abnormalities in prenatally stressed offspring reflect reduced numbers of both neurons and glia, in particular in the hippocampus and amygdala due, in part, to decreased neurogenesis (Coe *et al*, 2003; Fujioka *et al*, 2006; Kawamura *et al*, 2006; Kraszpulski *et al*, 2006; Lemaire *et al*, 2000; Rayen *et al*, 2011). Further, reduced synaptogenesis and hypo-myelination in limbic regions was found in male, but not in female, juvenile rats exposed to maternal stress (Murmu *et al*, 2006; Xu *et al*, 2013). Such changes appear to persist into adulthood, where, eg, decreased hippocampal cell proliferation and reduced dendritic length and complexity were detected in adult prenatally stressed male rats (Mandyam *et al*, 2008; Suenaga *et al*, 2012). These data suggest potential primary impairment of cell proliferation, growth, survival, and connectivity in the prenatally stressed brain, and more specifically in the male brain.




Figure 1. Morphology of the maternal–fetal interface: navigating a complex interaction between maternal and fetal compartments. Mid-sagittal schematic depicting the intricate arrangement of maternal tissue (blue), fetally derived trophoblasts (all five subtypes in purple), and fetal endothelial cells (tan) within the three major functional zones of the mature mouse placenta. The decidua, the most superficial layer from the maternal side (top), comprises maternal uterine and immune cells, as well as specialized glycogen-storing cells from the trophoblast lineage. The decidua is traversed by the maternal spiral arteries, veins, and the fetally derived endovascular trophoblasts that line it. The arrow indicates the direction of maternal blood flow. Bordering the decidua is the junctional zone (basal plate in humans), where maternal vasculature penetrates a layer of trophoblast giant cells and spongiotrophoblasts (extravillous cytotrophoblasts in humans) that secrete hormones to modulate maternal–fetal cross-talk, angiogenesis, and tissue remodeling. Finally, in the labyrinth zone (chorionic villous in humans), cytotrophoblasts and syncytiotrophoblasts residing between maternal blood spaces and fetal endothelial cells (inset) prevent direct blood contact while facilitating selective and essential nutrient/waste exchange. Environmental stimuli such as maternal stress can disrupt vital aspects of placental organogenesis and function, including decidual immune tolerance, vascularization and utero-placental blood flow, trophoblast hormone secretion, and nutrient exchange within the labyrinth zone. The neurodevelopmental consequences of stress depend on the maturational state of the entire maternal–placental–fetal unit at the time of exposure. Although implantation occurs early in gestation (embryonic day 4.5 in mice and the second week in humans), placental maturation and expansion continues throughout gestation, leaving this fetal lifeline and the somatic and germ cells it sustains continuously vulnerable to maternal stress signals.

PLACENTAL ORCHESTRATION OF FETAL BRAIN PROGRAMMING

The Programmatic Capacity of the Placenta

The mechanisms by which maternal stress confers lasting sex-specific neurobehavioral dysfunction likely initiate within the placenta. Stress-mediated perturbations of the maternal milieu must be conveyed to the embryo via interactions with the placenta. This transient organ is not simply a passive thoroughfare, but is an active maternal–fetal intermediary within which distinct functional zones collectively maintain intrauterine homeostasis (depicted in Figure 1). Although human and rodent placentas differ somewhat in their microstructure and developmental trajectory, the gross organization and functions of these zones are conserved (Georgiades *et al*, 2002). The decidua, comprising maternal uterine and immune cells, mediates immunological tolerance of the embryo (Arck and Hecher, 2013). Fetally derived trophoblast cells predominate in the basal plate (junctional zone in rodents), where they synthesize and secrete endocrine factors into both maternal and fetal circulations. Finally, in the chorionic villous (labyrinth zone in rodents), trophoblasts residing between the maternal and fetal vasculature control exchange of

nutrients, oxygen, and waste via facilitated diffusion and macro- and micronutrient transporters (Georgiades *et al*, 2002; Rossant and Cross, 2001; Watson and Cross, 2005). Disruption of these critical functions can have an impact on fetal development, including the brain and primordial germ cells. Importantly, sex-specific reprogramming in response to maternal stress likely arises, owing to sex differences in trophoblasts derived from male (XY) and female (XX) embryos. Sex differences in placental size and gene expression have been identified in normal human and rodent placentas, and sex-specific placental abnormalities predict offspring outcome in pregnancies complicated by maternal asthma and preeclampsia (Buckberry *et al*, 2014; Clifton, 2010; Gabory *et al*, 2013; Howerton *et al*, 2013; Mao *et al*, 2010; O’Connell *et al*, 2013; Sood *et al*, 2006).

Stress at the Dynamic Maternal–Fetal Interface

Placental function is regulated by the collective responses of maternal decidual cells, trophoblastic cells, and fetal endothelial cells to the local environment (Fowden *et al*, 2008). Thus, disruption of the maternal milieu by stress and other stimuli can influence vital aspects of placental structure and function, including integrity of the protective

transplacental barrier, nutrient and oxygen exchange, and placental endocrine action (Jansson and Powell, 2007; Myatt, 2006). The specific consequences likely depend on the timing of exposure, as stress signals interact with dynamic events ongoing in maternal, fetal, and placental compartments. Environmental stimuli may modulate the onset, offset, or duration of these sequential events, resulting in distinct programmatic outcomes.

For example, perturbations in early pregnancy are more likely to produce prolonged effects on placental function via interference with organizational processes including trophoblast differentiation and vascular remodeling (Watson and Cross, 2005). Such changes would then have the potential to impact neurodevelopment across the duration of pregnancy, likely initiating broad programmatic consequences. Epidemiological studies support this unique early gestational vulnerability, as increased schizophrenia and autism risk were specific to first-trimester maternal stress and viral infection exposure, respectively (Atladóttir *et al*, 2010; Khashan *et al*, 2008). In contrast, late-pregnancy perturbations may have a more transient impact on placental actions, by impairing nutrient delivery during periods of high fetal demand or disrupting processes such as neurogenesis, synaptogenesis, and early myelination in the fetal brain. This timing would likely introduce region and cell-type-specific effects on the developing brain that vary depending on species-specific neurodevelopmental trajectories (Andersen, 2003; Avishai-Eliner *et al*, 2002). The capacity for reprogramming, however, is not limited to fetal somatic cells. Beginning in the second week of gestation in mice (3–10 weeks post conception in humans), the newly specified primordial germ cells migrate and undergo a wave of epigenetic remodeling, permitting a unique window during which stress effects may be transmitted to subsequent generations via new epigenetic marks (Bale, 2015; De Felici, 2013; Gapp *et al*, 2014; Saitou and Yamaji, 2012). Supporting this, male mice exposed to early prenatal stress and presenting with disease-relevant endophenotypes, including HPA axis dysfunction and hypothalamic reprogramming, transmitted this phenotype to their offspring in the absence of direct stress re-exposure (Morgan and Bale, 2011).

Human and animal studies have demonstrated timing-dependent and sex-specific effects of maternal stress on placental size, efficiency, and gene expression (summarized in Table 1). Recent studies using transgenic mouse lines to selectively target stress-sensitive placental genes, including O-GlcNAc transferase, were able to recapitulate the effects of prenatal stress on hypothalamic programming and function, and thereby provide strong evidence for the importance of placental function in brain development (Howerton and Bale, 2014; Howerton *et al*, 2013). In the next sections, we describe the effects of maternal stress on aspects of placental structure and function in more detail, and discuss potential mechanisms by which these changes may subsequently reprogram the developing brain.

Transplacental Barrier Permeability

One prominent hypothesis posits that stress compromises the transplacental barrier and, in turn, increases fetal exposure to selectively permeable factors such as steroid hormones and exogenous teratogens (Aye and Keelan, 2013; Seckl and Holmes, 2007). In addition to the structural separation within the labyrinth zone that prevents direct contact between maternal and fetal blood supplies (depicted in Figure 1, inset), metabolizing enzymes localized within trophoblasts facilitate fetal protection from excess glucocorticoids and amines (Brown *et al*, 1996; Nguyen *et al*, 1999). These barrier enzymes can be sensitive to maternal stress, where, eg, reduced placental expression of the glucocorticoid-inactivating enzyme, 11 β -hydroxysteroid dehydrogenase type-2 (11 β HSD2), has been associated with maternal anxiety and depressed mood in humans, as well as with chronic maternal stress in rodents (Blakeley *et al*, 2013; Jensen Peña *et al*, 2012; Mairesse *et al*, 2007; O'Donnell *et al*, 2012; Pankevich *et al*, 2009; Ponder *et al*, 2011). The result is for a potential fetal glucocorticoid overexposure to impact ongoing developmental events including a restriction of fetal growth, premature maturation of proliferative neural precursors, and altered HPA axis development (Seckl and Holmes, 2007). Notably, these predicted effects are consistent with the reduced perinatal brain volumetric findings in association with maternal stress described earlier (Li *et al*, 2012; Lou *et al*, 1994; Qiu *et al*, 2013). In addition to the potential for actions of excess glucocorticoids on the developing fetal brain, excess glucocorticoids also act within the placenta to have an impact on endocrine functions, thus compromising placental growth, vascularization, and nutrient transport (Hewitt *et al*, 2006; Wyrwoll *et al*, 2009). Further, known sex differences in placental 11 β HSD2 expression, both at baseline and in response to environmental stimuli including stress, may contribute to sex-biased neurodevelopmental outcomes (Cuffe *et al*, 2011; Pankevich *et al*, 2009).

In addition to glucocorticoid release, stress increases levels of stress-related neurotransmitters, including serotonin, norepinephrine, and dopamine within both the maternal brain and circulation (Joëls and Baram, 2009). However, less is known about maternal stress effects on placental barrier proteins that may moderate passage of these amines to the fetus. Their fate within the placenta is controlled by their cell-surface transporters and intracellular metabolizing enzymes, such as monoamine oxidase (MAO) and catechol-O-methyl transferase, which regulate delivery to the fetus as well as local processes within the placenta (Nguyen *et al*, 1999). Recent human studies have associated maternal stress with increased levels of serotonin and norepinephrine transporters, and a downregulation of MAO in villous trophoblasts at term (Blakeley *et al*, 2013; Ponder *et al*, 2011). As these cells reside between the maternal and fetal vasculature, such changes in transport would increase their intrauterine availability, especially that of serotonin, which is synthesized in maternal, placental, and fetal compartments

TABLE 1 Summary of Maternal Stress Effects on the Placenta

Species	Stress type	Stress timing	Sex	Placenta phenotype	Reference
Human	Anxiety	36 wpc	?	↑ Umbilical artery resistance, redistribution of fetal blood flow	Sjöström <i>et al</i> , 1997
Human	Anxiety	28–36 wpc	?	↑ Uterine artery resistance	Teixeira <i>et al</i> , 1999
Human	Anxiety	20 wpc	?	No association with uterine artery resistance	Kent <i>et al</i> , 2002
Human	Life stress	30 wpc	♂ and ♀	↑ Placenta weight	Tegethoff <i>et al</i> , 2010
Human	Anxiety and depression	Second to third trimester	♂ and ♀	↑ SLC6A4, trending↑SLC6A2 and 11βHSD2	Ponder <i>et al</i> , 2011
Human	Anxiety	Term	♂ and ♀	↓ 11βHSD2	O'Donnell <i>et al</i> , 2012
Human	Depression	Term	♂ and ♀	Trending↓ 11βHSD2	O'Donnell <i>et al</i> , 2012
Human	Depression	Term	♂ and ♀	↓ MAOA in villous trophoblasts	Blakeley <i>et al</i> , 2013
Human	Anxiety	Term	♂ and ♀	Trending↓MAOA in villous trophoblasts	Blakeley <i>et al</i> , 2013
Human	Depression	Pregnancy	♂ and ♀	↑ Methylation of glucocorticoid receptor	Conradt <i>et al</i> , 2013
Human	Anxiety	Pregnancy	♂ and ♀	↑ Methylation of 11βHSD2	Conradt <i>et al</i> , 2013
Mouse	Chronic variable (1/day)	1–7 dpc	♂ and ♀	↑ IGF2 in ♂ and ♀, ↓ 11βHSD in ♀	Pankevich <i>et al</i> , 2009
Mouse	Chronic variable (1/day)	1–7 dpc	♂ and ♀	♂:↑PPARα, IGFBP1, GLUT4, HIF3α ♀:↓PPARα,↑DNMT1	Mueller and Bale, 2008
Mouse	Chronic variable (1/day)	1–7 dpc	♂ and ♀	↓ OGT in ♂ and ♀	Howerton <i>et al</i> , 2013
Mouse	Chronic variable (1/day)	1–7 dpc	♂ and ♀	♂:↑FASL, pro-inflammatory cytokines (IL6, IL1B, IL12RA, PTGS2), chemokines (CCR7, CCL5, CXCL10), cell surface antigens (H2-EB1, PTPRC), and endothelial molecules (EDN1, SELP). ♀:↑FASL,↓CCL2.	Bronson and Bale, 2014
Rat	Restraint (3/day)	10–21 dpc	♂	↓ GLUT1 and 11βHSD2,↑GLUT3 and GLUT4	Mairesse <i>et al</i> , 2007
Rat	Social defeat (1/day)+restraint (1/day)	5–20 dpc	?	↑ 11βHSD2 in low, but not high anxiety rat strains	Lucassen <i>et al</i> , 2009
Rat	Restraint (1/day)	14–20 dpc	?	↓ 11βHSD2,↑DNMT3a,↑11βHSD2 methylation	Jensen Peña <i>et al</i> , 2012

Abbreviations: 11βHSD, 11β-hydroxysteroid dehydrogenase 2; DNMT, DNA methyltransferase; dpc, days post conception; GLUT, glucose transporter; HIF, hypoxia-inducible factor; IGF, insulin-like growth factor; IGFBP, insulin-like growth factor binding protein; IL, interleukin; MAOA, monoamine oxidase A; OGT, O-GlcNAc transferase; PPAR, peroxisome proliferator-activated receptor; PTGS, prostaglandin; SLC6A4, serotonin transporter; SLC6A2, norepinephrine transporter; wpc, weeks post conception.

In human studies, stress timing refers to the gestational age of maternal stress assessment (wpc). In rodent studies, stress timing indicates the gestational age at maternal stress administration (dpc).

(Bonnin and Levitt, 2011; Nguyen *et al*, 1999; Verhaagh *et al*, 2001). Subsequent fetal overexposure may have deleterious effects on brain development, where, eg, serotonin excess impaired embryonic cortical interneuron migration in mice (Riccio *et al*, 2009; Velasquez *et al*, 2013). Serotonin, a potent vasoconstrictor, also elevates vascular resistance and reduces utero-placental blood flow, a mechanism thought to underlie hypertension in preeclampsia and gestational diabetes (Bolte *et al*, 2001; Li *et al*, 2014). Consistent with these vascular effects, maternal stress has been correlated with Doppler ultrasound indicators of restricted umbilical artery blood flow (Sjöström *et al*, 1997; Teixeira *et al*, 1999). Taken together, maternal stress-induced increased permeability of the transplacental barrier can influence fetal brain development via both direct and indirect actions, and aspects of placental function, such as nutrient transfer and endocrine actions.

Nutrient Exchange and Energy Homeostasis

The fetus relies on placental transfer of nutrients from the maternal circulation, which is achieved via diffusion or active transport across trophoblastic and endothelial cell membranes. The transfer capacity is determined by multiple factors including placenta size, vascularization, metabolism,

and the availability of maternal-facing and fetal-facing transmembrane transporters. These factors are sensitive to the forces of maternal supply and fetal demand, and to adverse intrauterine conditions (Fowden *et al*, 2006; Jansson and Powell, 2006). Clinical studies suggest that placental transport capacity may be perturbed by maternal stress, owing to the utero-placental vascular dysfunction described in the previous section, which can impair bidirectional exchange of flow-limited substrates including oxygen and carbon dioxide (Myatt, 2006). The link between vascular function and nutrient delivery has been well established in uterine artery ligation models of placental insufficiency, where fetal hypoxia and oxidative stress are associated with reduced hippocampal size, abnormal neural migration, and hypomyelination in the offspring (Basiliou *et al*, 2014; Lane *et al*, 2001; Reid *et al*, 2012). In addition, uterine artery ligation in rats disrupted striatal amino acid metabolism at term (Thordstein *et al*, 1992). Thus, fetal hypoxia, oxidative stress, and altered amino acid availability subsequent to stress-induced vascular dysfunction can adversely impact cell survival and differentiation within the fetal brain.

Although not yet investigated in humans, dysregulation of placental nutrient transporters has been demonstrated in animal models of prenatal stress (Mairesse *et al*, 2007; Mueller and Bale, 2008). Transporters are necessary for the

delivery of maternally derived glucose, amino acids, fatty acids, and cholesterol-containing lipoproteins, macronutrients essential for fetal development (Brett *et al*, 2014; Lager and Powell, 2012). Placental expression of the glucose transporter (GLUT) family is particularly sensitive to adverse conditions, such as psychosocial stress, diabetes, and malnutrition in both humans and rodents (Das *et al*, 1998; Illsley, 2000; Mairesse *et al*, 2007; Mueller and Bale, 2008). Further, maternal stress decreased GLUT1 in rat placentas at term (Mairesse *et al*, 2007; Mueller and Bale, 2008). As GLUT1 is the predominant isoform in late pregnancy, these data suggest that stress may reduce glucose transfer to the fetus (Brett *et al*, 2014; Illsley, 2000). In addition, the effects of stress on nutrient exchange are also sex dependent, where stress increased expression of GLUT4 as well as genes regulating fatty acid and oxygen availability in male, but not in female, placentas in rodents (Mairesse *et al*, 2007; Mueller and Bale, 2008). These changes, indicative of disrupted intrauterine energy homeostasis, were also associated with markers of delayed neurodevelopment in male neonates (Mueller and Bale, 2008).

Although changes in nutrient exchange in response to environmental stimuli promote adaptive advantages and maintenance of maternal–fetal homeostasis *in utero*, dysregulated availability of glucose, amino acids, and fatty acids initiates broad programmatic effects on the fetal brain (Innis, 2005; Morgane *et al*, 1993). For example, the fetus relies on fatty acids derived from the maternal circulation (Lager and Powell, 2012). Deficiency in long-chain polyunsaturated fatty acids during gestation, such as docosahexaenoic acid and arachidonic acid, which comprise 40%–50% of neuronal membrane phospholipids in the developing brain, resulted in altered neuronal membrane composition, increased neural inflammation, impaired microglia motility, and programmed long-term impairment in sensorimotor gating in offspring (Fedorova *et al*, 2009; Madore *et al*, 2014; Tam and Innis, 2006). Reduced fatty acid transport can also directly compromise placental function, owing to excess accumulation and subsequent oxidative stress (Jarvie *et al*, 2010). The high fetal demand for cholesterol in late gestation is met by both placental transfer from the maternal circulation and by fetal *de novo* synthesis (Lindegaard *et al*, 2008; Yoshida and Wada, 2005). Therefore, changes in placental cholesterol uptake and transfer are more likely to affect the fetal brain via further impact on placental function. For example, impaired trophoblast uptake of cholesterol may disrupt maternal–fetal cross-talk by reducing substrates for steroidogenesis as well as for synthesis of cholesterol-containing exosomes, non-hormonal communicators that regulate immune homeostasis (Beninson and Fleshner, 2014; Ouyang *et al*, 2014).

Endocrine Action

Bidirectional communication is achieved via hormone secretion by maternal-facing and fetal-facing trophoblasts.

These cells synthesize and secrete growth factors, immunomodulators, sex steroids, metabolic mediators such as leptin and lactogen, and neuromodulators such as corticotropin-releasing factor (CRF) and serotonin (reviewed in Bonnin and Levitt, 2011; Bowen *et al*, 2002; Fowden *et al*, 2014; Reis *et al*, 2001; Sagawa *et al*, 2002; Sandman, 2015). Placental hormones act as endocrine, paracrine, and autocrine modulators of maternal and fetal physiology throughout pregnancy, in particular during implantation, at parturition, and in response to intrauterine conditions including stress signals. Their synthesis is regulated, in part, by gestational age and fetal sex, and some placental hormones are species specific (Carter, 2012; Fowden *et al*, 2014).

Perturbation of placental endocrine actions by stress can have profound effects on fetal neurodevelopment and subsequent disease risk. The role of placental-derived CRF in this reprogramming has been thoroughly investigated (reviewed in Avishai-Eliner *et al*, 2002; Sandman, 2015). Although placental synthesis of CRF in rodents has not yet been demonstrated, the human placenta produces CRF beginning in the seventh gestational week, the levels of which exponentially increase in the maternal circulation as pregnancy progresses (Emanuel *et al*, 1994; Goland *et al*, 1988; Petraglia *et al*, 1987; Robinson *et al*, 1989). Placental CRF is also released into the fetal circulation, where it has influence on nervous system development including regulation of proliferation and survival of neural progenitors (Koutmani *et al*, 2013). The bioavailability of CRF during pregnancy is determined by levels of a circulating binding protein (CRF-BP) (Bowman *et al*, 2001). Stress-related changes in CRF and CRF-BP can lead to neurotoxic effects, in particular in HPA and limbic circuits (Avishai-Eliner *et al*, 2002; Fujioka *et al*, 1999). Supporting this, negative correlations between placental CRF levels and fetal startle responses suggest that CRF overexposure delays maturation in humans (Sandman, 2015).

PLACENTAL INTEGRATION OF STRESS SIGNALS

Signal Detection and Response Coordination

Information about adverse intrauterine status is communicated to the placenta by maternally and fetally derived endocrine factors, which together determine placental adaptive responses and resource re-allocation. The placenta expresses receptors for numerous hormones including glucocorticoids, insulin, insulin-like growth factors (IGFs), leptin, gonadal hormones, cytokines, and prostaglandins (Bodner *et al*, 1999; Fowden *et al*, 2014; Hiden *et al*, 2006; Keelan and Mitchell, 2007; McCormick *et al*, 1981; Unluedik *et al*, 2010). These signals are integrated by downstream cascades that ultimately influence placental growth, energy homeostasis, and endocrine action. For example, Jansson and Powell (2006) proposed that the mammalian target of rapamycin (mTOR) pathway serves as

a nutrient sensor that regulates transporter expression in order to facilitate compatibility between maternal supply and fetal demand. Circulating insulin and IGF1 activate the protein kinase activity of mTOR, leading to increased expression of genes promoting cell growth and metabolism (Roos *et al*, 2009).

Stress during pregnancy compromises the maternal hormonal milieu in humans and in animal models, where elevations in CRF, pro-inflammatory cytokines, and glucocorticoids have been reported (Parker and Douglas, 2010). The placenta integrates these signals and facilitates a coordinated, sex-specific response involving immune function, nutrient metabolism/transport, or its own endocrine actions (Fowden *et al*, 2014). For example, in mice, prenatal stress increased pro-inflammatory cytokines, including tumor necrosis factor- α and interleukin-6 in male, but not in female, placentas and the neurodevelopmental programming effects in this model were ameliorated by maternal anti-inflammatory treatment during stress exposure (Bronson and Bale, 2014; Mueller and Bale, 2008). The effects of immune dysregulation on the developing brain program endophenotypes of neurodevelopmental disorders, including autism and schizophrenia, and have been reviewed previously (Hsiao and Patterson, 2012).

A Common Programmatic Pathway

In addition to maternal psychosocial stress, fetal exposures involving stressful metabolic challenges are also associated with increased neuropsychiatric disease risk, including maternal diabetes, obesity, infection, and pre-eclampsia (Bale *et al*, 2010; Brown and Derkits, 2010; Walker *et al*, 2015). It is well established that maternal insulin is dysregulated in pregnancies complicated by diabetes, obesity, undernutrition, and preeclampsia, and insulin resistance also occurs in the placentas from diabetic, preeclamptic, and growth-restricted pregnancies (Colomiere *et al*, 2009; Rademacher *et al*, 2007; Scioscia *et al*, 2006; Street *et al*, 2011). Maternal infection during pregnancy increases cytokines within the placenta and is predicted to disrupt insulin signaling due to the inhibitory effects of cytokines on insulin action (Aguirre *et al*, 2002; Sykiotis and Papavassiliou, 2001; Tanti and Jager, 2009). This same mechanism is proposed to impair placental insulin signaling in pregnancies complicated by maternal psychosocial stress, as pro-inflammatory cytokines were increased in stress-exposed placentas and were associated with the male-biased endophenotype (Bronson and Bale, 2014). Programmatic effects in these conditions likely depend on the gestational age at which placental insulin signaling is affected, as insulin receptors are expressed in mammalian placental trophoblasts and fetal endothelial cells in a spatiotemporal pattern (Desoye *et al*, 1994). Functional analysis of insulin-responsive genes suggests that maternal insulin regulates placental metabolism (predominately of lipids and fatty acids) in early pregnancy, whereas fetal insulin

communicates demand for growth and cell proliferation/survival near term (Hiden *et al*, 2006). Therefore, insulin perturbation can influence placental function, nutrient availability, and intrauterine homeostasis throughout gestation (depicted in Figure 2) and vastly have an impact on neurodevelopmental programming.

Encoding of Stress Memories in the Epigenome

Epigenomic remodeling is increasingly recognized as a molecular bridge linking placental adaptive responses to adversity with long-term phenotypic outcomes. Epigenetic processes including DNA methylation, histone modifications, and changes in small noncoding RNA expression are dynamic mechanisms by which the environment can shape gene expression and placental function, and when maintained within the fetal germ cells can influence the phenotype of future generations (reviewed in Bale, 2015; Franklin *et al*, 2010; Maccani and Marsit, 2009; Marsit, 2015; Monk *et al*, 2012). Tight epigenetic regulation of gene expression is essential for normal developmental processes, including placentation, cell fate determination, genomic imprinting, and X-inactivation, in particular throughout critical developmental periods during which imprinted epigenetic marks are erased and re-established (Gabory *et al*, 2011; Rugg-Gunn, 2012).

Intrauterine conditions, especially perturbations in nutrient availability and metabolism, have dramatic effects on epigenetic machinery (Tarrade *et al*, 2015). For example, maternally derived micronutrients such as folate and choline serve as substrates for DNA methyltransferases (DNMT) and their deficiency decreases placental DNA methylation (Kim *et al*, 2009). In calorie-restricted as well as high-fat diet-exposed mice, distinct hypomethylation patterns were detected in male and female placentas, including at imprinted loci, suggesting sex-specific placental adaptive responses to nutritional state (Chen *et al*, 2013; Gallou-Kabani *et al*, 2010). Similarly, maternal stress in humans and rodents elicits sex-specific placental epigenetic adaptations, including changes in DNA methylation and DNMT expression that were associated with neurodevelopmental outcomes (Table 1, (Conradt *et al*, 2013; Jensen Peña *et al*, 2012; Mueller and Bale, 2008). Differential methylation of the placental 11 β HSD2 promoter has been the epigenetic mark most consistently correlated with gene expression levels and maternal stress (Conradt *et al*, 2013; Jensen Peña *et al*, 2012). Although the effects of stress on histone modifications and microRNA expression within the placenta have yet to be fully elucidated, it is clear that placental epigenetic processes maintain important sex differences in homeostatic strategies at the maternal–fetal interface. Sex-dependent molecular memories of the prenatal environment can then be encoded by the fetal epigenome and, when incorporated into the germline, communicated to future generations.




Figure 2. Insulin signaling as potential common programmatic placental pathway. A simplified schematic depicting the predicted impairment of insulin signaling within placentas complicated by diverse fetal exposures. In diabetes, obesity, and pre-eclampsia, changes in insulin receptor (InsR) localization, kinase activity, and substrate availability lead to placenta insulin resistance. Maternal stress and infection are predicted to also elicit insulin resistance, owing to the inhibitory effects of cytokines on insulin action. Predicted programmatic effects of placenta insulin resistance depend on exposure timing. In early pregnancy, InsRs are localized to maternal-facing trophoblasts and predominately regulate expression of genes related to metabolism of lipids and fatty acids. Such changes may have an impact on placental growth, trophoblast survival, and hormone secretion in early pregnancy. Later in gestation, their expression is restricted to the fetal endothelial cells where insulin communicates fetal demand for growth, cell proliferation, and cell survival.

CONCLUSIONS

Nearly 20%–40% of pregnancies are complicated by adverse intrauterine conditions, including maternal mental health disorders, diabetes, obesity, and preeclampsia (Ananth *et al*, 2013; Bennett *et al*, 2004; Dawson *et al*, 2015; DeSisto *et al*, 2014; Goodman *et al*, 2014; Rubertsson *et al*, 2014). These fetal exposures are significant risk factors for neuropsychiatric disease predisposition, in particular in male offspring. Growing evidence supports a critical role for the placenta in the deleterious and sex-specific effects of these fetal exposures on the developing brain. Although a great deal more investigation is necessary to elucidate the specific mechanisms by which the placenta imparts these critical signals to the developing embryo, the potential importance of the placenta as a biomarker of maternal stress and its potential for determining neurodevelopmental and neuropsychiatric disease risk is enormous.

FUNDING AND DISCLOSURE

The work described was funded by grants from NIH MH073030, MH087597, MH091258, MH099910 and MH104184. The authors declare no conflict of interest.

ACKNOWLEDGMENTS

We thank Ali Rodgers and Katie Morrison for reading of the manuscript.

REFERENCES

- Aguirre V, Werner ED, Giraud J, Lee YH, Shoelson SE, White MF (2002). Phosphorylation of Ser307 in insulin receptor substrate-1 blocks interactions with the insulin receptor and inhibits insulin action. *J Biol Chem* **277**: 1531–1537.
- Ananth CV, Keyes KM, Wapner RJ (2013). Pre-eclampsia rates in the United States, 1980–2010: age-period-cohort analysis. *BMJ* **347**: f6564.
- Andersen SL (2003). Trajectories of brain development: point of vulnerability or window of opportunity? *Neurosci Biobehav Rev* **27**: 3–18.
- Anderson GM, Jacobs-Stannard A, Chawarska K, Volkmar FR, Klinan HJ (2007). Placental trophoblast inclusions in autism spectrum disorder. *Biol Psychiatry* **61**: 487–491.
- Arck PC, Hecher K (2013). Fetomaternal immune cross-talk and its consequences for maternal and offspring's health. *Nat Med* **19**: 548–556.
- Atladóttir HO, Thorsen P, Østergaard L, Schendel DE, Lemcke S, Abdallah M *et al* (2010). Maternal infection requiring hospitalization during pregnancy and autism spectrum disorders. *J Autism Dev Disord* **40**: 1423–1430.
- Avishai-Eliner S, Brunson KL, Sandman CA, Baram TZ (2002). Stressed-out, or in (utero)? *Trends Neurosci* **25**: 518–524.
- Aye LMH, Keelan JA (2013). Placental ABC transporters, cellular toxicity and stress in pregnancy. *Chem Biol Interact* **203**: 456–466.
- Baio J (2012). Prevalence of autism spectrum disorders: autism and developmental disabilities monitoring network, 14 sites, United States, 2008. *MMWR Surveill Sum* **Volume 61** (Number 3): 1–19 Centers Dis Control Prev at <<http://eric.ed.gov/?id=ED530639>>.

- Bale TL (2011). Sex differences in prenatal epigenetic programming of stress pathways. *Stress* **14**: 348–356.
- Bale TL (2015). Epigenetic and transgenerational reprogramming of brain development. *Nat Rev Neurosci* **16**: 332–344.
- Bale TL, Baram TZ, Brown AS, Goldstein JM, Insel TR, McCarthy MM *et al* (2010). Early life programming and neurodevelopmental disorders. *Biol Psychiatry* **68**: 314–319.
- Barker DJ (1997). Maternal nutrition, fetal nutrition, and disease in later life. *Nutrition* **13**: 807–813.
- Basiliou A, Yager J, Fehlings MG (2014). Neurological outcomes of animal models of uterine artery ligation and relevance to human intrauterine growth restriction: a systematic review. *Dev Med Child Neurol* **57**: 420–430.
- Beninson LA, Fleschner M (2014). Exosomes: an emerging factor in stress-induced immunomodulation. *Semin Immunol* **26**: 394–401.
- Bennett HA, Einarson A, Taddio A, Koren G, Einarson TR (2004). Prevalence of depression during pregnancy: systematic review. *Obstet Gynecol* **103**: 698–709.
- Beversdorf DQ, Manning SE, Hillier A, Anderson SL, Nordgren RE, Walters SE *et al* (2005). Timing of prenatal stressors and autism. *J Autism Dev Disord* **35**: 471–478.
- Blakeley PM, Capron LE, Jensen AB, O'Donnell KJ, Glover V (2013). Maternal prenatal symptoms of depression and down regulation of placental monoamine oxidase A expression. *J Psychosom Res* **75**: 341–345.
- Bloom B, Cohen RA, Freeman G (2011). Summary health statistics for U.S. children: National Health Interview Survey, 2010. *Vital Health Stat* **10**: 1–80 at <<http://www.ncbi.nlm.nih.gov/pubmed/22338334>>.
- Bock J, Rether K, Gröger N, Xie L, Braun K (2014). Perinatal programming of emotional brain circuits: an integrative view from systems to molecules. *Front Neurosci* **8**: 11.
- Bock J, Wainstock T, Braun K, Segal M (2015). Stress in utero: prenatal programming of brain plasticity and cognition. *Biol Psychiatry* **78**: 315–326.
- Bodner J, Ebenbichler CF, Wolf HJ, Müller-Holzner E, Stanzl U, Gander R *et al* (1999). Leptin receptor in human term placenta: in situ hybridization and immunohistochemical localization. *Placenta* **20**: 677–682.
- Boite AC, Geijn HP, van Dekker GA (2001). Pathophysiology of preeclampsia and the role of serotonin. *Eur J Obstet Gynecol Reprod Biol* **95**: 12–21.
- Bonin A, Levitt P (2011). Fetal, maternal, and placental sources of serotonin and new implications for developmental programming of the brain. *Neuroscience* **197**: 1–7.
- Bowen JM, Chamley L, Keelan JA, Mitchell MD (2002). Cytokines of the placenta and extra-placental membranes: roles and regulation during human pregnancy and parturition. *Placenta* **23**: 257–273.
- Bowman ME, Lopata A, Jaffe RB, Golos TG, Wickings J, Smith R (2001). Corticotropin-releasing hormone-binding protein in primates. *Am J Primatol* **53**: 123–130.
- Brett KE, Ferraro ZM, Yockell-Lelièvre J, Gruslin A, Adamo KB (2014). Maternal-fetal nutrient transport in pregnancy pathologies: the role of the placenta. *Int J Mol Sci* **15**: 16153–16185.
- Bronson SL, Bale TL (2014). Prenatal stress-induced increases in placental inflammation and offspring hyperactivity are male-specific and ameliorated by maternal antiinflammatory treatment. *Endocrinology* **155**: 2635–2646.
- Brown AS, Derkots EJ (2010). Prenatal infection and schizophrenia: a review of epidemiologic and translational studies. *Am J Psychiatry* **167**: 261–280.
- Brown RW, Diaz R, Robson AC, Kotelevtsev Y V, Mullins JJ, Kaufman MH *et al* (1996). The ontogeny of 11 beta-hydroxysteroid dehydrogenase type 2 and mineralocorticoid receptor gene expression reveal intricate control of glucocorticoid action in development. *Endocrinology* **137**: 794–797.
- Brunton PJ, Russell JA (2010). Prenatal social stress in the rat programmes neuroendocrine and behavioural responses to stress in the adult offspring: sex-specific effects. *J Neuroendocrinol* **22**: 258–271.
- Buckberry S, Bianco-Miotto T, Bent SJ, Dekker GA, Roberts CT (2014). Integrative transcriptome meta-analysis reveals widespread sex-biased gene expression at the human fetal-maternal interface. *Mol Hum Reprod* **20**: 810–819.
- Buss C, Davis EP, Muftuler LT, Head K, Sandman CA (2010). High pregnancy anxiety during mid-gestation is associated with decreased gray matter density in 6–9-year-old children. *Psychoneuroendocrinology* **35**: 141–153.
- Buss C, Davis EP, Shahbaba B, Pruessner JC, Head K, Sandman CA (2012). Maternal cortisol over the course of pregnancy and subsequent child amygdala and hippocampus volumes and affective problems. *Proc Natl Acad Sci USA* **109**: E1312–E1319.
- Carter AM (2012). Evolution of placental function in mammals: the molecular basis of gas and nutrient transfer, hormone secretion, and immune responses. *Physiol Rev* **92**: 1543–1576.
- Chen P-Y, Ganguly A, Rubbi L, Orozco LD, Morselli M, Ashraf D *et al* (2013). Intrauterine caloric restriction affects placental DNA methylation and gene expression. *Physiol Genomics* **45**: 565–576.
- Clifton VL (2010). Review: Sex and the human placenta: mediating differential strategies of fetal growth and survival. *Placenta* **31 Suppl**: S33–S39.
- Coe CL, Kramer M, Czéh B, Gould E, Reeves AJ, Kirschbaum C *et al* (2003). Prenatal stress diminishes neurogenesis in the dentate gyrus of juvenile rhesus monkeys. *Biol Psychiatry* **54**: 1025–1034.
- Colomiere M, Permezel M, Riley C, Desoye G, Lappas M (2009). Defective insulin signaling in placenta from pregnancies complicated by gestational diabetes mellitus. *Eur J Endocrinol* **160**: 567–578.
- Conradt E, Lester BM, Appleton AA, Armstrong DA, Marsit CJ (2013). The roles of DNA methylation of NR3C1 and 11 β -HSD2 and exposure to maternal mood disorder in utero on newborn neurobehavior. *Epigenetics* **8**: 1321–1329.
- Cottrell EC, Seckl JR (2009). Prenatal stress, glucocorticoids and the programming of adult disease. *Front Behav Neurosci* **3**: 19.
- Cuffe JSM, Dickinson H, Simmons DG, Moritz KM (2011). Sex specific changes in placental growth and MAPK following short term maternal dexamethasone exposure in the mouse. *Placenta* **32**: 981–989.
- Darnaudéry M, Maccari S (2008). Epigenetic programming of the stress response in male and female rats by prenatal restraint stress. *Brain Res Rev* **57**: 571–585.
- Das UG, Sadiq HF, Soares MJ, Hay WW, Devaskar SU (1998). Time-dependent physiological regulation of rodent and ovine placental glucose transporter (GLUT-1) protein. *Am J Physiol* **274**: R339–R347.
- Davis EP, Pfaff D (2014). Sexually dimorphic responses to early adversity: implications for affective problems and autism spectrum disorder. *Psychoneuroendocrinology* **49**: 11–25.
- Davis EP, Sandman CA, Buss C, Wing DA, Head K (2013). Fetal glucocorticoid exposure is associated with preadolescent brain development. *Biol Psychiatry* **74**: 647–655.
- Dawson AL, Razzaghi H, Arth A, Canfield MA, Parker SE, Reefhuis J (2015). Maternal exposures in the National Birth Defects Prevention Study: time trends of selected exposures. *Birth Defects Res A Clin Mol Teratol*, **103**: 703–712.
- De Felici M (2013). Origin, migration and proliferation of human primordial germ cells. *Oogenesis* 19–37 (doi:10.1007/978-0-85729-826-3).
- DeSisto CL, Kim SY, Sharma AJ (2014). Prevalence estimates of gestational diabetes mellitus in the United States, Pregnancy Risk Assessment Monitoring System (PRAMS), 2007–2010. *Prev Chronic Dis* **11**: E104.
- Desoye G, Hartmann M, Blaschitz A, Dohr G, Hahn T, Kohnen G *et al* (1994). Insulin receptors in syncytiotrophoblast and fetal endothelium of human placenta. Immunohistochemical evidence for developmental changes in distribution pattern. *Histochemistry* **101**: 277–285.
- Du F, Cooper AJ, Thida T, Shinn AK, Cohen BM, Ongür D (2013). Myelin and axon abnormalities in schizophrenia measured with magnetic resonance imaging techniques. *Biol Psychiatry* **74**: 451–457.
- Dörner G (1973). [Possible significance of prenatal and/or perinatal nutrition for the pathogenesis of obesity]. *Acta Biol Med Ger* **30**: K19–K22.
- Emanuel RL, Robinson BG, Seely EW, Graves SW, Kohane I, Saltzman D *et al* (1994). Corticotrophin releasing hormone levels in human plasma and amniotic fluid during gestation. *Clin Endocrinol (Oxf)* **40**: 257–262.
- Fedorova I, Alveim AR, Hussein N, Salem N (2009). Deficit in prepulse inhibition in mice caused by dietary n-3 fatty acid deficiency. *Behav Neurosci* **123**: 1218–1225.
- Fowden AL, Forhead AJ, Coan PM, Burton GJ (2008). The placenta and intrauterine programming. *J Neuroendocrinol* **20**: 439–450.
- Fowden AL, Forhead AJ, Sferruzzi-Perri AN, Burton GJ, Vaughan OR (2014). Review: endocrine regulation of placental phenotype. *Placenta* **36 Suppl 1**: S50–S59.
- Fowden AL, Ward JW, Wooding FPB, Forhead AJ, Constanica M (2006). Programming placental nutrient transport capacity. *J Physiol* **572**: 5–15.
- Franklin TB, Russig H, Weiss IC, Gräff J, Linder N, Michalon A *et al* (2010). Epigenetic transmission of the impact of early stress across generations. *Biol Psychiatry* **68**: 408–415.
- Froehlich TE, Lanphear BP, Epstein JN, Barbaresi WJ, Katusic SK, Kahn RS (2007). Prevalence, recognition, and treatment of attention-deficit/hyperactivity disorder in a national sample of US children. *Arch Pediatr Adolesc Med* **161**: 857–864.
- Fujioka A, Fujioka T, Ishida Y, Maekawa T, Nakamura S (2006). Differential effects of prenatal stress on the morphological maturation of hippocampal neurons. *Neuroscience* **141**: 907–915.
- Fujioka T, Sakata Y, Yamaguchi K, Shibasaki T, Kato H, Nakamura S (1999). The effects of prenatal stress on the development of hypothalamic paraventricular neurons in fetal rats. *Neuroscience* **92**: 1079–1088.

- Gabory A, Attig L, Junien C (2011). Developmental programming and epigenetics. *Am J Clin Nutr* **94**: 1943S–1952S.
- Gabory A, Roseboom TJ, Moore T, Moore LG, Junien C (2013). Placental contribution to the origins of sexual dimorphism in health and diseases: sex chromosomes and epigenetics. *Biol Sex Differ* **4**: 5.
- Gallou-Kabani C, Gabory A, Tost J, Karimi M, Mayeur S, Lesage J *et al* (2010). Sex- and diet-specific changes of imprinted gene expression and DNA methylation in mouse placenta under a high-fat diet. *PLoS One* **5**: e14398.
- Gapp K, Ziegler L von, Tweedie-Cullen RY, Mansuy IM (2014). Early life epigenetic programming and transmission of stress-induced traits in mammals: how and when can environmental factors influence traits and their transgenerational inheritance? *Bioessays* **36**: 491–502.
- Georgiades P, Ferguson-Smith AC, Burton GJ (2002). Comparative developmental anatomy of the murine and human definitive placentae. *Placenta* **23**: 3–19.
- Gerardin P, Wendland J, Bodeau N, Galin A, Bialobos S, Tordjman S *et al* (2011). Depression during pregnancy: is the developmental impact earlier in boys? A prospective case-control study. *J Clin Psychiatry* **72**: 378–387.
- Goland RS, Wardlaw SL, Blum M, Tropper PJ, Stark RI (1988). Biologically active corticotropin-releasing hormone in maternal and fetal plasma during pregnancy. *Am J Obstet Gynecol* **159**: 884–890.
- Goodman JH, Chenausky KL, Freeman MP (2014). Anxiety disorders during pregnancy: a systematic review. *J Clin Psychiatry* **75**: e1153–e1184.
- Hewitt DP, Mark PJ, Waddell BJ (2006). Glucocorticoids prevent the normal increase in placental vascular endothelial growth factor expression and placental vascularity during late pregnancy in the rat. *Endocrinology* **147**: 5568–5574.
- Hidden U, Maier A, Bilban M, Ghaffari-Tabrizi N, Wadsack C, Lang I *et al* (2006). Insulin control of placental gene expression shifts from mother to foetus over the course of pregnancy. *Diabetologia* **49**: 123–131.
- Howerton CL, Bale TL (2014). Targeted placental deletion of OGT recapitulates the prenatal stress phenotype including hypothalamic mitochondrial dysfunction. *Proc Natl Acad Sci USA* **111**: 9639–9644.
- Howerton CL, Morgan CP, Fischer DB, Bale TL (2013). O-GlcNAc transferase (OGT) as a placental biomarker of maternal stress and reprogramming of CNS gene transcription in development. *Proc Natl Acad Sci USA* **110**: 5169–5174.
- Hsiao EY, Patterson PH (2012). Placental regulation of maternal-fetal interactions and brain development. *Dev Neurobiol* **72**: 1317–1326.
- Huttunen MO, Niskanen P (1978). Prenatal loss of father and psychiatric disorders. *Arch Gen Psychiatry* **35**: 429–431.
- Illsley NP (2000). Glucose transporters in the human placenta. *Placenta* **21**: 14–22.
- Innis SM (2005). Essential fatty acid transfer and fetal development. *Placenta* **26** (Suppl A): S70–S75.
- Jansson T, Powell TL (2006). IFPA 2005 Award in Placentology Lecture. Human placental transport in altered fetal growth: does the placenta function as a nutrient sensor? – a review. *Placenta* **27**(Suppl A): S91–S97.
- Jansson T, Powell TL (2007). Role of the placenta in fetal programming: underlying mechanisms and potential interventional approaches. *Clin Sci (Lond)* **113**: 1–13.
- Jarvie E, Hauguel-de-Mouzon S, Nelson SM, Sattar N, Catalano PM, Freeman DJ (2010). Lipotoxicity in obese pregnancy and its potential role in adverse pregnancy outcome and obesity in the offspring. *Clin Sci (Lond)* **119**: 123–129.
- Jensen Peña C, Monk C, Champagne FA (2012). Epigenetic effects of prenatal stress on 11 β -hydroxysteroid dehydrogenase-2 in the placenta and fetal brain. *PLoS One* **7**: e39791.
- Joëls M, Baram TZ (2009). The neuro-symphony of stress. *Nat Rev Neurosci* **10**: 459–466.
- Kapoor A, Kostaki A, Janus C, Matthews SG (2009). The effects of prenatal stress on learning in adult offspring is dependent on the timing of the stressor. *Behav Brain Res* **197**: 144–149.
- Kapoor A, Matthews SG (2005). Short periods of prenatal stress affect growth, behaviour and hypothalamo-pituitary-adrenal axis activity in male guinea pig offspring. *J Physiol* **566**: 967–977.
- Kapoor A, Matthews SG (2008). Prenatal stress modifies behavior and hypothalamic-pituitary-adrenal function in female guinea pig offspring: effects of timing of prenatal stress and stage of reproductive cycle. *Endocrinology* **149**: 6406–6415.
- Kawamura T, Chen J, Takahashi T, Ichitani Y, Nakahara D (2006). Prenatal stress suppresses cell proliferation in the early developing brain. *Neuroreport* **17**: 1515–1518.
- Keelan JA, Mitchell MD (2007). Placental cytokines and preeclampsia. *Front Biosci* **12**: 2706–2727.
- Kent A, Hughes P, Ormerod L, Jones G, Thilaganathan B (2002). Uterine artery resistance and anxiety in the second trimester of pregnancy. *Ultrasound Obstet Gynecol* **19**: 177–179.
- Khashan AS, Abel KM, McNamee R, Pedersen MG, Webb RT, Baker PN *et al* (2008). Higher risk of offspring schizophrenia following antenatal maternal exposure to severe adverse life events. *Arch Gen Psychiatry* **65**: 146–152.
- Kim J-M, Hong K, Lee JH, Lee S, Chang N (2009). Effect of folate deficiency on placental DNA methylation in hyperhomocysteinemic rats. *J Nutr Biochem* **20**: 172–176.
- King S, Laplante DP (2005). The effects of prenatal maternal stress on children's cognitive development: Project Ice Storm. *Stress* **8**: 35–45.
- Kinney DK, Miller AM, Crowley DJ, Huang E, Gerber E (2008a). Autism prevalence following prenatal exposure to hurricanes and tropical storms in Louisiana. *J Autism Dev Disord* **38**: 481–488.
- Kinney DK, Munir KM, Crowley DJ, Miller AM (2008b). Prenatal stress and risk for autism. *Neurosci Biobehav Rev* **32**: 1519–1532.
- Koutmani Y, Politis PK, Elkouris M, Agrogiannis G, Kemerli M, Patsouris E *et al* (2013). Corticotropin-releasing hormone exerts direct effects on neuronal progenitor cells: implications for neuroprotection. *Mol Psychiatry* **18**: 300–307.
- Kraszpulski M, Dickerson PA, Salm AK (2006). Prenatal stress affects the developmental trajectory of the rat amygdala. *Stress* **9**: 85–95.
- Lager S, Powell TL (2012). Regulation of nutrient transport across the placenta. *J Pregnancy* **2012**: 179827.
- Lane RH, Ramirez RJ, Sirka AE, Kloesz JL, McLaughlin MK, Gruetzmacher EM *et al* (2001). Uteroplacental insufficiency lowers the threshold towards hypoxia-induced cerebral apoptosis in growth-retarded fetal rats. *Brain Res* **895**: 186–193.
- Laplante DP, Brunet A, Schmitz N, Ciampi A, King S (2008). Project Ice Storm: prenatal maternal stress affects cognitive and linguistic functioning in 5½-year-old children. *J Am Acad Child Adolesc Psychiatry* **47**: 1063–1072.
- Lemaire V, Koehl M, Le Moal M, Abrous DN (2000). Prenatal stress produces learning deficits associated with an inhibition of neurogenesis in the hippocampus. *Proc Natl Acad Sci USA* **97**: 11032–11037.
- Li J, Olsen J, Vestergaard M, Obel C (2010). Attention-deficit/hyperactivity disorder in the offspring following prenatal maternal bereavement: a nationwide follow-up study in Denmark. *Eur Child Adolesc Psychiatry* **19**: 747–753.
- Li J, Wang Z-N, Chen Y-P, Dong Y-P, Shuai H-L, Xiao X-M *et al* (2012). Late gestational maternal serum cortisol is inversely associated with fetal brain growth. *Neurosci Biobehav Rev* **36**: 1085–1092.
- Li Y, Hadden C, Singh P, Mercado CP, Murphy P, Dajani NK *et al* (2014). GDM-associated insulin deficiency hinders the dissociation of SERT from ERp44 and down-regulates placental 5-HT uptake. *Proc Natl Acad Sci USA* **111**: E5697–E5705.
- Lindgaard ML, Wassif CA, Vaisman B, Amar M, Wasmuth E V, Shamburek R *et al* (2008). Characterization of placental cholesterol transport: ABCA1 is a potential target for in utero therapy of Smith-Lemli-Opitz syndrome. *Hum Mol Genet* **17**: 3806–3813.
- Lou HC, Hansen D, Nordentoft M, Pryds O, Jensen F, Nim J *et al* (1994). Prenatal stressors of human life affect fetal brain development. *Dev Med Child Neurol* **36**: 826–832.
- Lucassen PJ, Bosch OJ, Jousma E, Krömer SA, Andrew R, Seckl JR (2009). Prenatal stress reduces postnatal neurogenesis in rats selectively bred for high, but not low, anxiety: possible key role of placental 11 β -hydroxysteroid dehydrogenase type 2. *Eur J Neurosci* **29**: 97–103.
- Maccani MA, Marsit CJ (2009). Epigenetics in the placenta. *Am J Reprod Immunol* **62**: 78–89.
- Madore C, Nadjar A, Delpech J-C, Sere A, Aubert A, Portal C *et al* (2014). Nutritional n-3 PUFAs deficiency during perinatal periods alters brain innate immune system and neuronal plasticity-associated genes. *Brain Behav Immun* **41**: 22–31.
- Mairesse J, Lesage J, Breton C, Bréant B, Hahn T, Darnaudey M *et al* (2007). Maternal stress alters endocrine function of the feto-placental unit in rats. *Am J Physiol Endocrinol Metab* **292**: E1526–E1533.
- Mandyam CD, Crawford EF, Eisch AJ, Rivier CL, Richardson HN (2008). Stress experienced in utero reduces sexual dichotomies in neurogenesis, microenvironment, and cell death in the adult rat hippocampus. *Dev Neurobiol* **68**: 575–589.
- Mao J, Zhang X, Sieli PT, Faldut MT, Torres KE, Rosenfeld CS (2010). Contrasting effects of different maternal diets on sexually dimorphic gene expression in the murine placenta. *Proc Natl Acad Sci USA* **107**: 5557–5562.
- Markham JA, Koenig JI (2011). Prenatal stress: Role in psychotic and depressive diseases. *Psychopharmacology (Berl)* **214**: 89–106.
- Marsit CJ (2015). Influence of environmental exposure on human epigenetic regulation. *J Exp Biol* **218**: 71–79.
- McCormick PD, Razel AJ, Spelsberg TC, Coulam CB (1981). Evidence for an androgen receptor in the human placenta. *Am J Obstet Gynecol* **140**: 8–13.

- Monk C, Spicer J, Champagne FA (2012). Linking prenatal maternal adversity to developmental outcomes in infants: the role of epigenetic pathways. *Dev Psychopathol* **24**: 1361–1376.
- Morgan CP, Bale TL (2011). Early prenatal stress epigenetically programs dysmasculinization in second-generation offspring via the paternal lineage. *J Neurosci* **31**: 11748–11755.
- Morgane PJ, Austin-LaFrance R, Bronzino J, Tonkiss J, Diaz-Cintra S, Cintra L *et al* (1993). Prenatal malnutrition and development of the brain. *Neurosci Biobehav Rev* **17**: 91–128.
- Mueller BR, Bale TL (2007). Early prenatal stress impact on coping strategies and learning performance is sex dependent. *Physiol Behav* **91**: 55–65.
- Mueller BR, Bale TL (2008). Sex-specific programming of offspring emotionality after stress early in pregnancy. *J Neurosci* **28**: 9055–9065.
- Murmu MS, Salomon S, Biala Y, Weinstock M, Braun K, Bock J (2006). Changes of spine density and dendritic complexity in the prefrontal cortex in offspring of mothers exposed to stress during pregnancy. *Eur J Neurosci* **24**: 1477–1487.
- Myatt L (2006). Placental adaptive responses and fetal programming. *J Physiol* **572**: 25–30.
- Nguyen TT, Tseng YT, McGonnigal B, Stabila JP, Worrell LA, Saha S *et al* (1999). Placental biogenic amine transporters: in vivo function, regulation and pathobiological significance. *Placenta* **20**: 3–11.
- van Os J, Selten JP (1998). Prenatal exposure to maternal stress and subsequent schizophrenia. The May 1940 invasion of The Netherlands. *Br J Psychiatry* **172**: 324–326.
- Ouyang Y, Mouillet J-F, Coyne CB, Sadovsky Y (2014). Review: placenta-specific microRNAs in exosomes - good things come in nano-packages. *Placenta* **35 Suppl**: S69–S73.
- O'Connell BA, Moritz KM, Walker DW, Dickinson H (2013). Sexually dimorphic placental development throughout gestation in the spiny mouse (*Acomys cahirinus*). *Placenta* **34**: 119–126.
- O'Donnell KJ, Bugge Jensen A, Freeman L, Khalife N, O'Connor TG, Glover V (2012). Maternal prenatal anxiety and downregulation of placental 11 β -HSD2. *Psychoneuroendocrinology* **37**: 818–826.
- Pankevich DE, Mueller BR, Brockel B, Bale TL (2009). Prenatal stress programming of offspring feeding behavior and energy balance begins early in pregnancy. *Physiol Behav* **98**: 94–102.
- Parker VJ, Douglas AJ (2010). Stress in early pregnancy: maternal neuro-endocrine-immune responses and effects. *J Reprod Immunol* **85**: 86–92.
- Petraglia F, Sawchenko PE, Rivier J, Vale W (1987). Evidence for local stimulation of ACTH secretion by corticotropin-releasing factor in human placenta. *Nature* **328**: 717–719.
- Ponder KL, Salisbury A, McGonnigal B, Laliberte A, Lester B, Padbury JF (2011). Maternal depression and anxiety are associated with altered gene expression in the human placenta without modification by antidepressant use: implications for fetal programming. *Dev Psychobiol* **53**: 711–723.
- Qiu A, Anh TT, Li Y, Chen H, Rifkin-Graboi A, Broekman BFP *et al* (2015). Prenatal maternal depression alters amygdala functional connectivity in 6-month-old infants. *Transl Psychiatry* **5**: e508.
- Qiu A, Rifkin-Graboi A, Chen H, Chong Y-S, Kwek K, Gluckman PD *et al* (2013). Maternal anxiety and infants' hippocampal development: timing matters. *Transl Psychiatry* **3**: e306.
- Rademacher TW, Gumaa K, Scioscia M (2007). Preeclampsia, insulin signalling and immunological dysfunction: a fetal, maternal or placental disorder? *J Reprod Immunol* **76**: 78–84.
- Rayen I, Hove DL van den, Prickaerts J, Steinbusch HW, Pawluski JL (2011). Fluoxetine during development reverses the effects of prenatal stress on depressive-like behavior and hippocampal neurogenesis in adolescence. *PLoS One* **6**: e24003.
- Reid MV, Murray KA, Marsh ED, Golden JA, Simmons RA, Grinspan JB (2012). Delayed myelination in an intrauterine growth retardation model is mediated by oxidative stress upregulating bone morphogenetic protein 4. *J Neuropathol Exp Neurol* **71**: 640–653.
- Reis FM, Florio P, Cobellis L, Luisi S, Severi FM, Bocchi C *et al* (2001). Human placenta as a source of neuroendocrine factors. *Biol Neonate* **79**: 150–156.
- Riccio O, Potter G, Walzer C, Vallet P, Szabó G, Vutskits L *et al* (2009). Excess of serotonin affects embryonic interneuron migration through activation of the serotonin receptor 6. *Mol Psychiatry* **14**: 280–290.
- Rifkin-Graboi A, Bai J, Chen H, Hameed WB, Sim LW, Tint MT *et al* (2013). Prenatal maternal depression associates with microstructure of right amygdala in neonates at birth. *Biol Psychiatry* **74**: 837–844.
- Robinson BG, Arbiser JL, Emanuel RL, Majzoub JA (1989). Species-specific placental corticotropin releasing hormone messenger RNA and peptide expression. *Mol Cell Endocrinol* **62**: 337–341.
- Rodriguez A, Bohlin G (2005). Are maternal smoking and stress during pregnancy related to ADHD symptoms in children? *J Child Psychol Psychiatry* **46**: 246–254.
- Ronald A, Pennell CE, Whitehouse AJO (2010). Prenatal maternal stress associated with ADHD and autistic traits in early childhood. *Front Psychol* **1**: 223.
- Roos S, Powell TL, Jansson T (2009). Placental mTOR links maternal nutrient availability to fetal growth. *Biochem Soc Trans* **37**: 295–298.
- Rossant J, Cross JC (2001). Placental development: lessons from mouse mutants. *Nat Rev Genet* **2**: 538–548.
- Rubertsson C, Hellström J, Cross M, Sydsjö G (2014). Anxiety in early pregnancy: prevalence and contributing factors. *Arch Womens Ment Health* **17**: 221–228.
- Rugg-Gunn PJ (2012). Epigenetic features of the mouse trophoblast. *Reprod Biomed Online* **25**: 21–30.
- Sagawa N, Yura S, Itoh H, Kakui K, Takemura M, Nuamah MA *et al* (2002). Possible role of placental leptin in pregnancy: a review. *Endocrine* **19**: 65–71.
- Saitou M, Yamaji M (2012). Primordial germ cells in mice. *Cold Spring Harb Perspect Biol* **4**.
- Sandman CA (2015). Fetal exposure to placental corticotropin-releasing hormone (pCRH) programs developmental trajectories. *Peptides* 10.1016/j.peptides.2015.03.020 (in press).
- Sarkar S, Craig MC, Dell'Acqua F, O'Connor TG, Catani M, Deeley Q *et al* (2014). Prenatal stress and limbic-prefrontal white matter microstructure in children aged 6–9 years: a preliminary diffusion tensor imaging study. *World J Biol Psychiatry* **15**: 346–352.
- Schneider ML (1992). Prenatal stress exposure alters postnatal behavioral expression under conditions of novelty challenge in rhesus monkey infants. *Dev Psychobiol* **25**: 529–540.
- Schneider ML, Coe CL (1993). Repeated social stress during pregnancy impairs neuromotor development of the primate infant. *J Dev Behav Pediatr* **14**: 81–87.
- Schneider ML, Roughton EC, Koehler AJ, Lubach GR (1999). Growth and development following prenatal stress exposure in primates: an examination of ontogenetic vulnerability. *Child Dev* **70**: 263–274.
- Scioscia M, Gumaa K, Kunjara S, Paine MA, Selvaggi LE, Rodeck CH *et al* (2006). Insulin resistance in human preeclamptic placenta is mediated by serine phosphorylation of insulin receptor substrate-1 and -2. *J Clin Endocrinol Metab* **91**: 709–717.
- Seckl JR, Holmes MC (2007). Mechanisms of disease: glucocorticoids, their placental metabolism and fetal “programming” of adult pathophysiology. *Nat Clin Pract Endocrinol Metab* **3**: 479–488.
- Sjöström K, Valentin L, Thelin T, Marsál K (1997). Maternal anxiety in late pregnancy and fetal hemodynamics. *Eur J Obstet Gynecol Reprod Biol* **74**: 149–155.
- Sood R, Zehnder JL, Druzin ML, Brown PO (2006). Gene expression patterns in human placenta. *Proc Natl Acad Sci USA* **103**: 5478–5483.
- Street ME, Viani I, Ziveri MA, Volta C, Smerieri A, Bernasconi S (2011). Impairment of insulin receptor signal transduction in placentas of intra-uterine growth-restricted newborns and its relationship with fetal growth. *Eur J Endocrinol* **164**: 45–52.
- Suenaga T, Yukie M, Gao S, Nakahara D (2012). Sex-specific effects of prenatal stress on neuronal development in the medial prefrontal cortex and the hippocampus. *Neuroreport* **23**: 430–435.
- Sykietis GP, Papavassiliou AG (2001). Serine phosphorylation of insulin receptor substrate-1: a novel target for the reversal of insulin resistance. *Mol Endocrinol* **15**: 1864–1869.
- Tam O, Innis SM (2006). Dietary polyunsaturated fatty acids in gestation alter fetal cortical phospholipids, fatty acids and phosphatidylserine synthesis. *Dev Neurosci* **28**: 222–229.
- Tanti J-F, Jager J (2009). Cellular mechanisms of insulin resistance: role of stress-regulated serine kinases and insulin receptor substrates (IRS) serine phosphorylation. *Curr Opin Pharmacol* **9**: 753–762.
- Tarrade A, Panchenko P, Junien C, Gabory A (2015). Placental contribution to nutritional programming of health and diseases: epigenetics and sexual dimorphism. *J Exp Biol* **218**: 50–58.
- Tegethoff M, Greene N, Olsen J, Meyer AH, Meinschmidt G (2010). Maternal psychosocial stress during pregnancy and placenta weight: evidence from a national cohort study. *PLoS One* **5**: e14478.
- Teixeira JM, Fisk NM, Glover V (1999). Association between maternal anxiety in pregnancy and increased uterine artery resistance index: cohort based study. *BMJ* **318**: 153–157.
- Thordstein M, Andiné P, Lehmann A, Hagberg H (1992). Cerebral amino acids and energy metabolites in the growth retarded rat fetus under normoxia and hypoxia. *J Dev Physiol* **18**: 59–65.
- Unlagedik E, Alifaidy N, Holloway A, Lye S, Bocking A, Challis J *et al* (2010). Expression and regulation of prostaglandin receptors in the human placenta and fetal membranes at term and preterm. *Reprod Fertil Dev* **22**: 796–807.

- Velasquez JC, Goeden N, Bonnin A (2013). Placental serotonin: implications for the developmental effects of SSRIs and maternal depression. *Front Cell Neurosci* **7**: 47.
- Verhaagh S, Barlow DP, Zwart R (2001). The extraneuronal monoamine transporter Slc22a3/Orc13 co-localizes with the Maa0 metabolizing enzyme in mouse placenta. *Mech Dev* **100**: 127–130.
- Walder DJ, Laplante DP, Sousa-Pires A, Veru F, Brunet A, King S (2014). Prenatal maternal stress predicts autism traits in 6½ year-old children: Project Ice Storm. *Psychiatry Res* **219**: 353–360.
- Walker CK, Anderson KW, Milano KM, Ye S, Tancredi DJ, Pessah IN *et al* (2013). Trophoblast inclusions are significantly increased in the placentas of children in families at risk for autism. *Biol Psychiatry* **74**: 204–211.
- Walker CK, Krakowiak P, Baker A, Hansen RL, Ozonoff S, Hertz-Picciotto I (2015). Preeclampsia, placental insufficiency, and autism spectrum disorder or developmental delay. *JAMA Pediatr* **169**: 154–162.
- Watson ED, Cross JC (2005). Development of structures and transport functions in the mouse placenta. *Physiology (Bethesda)* **20**: 180–193.
- Weinstock M (2008). The long-term behavioural consequences of prenatal stress. *Neurosci Biobehav Rev* **32**: 1073–1086.
- Weinstock M (2011). Sex-dependent changes induced by prenatal stress in cortical and hippocampal morphology and behaviour in rats: an update. *Stress* **14**: 604–613.
- Wyrwoll CS, Seckl JR, Holmes MC (2009). Altered placental function of 11beta-hydroxysteroid dehydrogenase 2 knockout mice. *Endocrinology* **150**: 1287–1293.
- Xu J, Yang B, Yan C, Hu H, Cai S, Liu J *et al* (2013). Effects of duration and timing of prenatal stress on hippocampal myelination and synaptophysin expression. *Brain Res* **1527**: 57–66.
- Yoshida S, Wada Y (2005). Transfer of maternal cholesterol to embryo and fetus in pregnant mice. *J Lipid Res* **46**: 2168–2174.
- Zhang XY, Chen DC, Xiu MH, De Yang F, Haile CN, Kosten TA *et al* (2012). Gender differences in never-medicated first-episode schizophrenia and medicated chronic schizophrenia patients. *J Clin Psychiatry* **73**: 1025–1033.
- Zikopoulos B, Barbas H (2010). Changes in prefrontal axons may disrupt the network in autism. *J Neurosci* **30**: 14595–14609.