

Role of a Lateral Orbital Frontal Cortex-Basolateral Amygdala Circuit in Cue-Induced Cocaine-Seeking Behavior

Amy A Arguello¹, Ben D Richardson¹, Jacob L Hall¹, Rong Wang¹, Matthew A Hodges², Marshall P Mitchell¹, Garret D Stuber², David J Rossi¹ and Rita A Fuchs^{*,1}

¹Department of Integrative Physiology and Neuroscience, Washington State University, College of Veterinary Medicine, Pullman, WA, USA;

²Department of Psychiatry and Neuroscience Center, University of North Carolina at Chapel Hill, Chapel Hill, NC, USA

Cocaine addiction is a disease characterized by chronic relapse despite long periods of abstinence. The lateral orbitofrontal cortex (IOFC) and basolateral amygdala (BLA) promote cocaine-seeking behavior in response to drug-associated conditioned stimuli (CS) and share dense reciprocal connections. Hence, we hypothesized that monosynaptic projections between these brain regions mediate CS-induced cocaine-seeking behavior. Male Sprague-Dawley rats received bilateral infusions of a Cre-dependent adeno-associated viral (AAV) vector expressing enhanced halorhodopsin 3.0 fused with a reporter protein (NpHR-mCherry) or a control AAV (mCherry) plus optic fiber implants into the IOFC (Experiment 1) or BLA (Experiment 2). The same rats also received bilateral infusions of a retrogradely transported AAV vector expressing Cre recombinase (Retro-Cre-GFP) into the BLA (Experiment 1) or IOFC (Experiment 2). Thus, NpHR-mCherry or mCherry expression was targeted to IOFC neurons that project to the BLA or to BLA neurons that project to the IOFC in different groups. Rats were trained to lever press for cocaine infusions paired with 5-s CS presentations. Responding was then extinguished. At test, response-contingent CS presentation was discretely coupled with optogenetic inhibition (5-s laser activation) or no optogenetic inhibition while lever responding was assessed without cocaine/food reinforcement. Optogenetic inhibition of IOFC to BLA, but not BLA to IOFC, projections in the NpHR-mCherry groups disrupted CS-induced reinstatement of cocaine-seeking behavior relative to (i) no optogenetic inhibition or (ii) manipulations in mCherry control or (iii) NpHR-mCherry food control groups. These findings suggest that the IOFC sends requisite input to the BLA, via monosynaptic connections, to promote CS-induced cocaine-seeking behavior.

Neuropsychopharmacology (2017) **42**, 727–735; doi:10.1038/npp.2016.157; published online 21 September 2016

INTRODUCTION

Cocaine addiction is a chronic disease characterized by strong propensity for relapse upon exposure to previously cocaine-associated environmental stimuli even after long periods of abstinence (Ehrman *et al*, 1992). Thus, cue-induced relapse represents an enduring challenge for the successful treatment of cocaine addiction. Several brain regions have requisite roles in cocaine-seeking behavior produced by previously drug-paired conditioned stimuli (CS) in rodent models of cue-induced drug relapse (Bossert *et al*, 2013; Lasseter *et al*, 2010); however, few studies have aimed to identify functionally significant connections between these brain regions. As a result, our understanding of the neural circuitry by which drug-associated cues elicit motivation to seek cocaine is limited.

In the present study, we focused on the lateral orbitofrontal cortex (IOFC) and basolateral amygdala (BLA) given the importance of these brain regions in assessing the

motivational significance of drug-associated CSs. In support of this, cocaine-dependent individuals exhibit increased cerebral blood flow in the OFC and BLA upon exposure to images containing drug-related cues (Childress *et al*, 1999; Goldstein and Volkow, 2002). Moreover, GABA_B agonist treatment decreases OFC and BLA activation and self-reported craving in response to drug-related cues (Brebner *et al*, 2002; Young *et al*, 2014). In rats, GABA_{A/B} agonist-induced inhibition of the IOFC (Fuchs *et al*, 2004) or tetrodotoxin-induced inactivation of the BLA (Grimm and See, 2000) decreases CS-induced reinstatement of cocaine-seeking behavior, indicating that these brain regions critically control the retrieval and/or utilization of cocaine-CS associations that guide drug-seeking behavior.

The IOFC and BLA share dense reciprocal connections (Lasseter *et al*, 2014; Price, 2007) and may interact to promote CS-induced reinstatement of drug-seeking behavior. Functional disconnection studies have shown that the IOFC and BLA work together to encode stimulus-outcome associations (Saddoris *et al*, 2005), facilitate flexible decision-making (Zeeb and Winstanley, 2013), and control drug context-induced reinstatement of drug-seeking behavior (Lasseter *et al*, 2011). However, the reciprocal nature of anatomical connections between the IOFC and BLA has limited the interpretability of these functional disconnection

*Correspondence: Dr RA Fuchs, Department of Integrative Physiology and Neuroscience, Washington State University, College of Veterinary Medicine, PO Box 647620, Pullman, WA 99164-7620, USA, Tel: +509 335 6164, Fax: +509 335 4650, E-mail: ritafuchs@vetmed.wsu.edu
Received 5 May 2016; revised 8 August 2016; accepted 11 August 2016; accepted article preview online 18 August 2016

effects, as the direction of information transfer could not be determined using this methodology. Furthermore, the significance of IOFC–BLA interactions in CS-induced cocaine-seeking behavior, in particular, has not been investigated.

To separately investigate whether CS-induced cocaine-seeking behavior depends on monosynaptic input from the IOFC to the BLA or *vice versa*, we expressed cre-recombinase in adeno-associated viral (AAV) vector serotype 6 (Retro-Cre-GFP), which exhibits retrograde transport in rodents and primates (Salegio *et al*, 2013; San Sebastian *et al*, 2013). To permit optogenetic inhibition of either IOFC neurons that project to the BLA (Experiment 1) or BLA neurons that project to the IOFC (Experiment 2), Retro-Cre-GFP was infused at the terminal region, while a cre-dependent AAV containing enhanced halorhodopsin 3.0 fused to an mCherry reporter protein (NpHR-mCherry) and indwelling optic fibers were introduced at the cell body region of the target circuit. At test, optogenetic inhibition was restricted to the duration of CS-presentations to minimize the influence of manipulations on the drug-paired context, which may regain some of its motivational effects upon CS presentation and contribute to drug-seeking behavior through the activation of neural substrates that promote context-induced reinstatement (Fuchs *et al*, 2008; Remedios *et al*, 2014). Thus, the present study examined the specific contributions of monosynaptic projections between the IOFC and BLA to the motivational effects of a drug-paired CS.

MATERIALS AND METHODS

Animals

Male Sprague-Dawley rats (250–275 g; Harlan Laboratories, Livermore, CA; $N=45$) were individually housed in a climate-controlled vivarium on a reversed light–dark cycle. Rats were maintained on 20–25 g of rat chow per day with water available *ad libitum*. All protocols for rat housing and treatment were approved by the Institutional Animal Care and Use Committee and followed the *Guide for the Care and Use of Laboratory Rats* (Institute of Laboratory Animal Resources on Life Sciences, 2011).

Viral Constructs

To construct AAV6-EF1 α -Cre-recombinase-IRES2-GFP (Retro-Cre-GFP, Figure 1a), pCAG-Cre-GFP (Addgene, plasmid #26646, deposited by Dr Anjen Chenn) was overexpressed, purified, and then cloned into an AAV backbone by the University of North Carolina (UNC) BAC Recombineering Core. Key regions of the plasmid were sequenced to confirm correct orientation. The AAV was then packaged into serotype 6 by the UNC Vector Core. All viruses were purchased from the UNC Vector Core (titers: 4.0–6.7 $\times 10^{12}$ vm/ml) and delivered at 0.7 μ l/hemisphere (0.1 μ l/min). The infusion needles (33-gauge) remained in place for 10 min post infusion. Maximal AAV-mediated gene expression occurs ~2–3 weeks post-infection (Carlezon *et al*, 1997; Penrod *et al*, 2015).




Figure 1 Circuit specific, cre-dependent expression of NpHR-mCherry. (a) Diagrams of the AAV (serotype 5) expressing a cre-dependent, enhanced halorhodopsin 3.0 fused to mCherry (NpHR-mCherry) and AAV (serotype 6) expressing Cre-recombinase-IRES2-GFP (Retro-Cre-GFP) utilized in Experiments 1 and 2. Both AAVs contained transgenes under the control of an EF1 α promoter. Representative images showing (b) expression of NpHR-mCherry in the IOFC and (c) in the BLA. Scale bar = 200 μ m, $\times 10$ magnification. NpHR-mCherry expression was colocalized with immunoreactivity for (d) the excitatory marker, CaMKII, but not for (e) the GABAergic marker glutamic acid decarboxylase-67 (GAD-67). Scale bar = 50 μ m, $\times 20$ magnification.

Surgery

Forty-eight hours after a single food-training session (described in Supplementary Methods), rats were anesthetized with ketamine and xylazine (80.0 mg/kg and 5.0 mg/kg, i.p., respectively). AAV5-EF1 α -DIO-eNpHR3.0-mCherry

(NpHR-mCherry) or AAV5-EF1 α -DIO-mCherry (mCherry) was infused bilaterally into the IOFC (Experiment 1; +3.5 mm AP, +2.8 mm ML, -5.0 mm DV from bregma) or the BLA (Experiment 2; -2.7 mm AP, +5.0 mm ML, -8.7 mm DV from bregma) (Figure 1a). Retro-Cre-GFP was infused bilaterally into the BLA (Experiment 1) or IOFC (Experiment 2). Optical fibers (200 μ m diameter; Thor Labs, Newton, NJ) encased in ceramic ferrules (1.25 mm O.D., 230 μ m I.D.; Thor Labs) were assembled in house (Sparta *et al*, 2012) and implanted bilaterally, 0.5 mm dorsal to the NpHR-mCherry or mCherry virus infusion sites. Screws and cranioplastic cement secured the ferrules to the skull. Additional rats received NpHR-mCherry into the IOFC either alone or with Retro-Cre-GFP into the BLA for immunohistochemical verification experiments, or NpHR-mCherry into BLA with Retro-Cre-GFP into the IOFC for slice electrophysiology experiments (see Supplementary Methods). Intravenous (i.v.) catheters were constructed in-house (Fuchs *et al*, 2007) and were inserted into the right jugular vein and threaded subcutaneously to exit on the back. Catheter patency was tested and maintained as described in Supplementary Methods.

Cocaine Self-Administration Training

Drug (Figures 3a and 4a) or food (Supplementary Figure S4A) self-administration training was conducted in sound-attenuated operant conditioning chambers (Coulbourn Instruments, Allentown, PA) during the rats' dark cycle for 2 h each day. Throughout each session, a red house light (0.4 fc brightness) was illuminated on the wall opposite to levers. During the cocaine self-administration session, responses on a designated active lever resulted in 2-s cocaine infusions (cocaine hydrochloride dissolved in sterile saline; 0.15 mg/0.05 ml per infusion, i.v.; NIDA Drug Supply Program, Research Triangle Park, NC) under a fixed ratio 1 (FR1) reinforcement schedule with a 20-s timeout period. During the food self-administration sessions, active lever presses resulted in delivery of a single food pellet under gradually increasing FR schedules with 20-s timeout periods (see Supplementary Methods). Each reinforcer (drug or food) was paired with 5-s presentation of a CS complex (white stimulus light above the active lever plus a 75 dB, 1 kHz tone). Active lever responses during the timeout periods and inactive lever responses during the session had no programmed consequences. Training was terminated when rats reached a criterion of ≥ 10 cocaine infusions per session on ≥ 10 days. Data collection and reinforcer delivery were controlled using Graphic State Notation software version 4.1.04 (Coulbourn Instruments).

Extinction Training

After the completion of drug or food self-administration training, all rats received daily 2-h extinction training sessions on at least 7 days until they reached the extinction criterion (≤ 25 active lever responses per session on two consecutive days). During the sessions, lever responses had no programmed consequences. During the last three sessions, the rats' indwelling optic fibers were connected bilaterally to mock optical patch cords to acclimatize them to the apparatus.

Testing

At test, all rats' optic fibers were connected bilaterally to mono-fiber optic patch cords (50 μ m I.D.; Doric Lenses, Quebec, CAN) encased in spring leashes (Plastics One). The patch cords were connected to one end of an optical commutator (1 \times 2 Fiber Optic Rotary Joint; Doric Lenses), to allow for free movement of rats and provide bilateral division of laser light from a single source. The other end of the optical commutator was connected to a solid state laser (Shanghai Laser & Optics Century, Shanghai, China) via an optic patch cable (105 μ m, 3 m; Thor Labs). The laser was located outside of the sound-attenuation chamber. Laser light (532 nm) was calibrated to 10–12 mW intensity at the tip of the patch cords immediately before testing (Sparta *et al*, 2012).

During two 1-h test sessions, active lever presses resulted in 5-s presentations of the previously cocaine-paired or food-paired CS either alone (Laser OFF) or discretely coupled with laser activation (Laser ON; continuous for 5 s). Each CS presentation initiated a 20-s timeout period during which active lever presses had no programmed consequences. The order of Laser ON and Laser OFF test sessions was counterbalanced, and the test sessions were separated by additional extinction sessions until active lever responding was ≤ 25 per session on two consecutive days.

Histology and Data Analysis

Rats were overdosed using ketamine and xylazine (240 mg/kg and 15 mg/kg, i.v., respectively) and transcardially perfused. Brain tissue was collected for assessment of optic fiber placement, immunohistochemical experiments, and whole-cell electrophysiological recordings (see Supplementary Methods). Rats with insufficient mCherry expression, off-target mCherry or GFP expression, or misplaced optic fibers were excluded from data analysis. All data were analyzed using analyses of variance (ANOVA) or *t*-tests, when appropriate (see Supplementary Methods). *Post-hoc* comparisons were done using Tukey's tests. Alpha was set at 0.05.

RESULTS

Histology

In the virus verification experiments, NpHR-mCherry expression was not observed in the IOFC unless Retro-Cre-GFP was infused into the BLA (Supplementary Figure S1A and B). Expression was robust within both the somal and dendritic compartments of cells within the IOFC and BLA, (Figure 1b and c), with similar numbers of mCherry-immunoreactive cells visible in each target region (Supplementary Figure S1G). NpHR-mCherry expression was exclusive to glutamatergic neurons, as indicated by co-localization with the excitatory marker calcium/calmodulin-dependent protein kinase II (CaMKII, Figure 1d) but no co-localization with the GABAergic marker glutamic acid decarboxylase-67 (GAD-67, Figure 1e). In Experiment 1, optic fiber tracts and NpHR-mCherry/mCherry expression were restricted to the IOFC (Figure 2a) and Retro-Cre-GFP expression was restricted to the BLA (Supplementary




Figure 2 Virus spread and optic fiber placements. Schematics depicting mCherry and NpHR-mCherry viral spread (shaded areas) and the most ventral point of optic fiber tracts for rats that received mCherry (white circles) or NpHR-mCherry (black circles) into the (a) IOFC (mCherry, $n=8$; NpHR-mCherry, $n=8$) or (b) BLA (mCherry, $n=6$; NpHR-mCherry, $n=6$). Numbers denote distance from bregma in mm on the schematics modified from the rat brain atlas of Paxinos and Watson (1997).

Figure S2A) ($n=8$ /group). In Experiment 2, optic fiber tracts and NpHR-mCherry/mCherry expression were restricted to BLA (Figure 2b) and Retro-Cre-GFP expression was restricted to the IOFC (Supplementary Figure S2B) ($n=6$ /group). mCherry expression was not visible in terminal regions (Supplementary Figure S1C–F).

Behavioral History and Inactive Lever Presses

The groups did not differ in drug or food intake or lever responding during self-administration and extinction training. They also did not differ in inactive lever responding during any phase of the experiment (Supplementary Figure S3). These findings and the results of the food control experiment (Supplementary Figure S4) are reported in Supplementary Results.

IOFC → BLA Circuit Inhibition Disrupts CS-Induced Cocaine-Seeking Behavior

Cocaine-seeking behavior depended on laser manipulation and opsin expression (ie, group: mCherry vs NpHR-mCherry; 2×2 ANOVA laser \times group interaction: $F_{1,14}=4.93$, $P=0.04$; group main effect: $F_{1,14}=5.38$, $P=0.04$; laser main effect: $F_{1,14}=2.75$, $P=0.12$). In the mCherry control group, active lever responding did not differ across the Laser ON and Laser OFF tests (Figure 3b). Conversely, in the NpHR-mCherry group, responding during the Laser ON test was lower than during the Laser OFF test ($P<0.05$) and compared with the mCherry group during the Laser ON test ($P<0.01$). A time course analysis of behavior during the Laser ON test indicated time-dependent group differences (2×3 ANOVA time \times group interaction: $F_{2,14}=3.87$, $P=0.03$; time main effect: $F_{2,14}=17.64$, $P=0.0001$; group main effect, $F_{1,14}=9.15$, $P=0.01$). Specifically, the NpHR-mCherry group exhibited fewer responses




Figure 3 Effect of optogenetic inhibition of IOFC afferents to the BLA on CS-induced reinstatement of cocaine-seeking behavior. (a) Schematic depicting the timeline of the behavioral experiments: cocaine self-administration training (SA), extinction training (EXT), and counterbalanced CS-induced reinstatement tests (1 h each) with or without laser stimulation. (b) Active lever responses (mean \pm SEM) during each phase of the experiment for the mCherry ($n=8$) and NpHR-mCherry ($n=8$) groups. (c) Time course of active lever responses (mean \pm SEM) during each 20-min bin of the Laser ON test. (d) Response rates (mean responses/second \pm SEM) on the active lever during the Laser ON test when active lever presses were reinforced with CS presentation (REIN) or were not reinforced during the 5-s CS presentation (CS) and the remaining 15-s (No CS) of the timeout period. (e) Total number of CS-reinforced active lever responses during the Laser ON test. Symbols represent difference between mCherry and NpHR-mCherry groups (*, b, c: Tukey's test, $P<0.05$; d, e: t -test, $P<0.05$), between the Laser OFF and Laser ON test sessions (#, Tukey's test, $P<0.05$), and change from Bin 1 (‡, Tukey's test, $P<0.05$).

than the mCherry group during the first 20 min of the Laser ON test (Figure 3c; $P<0.01$), after which responding decreased (Bin 1 > Bins 2–3; $P<0.01$).

Additional analyses evaluated possible changes in active lever response rate in relation to the termination of laser stimulation (Figure 3d). The NpHR-mCherry group exhibited lower CS-reinforced response rates than the mCherry group ($t_{14}=3.08$, $P=0.008$). Non-reinforced response rates did not differ between the groups during CS-coupled laser stimulation ($t_{14}=1.88$, $P=0.08$) or during the remainder of the timeout period ($t_{14}=0.42$, $P=0.68$), suggesting that responding did not rebound after laser stimulation was terminated. Furthermore, the NpHR-mCherry group exhibited fewer CS-reinforced responses than the mCherry group during the Laser ON test (Figure 3e; $t_{14}=3.01$, $P=0.009$).

BLA → IOFC Circuit Inhibition Fails to Alter CS-Induced Cocaine-Seeking Behavior

There were no differences in cocaine-seeking behavior between the groups (Figure 4b; 2×2 ANOVA laser \times group




Figure 4 Effect of optogenetic inhibition of BLA afferents to the IOFC on CS-induced reinstatement of cocaine-seeking behavior. (a) Schematic depicting the timeline of the behavioral experiments: cocaine self-administration training (SA), extinction training (EXT), and counterbalanced CS-induced reinstatement tests with or without laser stimulation (1 h each). (b) Active lever responses (mean ± SEM) during each phase of the experiment for the mCherry ($n=6$) and NpHR-mCherry ($n=6$) groups. (c) Time course of active lever responses (mean ± SEM) during each 20-min bin of the Laser ON test. (d) Response rates (mean responses/second ± SEM) on the active lever during the Laser ON test when active lever presses were reinforced with CS presentation (REIN) or were not reinforced during the 5-s CS presentation (CS) and the remaining 15-s (No CS) of the timeout period. (e) Total number of CS-reinforced active lever responses during the Laser ON test. Symbols represent change from Bin 1 (\ddagger , Tukey's test, $P<0.05$).

interaction, $F_{1,10}=0.46$, $P=0.51$, group main effect, $F_{1,10}=0.19$, $P=0.67$; laser main effect, $F_{1,14}=0.003$, $P=0.96$). A time course analysis of behavior during the Laser ON test revealed a time main effect only (Figure 4c; 2×3 ANOVA time main effect, $F_{2,10}=7.26$, $P=0.004$; laser \times group interaction, $F_{2,10}=0.98$, $P=0.39$; group main effect, $F_{1,10}=0.78$, $P=0.40$). Thus, active lever responding declined in both groups during the Laser ON test session (Bin 1 > Bin 2–3; $P<0.05$).

In terms of active lever response rates (Figure 4d), no differences were observed between the NpHR-mCherry and mCherry groups in CS-reinforced ($t_{10}=0.51$, $P=0.62$) and non-reinforced response rates during CS-coupled laser stimulation ($t_{10}=0.46$, $P=0.66$) or during the remainder of the timeout period ($t_{10}=1.11$, $P=0.29$). Thus, responding did not rebound after laser stimulation was terminated. Furthermore, no differences were observed between the groups in the total number of CS-reinforced responses during the Laser ON test (Figure 4e; $t_{10}=0.52$, $P=0.62$) even though BLA neurons exhibited light-induced hyperpolarization (Figure 5a and b) and inhibition of evoked action potential firing *ex vivo* (Figure 5c–e) (see Supplementary Results).

DISCUSSION

The present study examined the direction-specific contribution of monosynaptic projections between the IOFC and BLA in cue-induced cocaine-seeking behavior. To permit the selective inhibition of these circuits, a Retro-Cre-GFP was infused at the terminal region, and a Cre-dependent NpHR-mCherry was infused at the cell body region, of the target circuit. Cre-dependent expression of mCherry was observed in the cell body regions of the IOFC (Experiment 1) and BLA (Experiment 2) (Figure 1). Importantly, optogenetic inhibition of IOFC afferents to the BLA impaired CS-induced reinstatement of cocaine-seeking behavior compared with no optogenetic inhibition (Figure 3) and to manipulations in the mCherry control groups that did not express the opsin (Figure 3) and in the NpHR-mCherry groups in the food control experiment (Supplementary Figure S4). Optogenetic inhibition of BLA afferents to the IOFC did not impair CS-induced cocaine-seeking behavior (Figure 4) despite eliciting light-induced hyperpolarization and inhibition of evoked action potential firing (Figure 5).

Direct Input from the IOFC to the BLA is Necessary for CS-Induced Cocaine-Seeking Behavior

The present study demonstrates that the IOFC and BLA interact in a direction-specific manner to promote cue-induced reinstatement, extending localization studies that focused on the individual contributions of these brain regions to CS-induced reinstatement (Fuchs *et al*, 2004; Grimm and See, 2000) and disconnection studies that showed their interaction in drug context-induced reinstatement (Lasseter *et al*, 2011). In addition to the interpretational difficulty stemming from interhemispheric connections between the BLA and IOFC, results from functional disconnection did not provide insight into the direction of information processing between these brain regions. In contrast, the approach utilized in the present study has allowed us to demonstrate that information transfer from the IOFC to the BLA, in particular, is necessary for CS-induced reinstatement.

Importantly, the decrease in cocaine-seeking behavior following optogenetic inhibition of IOFC afferents to the BLA was not mediated by the inhibition of postsynaptic BLA neurons that project back to the IOFC (ie, via a IOFC \rightarrow BLA \rightarrow IOFC functional loop) as inhibition of BLA projections to the IOFC failed to alter this behavior (Figure 4b). In line with previous studies (Calu *et al*, 2013), behavioral impairment was also not due to hypoactivity or nonspecific performance deficits associated with AAV or laser exposure. Consistent with this, optogenetic inhibition of IOFC projections to the BLA failed to alter responding on the inactive lever (Supplementary Figure S3) or on the active lever during the timeout periods (Figure 3d). Furthermore, laser exposure failed to alter responding on the active lever in the mCherry group in Experiment 1 (Figure 3b) or in the NpHR-mCherry group that received optogenetic inhibition of BLA projections to the IOFC in Experiment 2 (Figure 4b). Finally, it is unlikely that IOFC \rightarrow BLA circuit inhibition produced a nonspecific impairment in reversal learning (Schoenbaum *et al*, 2007) or in the processing of conditioned reinforcement (Roberts *et al*, 2007) because optogenetic manipulation failed




Figure 5 Effects of optogenetic manipulation on neuronal membrane potential and evoked action potential firing in BLA afferents to the IOFC. (a) Overlay of current-clamp recording of BLA neuronal membrane potential in response to three consecutive light pulses (560 nm, 5 s, shaded area). (b) Mean change in membrane potential (\pm SEM) in NpHR-mCherry-expressing neurons ($n=5$) at the peak and plateau of the light response. (c, d) Representative recordings of action potential firing evoked by current injection steps in the absence (c) and presence (d) of a light pulse (560 nm, 1 s, shaded area). (e) Current injection-evoked firing rates (mean \pm SEM, normalized to maximum rate in 'Light OFF' condition) in NpHR-mCherry-expressing BLA neurons in the presence and absence of a light pulse (560 nm, 1 s). Symbols represent difference relative to baseline membrane potential ($\#$, t -tests, $P<0.05$), relative to the Light OFF condition ($*$, ANOVA light main effect, $P<0.05$), and relative to current steps 0–75 pA (\wedge , Tukey's tests, $P<0.05$).

to alter food CS-induced food-seeking behavior (Supplementary Figure S4). Therefore, inhibition of IOFC afferents to the BLA selectively altered the motivational effects of the cocaine-associated CS.

As IOFC projection neurons are glutamatergic, our findings seem inconsistent with an earlier report that glutamate receptor antagonism in the BLA failed to inhibit CS-induced cocaine-seeking behavior (See *et al*, 2001). Null effects in that pharmacological study may have resulted from insufficient dosing in the BLA. Alternatively, CS-induced reinstatement may require metabotropic glutamate receptor stimulation within the BLA, similar to cocaine-conditioned place preference (Krishnan *et al*, 2011). Overall, the present findings suggest that the IOFC and BLA interact, and specifically the IOFC provides critical input to the BLA via monosynaptic projections, to stimulate CS-induced cocaine-seeking behavior.

The Role of the BLA in CS-Induced Cocaine-Seeking Behavior

The importance of the BLA in CS-induced motivated behavior is well recognized, yet optogenetic inhibition of BLA afferents to the IOFC failed to alter CS-induced cocaine-seeking behavior in the present study (Figure 4b). This unexpected finding may have resulted from possible off-target effects of our optogenetic manipulation (see review by Otchy *et al*, 2015). Mitigating this possibility, previous localization studies that utilized NMDA lesions, or tetrodotoxin-induced or lidocaine-induced neural inactivation, targeted BLA neurons in general (Grimm and See, 2000; Kantak *et al*, 2002; Meil and See, 1997; Whitelaw *et al*, 1996). Some of these manipulations inhibited fibers of passage and axon terminals, including IOFC terminals within the BLA, which would have impaired CS-induced cocaine-seeking behavior based on the findings reported here. Furthermore, earlier optogenetic studies targeted BLA neuronal populations that project to the nucleus accumbens core (NACC) or prelimbic cortex (PrL) and have requisite roles in CS-induced reinstatement (Stefanik and Kalivas, 2013). One population of BLA neurons that projects to the PrL and/or NACC also sends collaterals to the agranular insular cortex (Shinonaga *et al*, 1994), a region adjacent to the IOFC. In comparison, it is unclear whether BLA neurons that project to the IOFC possess direct connections with the PrL or NACC as we did not observe mCherry expression in the IOFC or these putative terminal regions (Supplementary Figure S1E and F) likely because mCherry expression in terminal regions requires more than 4–5 weeks to develop. However, our findings indicate that the activity of the BLA \rightarrow IOFC circuit is not functionally required for CS-induced reinstatement, independent of whether component BLA neurons are anatomically connected with the PrL or NACC.

Further supporting this conclusion, the null effect observed following optogenetic inhibition of BLA afferents to the IOFC was not due to insufficient opsin expression as the number of mCherry expressing cells was similar in the BLA and IOFC (Supplementary Figure S2G). Furthermore, halorhodopsin-expressing cell bodies in the BLA exhibited reliable light-induced hyperpolarization (Figure 5a and b) and inhibition of evoked action potential firing (Figure 5c–e). Termination of optogenetic inhibition did not

generally induce rebound excitation (Figure 5a; Raimondo *et al*, 2012). Finally, our behavioral findings conclusively demonstrated that the NpHR-mCherry and mCherry groups did not differ in response rate immediately after CS-coupled laser stimulation was discontinued (Figure 4d). Accordingly, rebound in neural activity was not responsible for the null effect in cocaine-seeking behavior. Based on these considerations, we conclude that requisite monosynaptic communication between the IOFC and BLA is unidirectional, and the contribution of the BLA → IOFC circuit to CS-induced reinstatement is not critical. This does not rule out that the BLA → IOFC circuit may make nonessential contributions that are masked by intact information transfer from the BLA to the IOFC through multi-synaptic, parallel circuits. Overall, direct BLA input to brain regions *other than the IOFC* is sufficient (present study) and necessary (eg, NAcC and PrL; Stefanik and Kalivas, 2013) for CS-induced reinstatement.

Selective Optogenetic Inhibition of CS-Induced Motivated Behavior

A unique methodological feature of the present study was optogenetic inhibition coupled with discrete CS presentation. In comparison, pharmacological manipulations inhibit neural activity for extended time periods, and thus possibly disrupt the cognitive representation or processing of the drug-paired CS and context alike. While contextual background stimuli acquire more modest motivational effects than CSs due to weaker temporal contiguity with the effects of cocaine, contextual stimuli facilitate drug-seeking behavior (Fuchs *et al*, 2008). For instance, response-contingent presentation of an alcohol-associated CS elicits more drug-seeking behavior in an alcohol-associated context than in a control context (Remedios *et al*, 2014). Following extinction training, such motivational effects of the context may be renewed by CS presentation. In the present study, optogenetic inhibition did not alter cocaine-paired context-induced motivation as the NpHR-mCherry and mCherry groups did not differ in responding during the timeout period, when CS presentations were unavailable, and re-obtained the extinction criterion at similar rates after a Laser ON test (see Supplementary Results). Thus, the robust effect of CS-coupled optogenetic inhibition highlights the degree to which IOFC afferents to the BLA control the motivational effects of cocaine-paired CSs in particular. To further explore the putative contributions of this neural circuit to the motivational effects of the drug-paired context during CS-induced reinstatement, a control group receiving laser stimulation exclusively unpaired from CS presentation will be a useful addition in future studies.

Combinatorial Viral Approach

Another novel methodological component of the present study was the use of a combinatorial viral approach. Retro-Cre-GFP was packaged in AAV serotype 6 which exhibits efficient retrograde transport via mechanisms that are likely dependent on dynein and kinesin-2 proteins (Murlidharan *et al*, 2014; Rothermel *et al*, 2013; Salegio *et al*, 2013). Similar studies have used recombinant canine

adenovirus 2 (CAV2) or AAV serotype 5 to manipulate neural circuits (Beyeler *et al*, 2016; Marchant *et al*, 2015). Most commonly, studies have utilized methodology that involves optogenetic inhibition at the virus infusion site (to target discrete brain nuclei) or at the terminal region of the target circuit following virus injection at the cell body region, even though these approaches are more prone to affect fibers of passage (Beyeler *et al*, 2016; Stamatakis and Stuber, 2012). In particular, the combinatorial viral approach is essential (a) to manipulate proteins that are not expressed at the terminal region of target circuits and (b) to mitigate the current scarcity of Cre-driver lines and unavailability of Cre-dependent opsins packaged in retrogradely transported viruses (Saunders *et al*, 2015; Witten *et al*, 2011; Warden *et al*, 2014). One caveat is that retrograde transport efficiency in various AAV serotypes appears to vary across species (Aschauer *et al*, 2013; Rothermel *et al*, 2013). The current viral approach is challenging as it requires co-infection of neurons by two viruses, but it can also be employed to selectively manipulate subpopulations of neurons using unique promoter combinations. Therefore, this combinatorial viral approach is an important tool for circuit-specific manipulations.

Conclusions: The Position of the IOFC and BLA in the CS-induced Relapse Circuitry

The present findings refine our understanding of the neural circuitry of cocaine-seeking behavior. According to previous studies, optogenetic inhibition of PrL to NAcC, BLA to PrL, or BLA to NAcC projections disrupts CS-induced reinstatement (Stefanik and Kalivas, 2013; Stefanik *et al*, 2013b), while optogenetic inhibition of NAcC to dorsolateral ventral pallidum (dlVP) or VTA to NAcC projections impairs CS + cocaine-primed reinstatement of cocaine-seeking behavior (Stefanik *et al*, 2013a). Other studies indicate that muscarinic cholinergic and dopamine receptor stimulation in the BLA mediates CS-induced reinstatement (See *et al*, 2001, 2003). Together these and the present findings are consistent with a circuitry model in which the BLA assesses the motivational significance of cocaine-paired CSs at the time of testing (Fuchs *et al*, 2002) based on input from the IOFC and VTA (See *et al*, 2001). The BLA may transmit the results of these computations to the PrL, which holds executive function (Lasseter *et al*, 2010) and integrates this information with input from the VTA (Ciccocioppo *et al*, 2001). Subsequently, the PrL mediates response selection and initiation via the NAcC-dlVP circuit (Kalivas and McFarland, 2003). However, response execution at the level of the NAcC occurs with additional requisite input from the BLA and VTA, and this may permit more nuanced control over behavior in complex situations where multiple reinforcers and response opportunities exist.

FUNDING AND DISCLOSURE

This work was supported by NIDA grants R01 DA025646 (RAF), R01 DA032750 (GDS), and K99 DA037271 (AAA), and NIAAA grant R01 AA012439 (DJR). The authors declare no conflict of interest.

ACKNOWLEDGMENTS

We would like to thank Carey M Lyons, Jessica A Higginbotham, Randall L Ung, and Pranish A Kantak for technical assistance, Dr JrGang Cheng and Xinghua Zeng for cloning and packing the cre-recombinase virus, and Drs Anjen Chenn and Karl Deisseroth for viral constructs.

REFERENCES

- Aschauer DF, Kreuz S, Rumpel S (2013). Analysis of transduction efficiency, tropism and axonal transport of aav serotypes 1, 2, 5, 6, 8 and 9 in the mouse brain. *PLoS One* **8**: e76310.
- Beyeler A, Namburi P, Glober GF, Simonnet C, Calhoon GG, Conyers GF et al (2016). Divergent routing of positive and negative information from the amygdala during memory retrieval. *Neuron* **16**: 00183–00185.
- Bossert JM, Marchant NJ, Calu DJ, Shaham Y (2013). The reinstatement model of drug relapse: recent neurobiological findings, emerging research topics, and translational research. *Psychopharmacology (Berl)* **229**: 453–476.
- Brebner K, Childress AR, Roberts DCS (2002). A potential role for GABA(B) agonists in the treatment of psychostimulant addiction. *Alcohol Alcohol* **37**: 478–484.
- Calu DJ, Kawa AB, Marchant NJ, Navarre BM, Henderson MJ, Chen B et al (2013). Optogenetic inhibition of dorsal medial prefrontal cortex attenuates stress-induced reinstatement of palatable food seeking in female rats. *J Neurosci* **33**: 214–226.
- Carlezon WAJ, Boundy VA, Haile CN, Lane SB, Kalb RG, Neve RL et al (1997). Sensitization to morphine induced by viral-mediated gene transfer. *Science* **277**: 812–814.
- Childress AR, Mozley PD, McElgin W, Fitzgerald J, Reivich M, O'Brien CP (1999). Limbic activation during cue-induced cocaine craving. *Am J Psychiatry* **156**: 11–18.
- Ciccocioppo R, Sanna PP, Weiss F (2001). Cocaine-predictive stimulus induced drug-seeking behavior and neural activation in limbic brain regions after multiple months of abstinence: reversal by D(1) antagonists. *Proc Natl Acad Sci USA* **98**: 1976–1981.
- Ehrman RN, Robbins SJ, Childress AR, O'Brien CP (1992). Conditioned responses to cocaine-related stimuli in cocaine abuse patients. *Psychopharmacology (Berl)* **107**: 523–529.
- Fuchs RA, Eaddy JL, Su Z-I, Bell GH (2007). Interactions of the basolateral amygdala with the dorsal hippocampus and dorsomedial prefrontal cortex regulate drug context-induced reinstatement of cocaine-seeking in rats. *Eur J Neurosci* **26**: 487–498.
- Fuchs RA, Lasseter HC, Ramirez DR, Xie X (2008). Relapse to drug seeking following prolonged abstinence: the role of environmental stimuli. *Drug Discov Today Dis Models* **5**: 251–258.
- Fuchs RA, Evans KA, Parker MP, See RE (2004). Differential involvement of orbitofrontal cortex subregions in conditioned cue-induced and cocaine-primed reinstatement of cocaine seeking in rats. *J Neurosci* **24**: 6600–6610.
- Fuchs RA, Weber SM, Rice HJ, Neisewander JL (2002). Effects of excitotoxic lesions of the basolateral amygdala on cocaine-seeking behavior and cocaine conditioned place preference in rats. *Brain Res* **929**: 15–25.
- Goldstein RZ, Volkow ND (2002). Drug addiction and its underlying neurobiological basis: neuroimaging evidence for the involvement of the frontal cortex. *Am J Psychiatry* **159**: 1642–1652.
- Grimm JW, See RE (2000). Dissociation of primary and secondary reward-relevant limbic nuclei in an animal model of relapse. *Neuropsychopharmacology* **22**: 473–479.
- Kalivas PW, McFarland K (2003). Brain circuitry and the reinstatement of cocaine-seeking behavior. *Psychopharmacology (Berl)* **168**: 44–56.
- Kantak KM, Black Y, Valencia E, Green-Jordan K, Eichenbaum HB (2002). Dissociable effects of lidocaine inactivation of the rostral and caudal basolateral amygdala on the maintenance and reinstatement of cocaine-seeking behavior in rats. *J Neurosci* **22**: 1126–1136.
- Krishnan B, Genzer KM, Pollandt SW, Liu J, Gallagher JP, Shinnick-Gallagher P (2011). Dopamine-induced plasticity, phospholipase D (PLD) activity and cocaine-cue behavior depend on PLD-linked metabotropic glutamate receptors in amygdala. *PLoS One* **6**: e25639.
- Lasseter HC, Wells AM, Xie X, Fuchs RA (2011). Interaction of the basolateral amygdala and orbitofrontal cortex is critical for drug context-induced reinstatement of cocaine-seeking behavior in rats. *Neuropsychopharmacology* **36**: 711–720.
- Lasseter HC, Xie X, Arguello AA, Wells AM, Hodges MA, Fuchs RA (2014). Contribution of a mesocorticolimbic subcircuit to drug context-induced reinstatement of cocaine-seeking behavior in rats. *Neuropsychopharmacology* **39**: 660–669.
- Lasseter HC, Xie X, Ramirez DR, Fuchs RA (2010). Prefrontal cortical regulation of drug seeking in animal models of drug relapse. *Curr Top Behav Neurosci* **3**: 101–117.
- Marchant NJ, Whitaker LR, Bossert JM, Harvey BK, Hope BT, Kaganovsky K et al (2015). Behavioral and physiological effects of a novel kappa opioid receptor based DREADD in rats. *Neuropsychopharmacology* **41**: 402–409.
- Meil WM, See RE (1997). Lesions of the basolateral amygdala abolish the ability of drug associated cues to reinstate responding during withdrawal from self-administered cocaine. *Behav Brain Res* **87**: 139–148.
- Murlidharan G, Samulski RJ, Asokan A (2014). Biology of adeno-associated viral vectors in the central nervous system. *Front Mol Neurosci* **7**: 76.
- Otchy TM, Wolff SB, Rhee JY, Pehlevan C, Kawai R, Kempf A et al (2015). Acute off-target effects of neural circuit manipulations. *Nature* **528**: 358–363.
- Paxinos G, Watson C. *The Rat Brain in Stereotaxic Coordinates, Compact* 3rd edn. Academic Press: San Diego, CA, USA, (1997).
- Penrod RD, Wells AM, Carlezon WAJ, Cowan CW (2015). Use of adeno-associated and herpes simplex viral vectors for *in vivo* neuronal expression in mice. *Curr Protoc in Neurosci* **73**: 1–31.
- Price JL (2007). Definition of the orbital cortex in relation to specific connections with limbic and visceral structures and other cortical regions. *Ann N Y Acad Sci* **1121**: 54–71.
- Raimondo JV, Kay L, Ellender TJ, Akerman CJ (2012). Optogenetic silencing strategies differ in their effect on inhibitory synaptic transmission. *Nat Neurosci* **15**: 1102–1104.
- Remedios J, Woods C, Tardif C, Janak PH, Chaudhri N (2014). Pavlovian-conditioned alcohol-seeking behavior in rats is invigorated by the interaction between discrete and contextual alcohol cues: implications for relapse. *Brain Behav* **4**: 278–289.
- Roberts AC, Reekie Y, Braesicke K (2007). Synergistic and regulatory effects of orbitofrontal cortex on amygdala-dependent appetitive behavior. *Ann NY Acad Sci* **1121**: 297–319.
- Rothermel M, Brunert D, Zabawa C, Diaz-Quesada M, Wachowiak M (2013). Transgene expression in target-defined neuron populations mediated by retrograde infection with adeno-associated viral vectors. *J Neurosci* **33**: 15195–15206.
- Saddoris MP, Gallagher M, Schoenbaum G (2005). Rapid associative encoding in basolateral amygdala depends on connections with orbitofrontal cortex. *Neuron* **46**: 321–331.
- Salegio EA, Samaranch L, Kells AP, Mittermeyer G, San Sebastian W, Zhou S et al (2013). Axonal transport of adeno-associated viral vectors is serotype-dependent. *Gene Ther* **20**: 348–352.
- San Sebastian W, Samaranch L, Heller G, Kells AP, Bringas J, Pivrotto P et al (2013). Adeno-associated virus type 6 is retrogradely transported in the non-human primate brain. *Gene Ther* **20**: 1178–1183.
- Saunders BT, Richard JM, Janak PH (2015). Contemporary approaches to neural circuit manipulation and mapping: focus

- on reward and addiction. *Philos Trans R Soc Lond B Biol Sci* **370**: 20140210.
- Schoenbaum G, Saddoris MP, Stalnaker TA (2007). Reconciling the roles of orbitofrontal cortex in reversal learning and the encoding of outcome expectancies. *Ann NY Acad Sci* **1121**: 320–335.
- See RE, McLaughlin J, Fuchs RA (2003). Muscarinic receptor antagonism in the basolateral amygdala blocks acquisition of cocaine-stimulus association in a model of relapse to cocaine-seeking behavior in rats. *Neuroscience* **117**: 477–483.
- See RE, Kruzich PJ, Grimm JW (2001). Dopamine, but not glutamate, receptor blockade in the basolateral amygdala attenuates conditioned reward in a rat model of relapse to cocaine-seeking behavior. *Psychopharmacology (Berl)* **154**: 301–310.
- Shinonaga Y, Takada M, Mizuno N (1994). Topographic organization of collateral projections from the basolateral amygdaloid nucleus to both the prefrontal cortex and nucleus accumbens in the rat. *Neuroscience* **58**: 389–397.
- Sparta DR, Stamatakis AM, Phillips JL, Hovelsø N, Zessen R van, Stuber GD (2012). Construction of implantable optical fibers for long-term optogenetic manipulation of neural circuits. *Nat Protoc* **7**: 12–23.
- Stamatakis AM, Stuber GD (2012). Optogenetic strategies to dissect the neural circuits that underlie reward and addiction. *Cold Spring Harb Perspect Med* **2**: pii011924.
- Stefanik MT, Kalivas PW (2013). Optogenetic dissection of basolateral amygdala projections during cue-induced reinstatement of cocaine seeking. *Front Behav Neurosci* **7**: 213.
- Stefanik MT, Kupchik YM, Brown RM, Kalivas PW (2013a). Optogenetic evidence that pallidal projections, not nigral projections, from the nucleus accumbens core are necessary for reinstating cocaine seeking. *J Neurosci* **33**: 13654–13662.
- Stefanik MT, Moussawi K, Kupchik YM, Smith KC, Miller RL, Huff ML et al (2013b). Optogenetic inhibition of cocaine seeking in rats. *Addict Biol* **18**: 50–53.
- Warden MR, Cardin JA, Deisseroth K (2014). Optical neural interfaces. *Annu Rev Biomed Eng* **16**: 103–129.
- Whitelaw RB, Markou A, Robbins TW, Everitt BJ (1996). Excitotoxic lesions of the basolateral amygdala impair the acquisition of cocaine-seeking behaviour under a second-order schedule of reinforcement. *Psychopharmacology (Berl)* **127**: 213–224.
- Witten IB, Steinberg EE, Lee SY, Davidson TJ, Zalocusky KA, Brodsky M et al (2011). Recombinase-driver rat lines: tools, techniques, and optogenetic application to dopamine-mediated reinforcement. *Neuron* **72**: 721–733.
- Young KA, Franklin TR, Roberts DCS, Jagannathan K, Suh JJ, Wetherill RR et al (2014). Nipping cue reactivity in the bud: baclofen prevents limbic activation elicited by subliminal drug cues. *J Neurosci* **34**: 5038–5043.
- Zeeb FD, Winstanley CA (2013). Functional disconnection of the orbitofrontal cortex and basolateral amygdala impairs acquisition of a rat gambling task and disrupts animals' ability to alter decision-making behavior after reinforcer devaluation. *J Neurosci* **33**: 6434–6443.

Supplementary Information accompanies the paper on the Neuropsychopharmacology website (<http://www.nature.com/npp>)