

# A Pharmacogenetic ‘Restriction-of-Function’ Approach Reveals Evidence for Anxiolytic-Like Actions Mediated by $\alpha 5$ -Containing GABA<sub>A</sub> Receptors in Mice

Lauren M Behlke<sup>1,2,6</sup>, Rachel A Foster<sup>1,2,6</sup>, Jing Liu<sup>1,2,6</sup>, Dietmar Benke<sup>3</sup>, Rebecca S Benham<sup>1,2</sup>, Anna J Nathanson<sup>1,2</sup>, Benjamin K Yee<sup>4</sup>, Hanns Ulrich Zeilhofer<sup>3,5</sup>, Elif Engin<sup>1,2</sup> and Uwe Rudolph<sup>\*,1,2</sup>

<sup>1</sup>Laboratory of Genetic Neuropharmacology, McLean Hospital, Belmont, MA, USA; <sup>2</sup>Department of Psychiatry, Harvard Medical School, Boston, MA, USA; <sup>3</sup>Institute of Pharmacology and Toxicology, University of Zurich, Zurich, Switzerland; <sup>4</sup>Department of Rehabilitation Sciences, The Hong Kong Polytechnic University, Hong Kong, Hong Kong; <sup>5</sup>Institute of Pharmaceutical Sciences, Swiss Federal Institute of Technology (ETH) Zurich, Zürich, Switzerland

Benzodiazepines have been widely used for their anxiolytic actions. However, the contribution of GABA<sub>A</sub> receptor subtypes to anxiolysis is still controversial. Studies with mutant mice harboring diazepam-insensitive  $\alpha$ -subunits  $\alpha 1$ ,  $\alpha 2$ ,  $\alpha 3$ , or  $\alpha 5$  have revealed that  $\alpha 2$ -containing GABA<sub>A</sub> receptors ( $\alpha 2$ -GABA<sub>A</sub>Rs) are required for diazepam-induced anxiolysis, with no evidence for an involvement of any other  $\alpha$ -subunit, whereas TP003, described as a selective modulator of  $\alpha 3$ -containing GABA<sub>A</sub> receptors, was shown to be anxiolytic. Here, we describe a novel, systematic approach to evaluate the role of positive allosteric modulation of each of the four diazepam-sensitive  $\alpha$ -subtypes in anxiety-related behavioral paradigms. By combining H to R point mutations in three out of the four diazepam-sensitive  $\alpha$ -subunits in mice with a 129X1/SvJ background, diazepam becomes a subtype-specific modulator of the remaining non-mutated  $\alpha$ -subtype. Modulation of  $\alpha 5$ -GABA<sub>A</sub>Rs, but not of  $\alpha 2$ -GABA<sub>A</sub>Rs, increased the time in the light side of the light–dark box as well as open-arm exploration in the elevated plus maze. In contrast, modulation of  $\alpha 3$ -GABA<sub>A</sub>Rs decreased open-arm exploration, whereas modulation of  $\alpha 2$ -GABA<sub>A</sub>Rs increased time in the center in the open-field test. Modulation of any single  $\alpha$ -subtype had no effect on stress-induced hyperthermia. Our results provide evidence that modulation of  $\alpha 5$ -GABA<sub>A</sub>Rs elicits anxiolytic-like actions, whereas our data do not provide evidence for an anxiolytic-like action of  $\alpha 3$ -GABA<sub>A</sub>Rs. Thus,  $\alpha 5$ -GABA<sub>A</sub>Rs may be suitable targets for novel anxiolytic drugs.

*Neuropsychopharmacology* (2016) **41**, 2492–2501; doi:10.1038/npp.2016.49; published online 11 May 2016

## INTRODUCTION

Benzodiazepines—pharmacological agents whose primary mechanism of action is the positive allosteric modulation of GABA<sub>A</sub> receptors—have been a major pharmacotherapeutic tool to manage pathological anxiety in the clinic for decades (Shader and Greenblatt, 1993). Although highly effective in reducing anxiety symptoms, the long-term use of benzodiazepines has been fraught with an unfavorable side-effect profile (eg, sedation and dependence liability) because of their nonselective action via multiple GABA<sub>A</sub> receptor subtypes. GABA<sub>A</sub> receptors are heteropentamers that can be configured from a repertoire of at least 19 subunits and are often classified into GABA<sub>A</sub> receptor subtypes based on their  $\alpha$ -subunit ( $\alpha 1$ – $\alpha 6$ ), which is an essential component of the benzodiazepine binding site (Rudolph and Knoflach, 2011).

Classical benzodiazepines such as diazepam bind to and modulate four out of these six subtypes ( $\alpha 1$ -,  $\alpha 2$ -,  $\alpha 3$ -, or  $\alpha 5$ -containing GABA<sub>A</sub> receptors, which are from here on referred to as  $\alpha 1$ -,  $\alpha 2$ -,  $\alpha 3$ -, or  $\alpha 5$ -GABA<sub>A</sub>Rs), leading to both therapeutically desired and undesired effects.

The pharmacological functions of GABA<sub>A</sub> receptor subtypes have so far been deduced from studies using subtype-selective compounds, GABA<sub>A</sub> receptor subunit knockout mice, and GABA<sub>A</sub> receptor subunit knock-in mice with mutations preventing binding of benzodiazepines while leaving the sensitivity for the physiological neurotransmitter GABA intact (Rudolph and Knoflach, 2011). Studies with  $\alpha$ -subunit knock-in mice with diazepam-insensitive  $\alpha$ -subunits, that is,  $\alpha 1$ (H101R),  $\alpha 2$ (H101R),  $\alpha 3$ (H126R), and  $\alpha 5$ (H105R) mice, demonstrated that the modulation of  $\alpha 2$ -GABA<sub>A</sub>Rs, but not of  $\alpha 1$ -,  $\alpha 3$ -, or  $\alpha 5$ -GABA<sub>A</sub>Rs, is required for the anxiolytic-like effects of benzodiazepines (Crestani *et al*, 2002; Low *et al*, 2000; McKernan *et al*, 2000; Morris *et al*, 2006; Rudolph *et al*, 1999; Smith *et al*, 2012; see also Engin *et al*, 2016). In an apparent contrast, the compound TP003 (4,2'-difluoro-5'-[8-fluoro-7-(1-hydroxy-1-methylethyl)imidazo[1,2-*a'*]pyridin-3-yl]biphenyl-2-carbonitrile), reported to be selective for  $\alpha 3$ -GABA<sub>A</sub>Rs in *in vitro* assays on recombinant receptors,

\*Correspondence: Dr U Rudolph, Laboratory of Genetic Neuropharmacology, McLean Hospital, Mailstop No. 145, 115 Mill Street, Belmont, MA 02478-1064, USA, Tel: +1 617 855 2088, Fax: +1 617 855 2012, E-mail: urudolph@mclean.harvard.edu

<sup>6</sup>These authors contributed equally to this work.

Received 30 December 2015; revised 1 April 2016; accepted 6 April 2016; accepted article preview online 12 April 2016

was found to have anxiolytic-like effects in several species (Dias *et al*, 2005), which led to the widely accepted conclusion that  $\alpha 3$ -GABA<sub>A</sub>Rs may also mediate anxiolytic effects of benzodiazepines (eg, Korpi *et al*, 2015; Sigel and Steinmann, 2012). However, two recent studies were unable to reproduce the  $\alpha 3$ -selectivity of TP003 in recombinant receptors (Christian *et al*, 2015; de Lucas *et al*, 2015), raising the question whether the anxiolytic-like action of TP003 is really dependent on modulation of  $\alpha 3$ -GABA<sub>A</sub>Rs, and, on a broader scale, whether  $\alpha 3$ -GABA<sub>A</sub>Rs are involved at all in the modulation of anxiety-related behaviors. Furthermore, a conditional deletion of the  $\alpha 5$ -subunit in PKC $\delta$ + neurons in the central amygdala resulted in increased anxiety in the open-field (OF) and elevated plus maze (EPM) tests (Botta *et al*, 2015), suggesting a role for  $\alpha 5$ -GABA<sub>A</sub>Rs in anxiety regulation. However, it has not been examined whether systemic positive allosteric modulation of  $\alpha 5$ -GABA<sub>A</sub>Rs leads to anxiolysis.

Thus, the role of different GABA<sub>A</sub> receptor subtypes in anxiety and anxiolytic drug action is still unclear, and the findings from studies using pharmacological agents have been inconclusive, as the currently available 'subtype-selective' compounds are not truly specific for a given  $\alpha$ -subtype. The current study is designed to evaluate whether highly specific positive allosteric modulation of any individual GABA<sub>A</sub> receptor subtype is sufficient to elicit anxiolytic-like actions using a combined pharmacological and genetic engineering approach, which overcomes the selectivity limitations of earlier pharmacological studies.

## MATERIALS AND METHODS

### Animals

All experiments and procedures were approved by Kantonales Veterinaramt Zurich or the McLean Hospital Institutional Animal Care and Use Committee following guidelines in the National Research Council Guide for the Care and Use of Laboratory Animals (8th edition). Mice (bred on the 129X1/SvJ background) carry three out of the four point mutations  $\alpha 1$ (H101R),  $\alpha 2$ (H101R),  $\alpha 3$ (H126R), and  $\alpha 5$ (H105R), which render the respective GABA<sub>A</sub> receptors insensitive to modulation by diazepam (Crestani *et al*, 2002; Low *et al*, 2000; Rudolph *et al*, 1999). We refer to these mice as DZ $\alpha 1$ , DZ $\alpha 2$ , DZ $\alpha 3$  and DZ $\alpha 5$  based on which of these  $\alpha$ -subunits is not mutated and thus diazepam-sensitive (previously described as HRRR, RHRR, RRHR and RRRH, respectively, in Ralvenius *et al*, 2015). Subjects were male mice aged 8–16 weeks at the time of testing and group-housed with 2–5 mice per cage in Super Mouse 750 cages. Cohort 1 mice were transferred from our breeding colony (12 : 12-h light–dark cycle with lights on at 0700 hours) to a housing room in our behavioral suite where they were kept on a regular 12 : 12-h light–dark cycle (lights on at 0700 hours). Following the stress-induced hyperthermia (SIH) test, the mice were moved to a different housing room where they were kept on a 12 : 12-h reverse light–dark cycle (lights on at 1900 hours) for 3 weeks, after which they were tested in the light–dark box (LDB). In the LDB, animals were balanced for previous drug history. Cohort 2 was moved to the reverse light–dark cycle housing room and were habituated to this cycle for 3 weeks before they were tested in the EPM. Cohort 3 was habituated to the reverse

light–dark cycle for at least 3 weeks before undergoing novel OF, forced-swim test (FST), and tail suspension test (TST), with a 1-week hiatus between the tests. The order of tests was counterbalanced for different groups of mice. A fourth cohort was habituated in our regular light–dark cycle behavioral suite for 1 week before undergoing the conditioned place preference (CPP) paradigm.

### Drugs

Diazepam (Sigma-Aldrich, St Louis, MO) was dissolved in 10% 2-hydroxypropyl- $\beta$ -cyclodextrin (Sigma-Aldrich). Diazepam (3 mg/kg) and vehicle were administered in a volume of 10 ml/kg via PO injection 1 h before testing (SIH) or intraperitoneal injection (2 or 10 mg/kg) 30 min before behavioral testing (LDB, EPM, OF, FST, TST). Cocaine (Sigma-Aldrich) was dissolved in 0.9% saline and administered intraperitoneally immediately before conditioning at 20 mg/kg.

### Autoradiography

Mice were deeply anesthetized with isoflurane and killed by decapitation. The dissected brains were immediately frozen in powdered dry ice and stored at  $-80^{\circ}\text{C}$  until used. Parasagittal cryostat-cut sections (16  $\mu\text{m}$ ) were thawed, washed two times for 10 min at  $4^{\circ}\text{C}$  in 50 mM Tris-HCl (pH 7.4), and dried under a stream of cold air. The sections were incubated for 90 min at  $4^{\circ}\text{C}$  with 5 nM [ $^3\text{H}$ ]flumazenil (50 Ci/mmol; ANAWA Trading SA, Wangen, Switzerland) or 8.8 nM [ $^3\text{H}$ ]Ro15-4513 (22.7 Ci/mmol; Perkin-Elmer, Schwerzenbach, Switzerland) diluted in 50 mM Tris-HCl (pH 7.4). Nonspecific [ $^3\text{H}$ ]flumazenil or [ $^3\text{H}$ ]Ro15-4513 binding was determined in the presence of 10  $\mu\text{M}$  clonazepam and 10  $\mu\text{M}$  flumazenil, respectively. The incubation was terminated by washing the sections three times for 20 s in ice-cold 50 mM Tris-HCl (pH 7.4). The sections were thoroughly dried and exposed to a tritium-sensitive phosphor imaging screen along with [ $^3\text{H}$ ]-standards for 5–7 days. The screens were scanned with a Packard Cyclone Storage Phosphor imager and images were quantified using Opti-Quant (version 4.0). Sections from all genotypes were processed and exposed in parallel and are therefore directly comparable.

### Behavioral Experiments

**Novel OF.** The OF apparatus was a transparent Plexiglas box (42 cm  $\times$  42 cm  $\times$  31 cm), evenly illuminated at 100 lx. Mice were placed in one corner and allowed to explore freely for 30 min while their distance traveled (cm) was recorded and analyzed using EthoVision XT video tracking system (Noldus Information Technology, Wageningen, The Netherlands). The center zone was 20 cm  $\times$  20 cm. The 408 s (22.7% of the time) in the center zone was determined as the chance level based on the sizes of the center zone and the full chamber assuming purely probabilistic movement of the mice.

**Elevated plus maze.** The EPM apparatus was elevated 1 m above the floor and consisted of two open arms (35 cm long  $\times$  6 cm wide), two closed arms (35 cm long  $\times$  6 cm

wide × 20 cm high), and one center area (5 cm × 5 cm). When all mice were tested under the same lighting conditions (Supplementary Figure S1), the vehicle-treated wild-type (WT) mice appeared to spend more time in the open arms than in any of the vehicle-treated mutant mice. Although this trend did not reach statistical significance in this pilot experiment, we predicted it could become a significant confound in a complete data set. Thus, each genotype was tested under conditions that resulted in a percent time in light that would avoid any floor or ceiling effects (≈20%) after vehicle injection. All mice except WT were habituated to the testing room overnight, and all mice including WT were habituated to the appropriate lighting in the testing room for an hour before testing. Illumination in the open arms was 20 lx for WT mice, 10 lx for DZα1, DZα3, and DZα5 mice, and red light for DZα2 mice. For additional reduction of baseline anxiety in DZα1, DZα2, DZα3, and DZα5 mice, each mouse was placed in a new cage after completion of the behavioral test to separate naïve mice from those who had experienced the EPM test. Each mouse was placed in the maze facing an open arm and allowed to freely explore the maze for 5 min. Behavioral activity was recorded with the EthoVision XT video tracking system. After each trial, the maze was wiped down with 70% ethanol and allowed to dry completely. The percent time in open arms ((time in open arms/5 min) × 100) and the percent open arm entries ((open arm entries)/(open arm entries + closed arm entries) × 100) were calculated as measures of anxiolysis. In addition, total distance traveled in the maze during the test was used to determine the effect of diazepam on locomotion during EPM. The number of total arm entries was recorded as an additional measure of within-test locomotor activity.

**Light–dark box test.** The LDB apparatus consisted of one clear, brightly lit (250 lx) chamber (28 cm × 28 cm × 31 cm) and one smaller dark (<10 lx) chamber (14 cm × 14 cm × 31 cm) connected by a square opening between the chambers (5 cm × 5 cm). As the lit chamber is two times as large as the dark chamber, mice would spend 67% time in the lit chamber if the distribution of time between the chambers was random ('chance level'). All mice were habituated to the test room overnight and habituated to the appropriate ambient lighting an hour before testing. At the start of the test, each mouse was placed within the dark chamber and allowed to freely explore the two chambers for 6 min. Each mouse's activity in the visible clear chamber was recorded with the EthoVision XT video tracking system. Between each trial, the chambers were wiped down with 70% ethanol and allowed to dry completely. The percent time in light ((time in clear chamber)/6 min × 100) was calculated as a measure of anxiolysis.

### Stress-Induced Hyperthermia

Mice were housed in the room where the SIH test occurred for 1 week before the test, and were single-housed 24 h before the start of the test. Diazepam was administered 1 h before first rectal temperature reading (*T*<sub>1</sub>). The second temperature (*T*<sub>2</sub>) was taken 10 min after *T*<sub>1</sub>. The change in temperature (*T*<sub>2</sub> – *T*<sub>1</sub>) induced by stress of *T*<sub>1</sub> was calculated and used to measure the autonomic response to stress.

### FST and TST

These tests were performed as described previously (Vollenweider *et al*, 2011). Diazepam or vehicle were administered intraperitoneally 30 min before testing.

### Statistical Analysis

For autoradiography quantification, each subregion was analyzed separately using a one-way analysis of variance (ANOVA). Only the [<sup>3</sup>H]flumazenil binding in the hippocampus data set passed the Shapiro–Wilk normality test, and thus this was the only data set that was analyzed with a parametric ANOVA and *post hoc* Tukey's test. All other quantification data sets were analyzed with a Kruskal–Wallis ANOVA on ranks followed by *post hoc* Dunn's method when necessary. Owing to the between-genotype variance in behavioral responses with vehicle injection during LDB and EPM, statistical analyses were only performed for within-genotype drug effects but not between-genotype effects. For OF, EPM, and LDB, each genotype was analyzed separately using a one-way ANOVA followed by the *post hoc* Tukey's test for pairwise comparison. For data sets that failed the Shapiro–Wilk normality test or the equal variance test (*p* < 0.05), the Kruskal–Wallis one-way ANOVA on ranks was used followed by *post hoc* pairwise comparison using Tukey's test when the treatment group sizes were equal and Dunn's method when the treatment group sizes were unequal. This nonparametric test was used in LDB data for DZα1, DZα2, and DZα3, and in EPM for DZα2 and DZα3 in % time in open arms, DZα5 distance traveled and DZα1 in % time, % entries, distance traveled, and total entries. For OF, the nonparametric analysis was used for WT center zone duration, DZα1 distance travelled and center duration, DZα2 for distance travelled, and DZα5 for center duration. For CPP (see Supplementary Methods), each genotype was analyzed using a two-way repeated-measures ANOVA. For SIH, the drug effect of each genotype was analyzed using separate *t*-tests for each genotype. The DZα3 SIH data set did not pass the Shapiro–Wilk normality test, and thus it was analyzed with a Mann–Whitney rank-sum test. In all statistical analyses, results were considered significant when *p* < 0.05.


## RESULTS

To investigate the pharmacological effects of highly specific positive allosteric modulation of single GABA<sub>A</sub> receptor subtypes, we used four gene-targeted mouse lines in which three out of the four diazepam-sensitive α-subunits were rendered insensitive to modulation by diazepam, which we refer to as DZα1, DZα2, DZα3, or DZα5 mice (the 'x' in 'DZαx' indicates the non-mutated α subunit) (Ralvenius *et al*, 2015).

### Receptor Autoradiography

To visualize the benzodiazepine binding sites in mice carrying triple point mutations and WT mice, we performed receptor autoradiography with [<sup>3</sup>H]flumazenil (Figure 1a). [<sup>3</sup>H]flumazenil binds with high affinity only to the GABA<sub>A</sub> receptors containing the WT α1-, α2-, α3-, or α5-subunits.





**Figure 1** Receptor autoradiography demonstrating specificity of pharmacogenetic 'restriction-of-function' approach. (a) [<sup>3</sup>H]flumazenil binding to parasagittal brain sections derived from wild-type (WT, 129X1/Sv) and triple mutant mice (DZα1, DZα2, DZα3, and DZα5). The α-subunit indicated for triple mutant mice is diazepam-sensitive. Nonspecific [<sup>3</sup>H]flumazenil binding was assessed in the presence of 10 μM clonazepam. Sections from all genotypes were processed in parallel and exposed to the same phosphorimaging screen. Therefore, the gray levels of sections from the different genotypes reflect the abundance of the respective receptor subtype(s) and can be directly compared. Scale bar = 2 mm. (b) Quantification of [<sup>3</sup>H]flumazenil binding in WT and triple mutant mice. [<sup>3</sup>H]flumazenil binding (nCi/mg) is expressed as mean (± SEM); n = 5–6 per genotype. \*p < 0.05, \*\*p < 0.01, and \*\*\*p < 0.001. (c) [<sup>3</sup>H]Ro15-4513 binding to parasagittal brain sections derived from WT (129X1/Sv) and triple mutant mice (DZα1, DZα2, DZα3, and DZα5). Nonspecific [<sup>3</sup>H]Ro15-4513 binding was assessed in the presence of 10 μM flumazenil. Sections from all genotypes were processed in parallel and exposed to the same phosphorimaging screen. Therefore, the gray levels of sections from the different genotypes reflect the abundance of the respective receptor subtype(s) and can be directly compared. Scale bar = 2 mm. (d) Quantification of [<sup>3</sup>H]Ro15-4513 binding in WT and triple mutant mice. [<sup>3</sup>H]Ro15-4513 binding (nCi/mg) is expressed as mean (± SEM); n = 6 per genotype. Cx, cortex; Hi, hippocampus.


The distribution of [<sup>3</sup>H]flumazenil binding observed in the DZα1, DZα2, DZα3, and DZα5 triple mutant mice corresponds well to the distribution of the diazepam-sensitive α-subunits seen in immunohistochemistry (Supplementary Figure S2), demonstrating the specificity of [<sup>3</sup>H]flumazenil binding to the non-mutated α-subtypes in the triple mutant mice. Quantification of [<sup>3</sup>H]flumazenil binding in total brain (H (4) = 26.253, p < 0.001) as well as in hippocampal (F (4, 24) = 72.281, p < 0.001) and cortical subregions (H (4) = 25.726, p < 0.001) further supports the specificity of [<sup>3</sup>H]flumazenil binding in the triple mutant mice (Figure 1b). DZα1 and DZα2 mice only showed a statistically significant decrease in [<sup>3</sup>H]flumazenil binding in the hippocampus (DZα1, p < 0.01; DZα2 p < 0.001), whereas DZα3 and DZα5 mice showed statistically significant decreases in the hippocampus (p < 0.001 for both), cortex (p < 0.05 for both), and total brain (p < 0.05 for both). The sum of [<sup>3</sup>H]flumazenil binding for DZα1, DZα2, DZα3, and

DZα5 mice is more than [<sup>3</sup>H]flumazenil binding in WT mice, which is likely due to GABA<sub>A</sub> receptor complexes that contain two different α-subunits (Benke *et al*, 2004).

To visualize binding to all GABA receptors, we performed additional receptor autoradiography with [<sup>3</sup>H]Ro15-4513 (Figure 1c). The distribution of [<sup>3</sup>H]Ro15-4513 binding in all of the triple mutant mice corresponds well to the [<sup>3</sup>H]Ro15-4513 binding in the WT mice, suggesting that total GABA<sub>A</sub> receptor expression levels remain unchanged in the triple mutant mice. Quantification of [<sup>3</sup>H]Ro15-4513 binding demonstrates no significant difference in total GABA<sub>A</sub> receptor levels between any of the triple mutants and WT (Figure 1d).

#### LDB Test

We assessed the effect of diazepam on unconditioned anxiety in the LDB test (Figure 2). One-way ANOVAs indicated a



**Figure 2** Light-dark box (LDB) test. Percent time in the light during LDB test after an injection of vehicle or diazepam (2 and 10 mg/kg) in wild-type (WT) (129X1/Sv) and triple mutant mice. The dotted line ('chance level') indicates the percent time in the light chamber that would be expected if the distribution of time between the chambers was random (67%). Percent time in the light side is expressed as mean ( $\pm$  SEM);  $n = 11$ – $12$  per treatment group. \* $p < 0.05$ .

significant effect of diazepam on the percent time spent in light for DZ $\alpha$ 5 mice ( $F(2,33) = 5.384$ ,  $p = 0.009$ ), and *post hoc* analysis revealed that 10 mg/kg diazepam significantly increased percent time in light in DZ $\alpha$ 5 mice compared with vehicle ( $p < 0.05$ ). There was no significant effect of diazepam on percent time in light in either DZ $\alpha$ 2 mice ( $H(2) = 5.264$ ,  $p = 0.072$ ) or DZ $\alpha$ 3 mice ( $H(2) = 2.803$ ,  $p = 0.246$ ). In DZ $\alpha$ 1 mice, diazepam had a significant effect on percent time in light ( $H(2) = 10.47$ ,  $p = 0.007$ ). *Post hoc* analysis demonstrated that 10 mg/kg decreased the percent time in light ( $p < 0.05$ ), an effect that is likely driven by the strong sedation that diazepam elicits in these mice (Ralvenius et al., 2015). Thus, a statistically significant anxiolytic-like effect was only observed in DZ $\alpha$ 5 mice, indicating that positive allosteric modulation of  $\alpha$ 5-GABA<sub>A</sub>Rs is sufficient to induce an anxiolytic-like effect in this test.

### EPM Test

In a second test of unconditioned anxiety, the EPM test (Figure 3), diazepam caused a significant effect on percent time spent in open arms ( $F(2,33) = 19.441$ ,  $p < 0.001$ ) and percent entries into open arms ( $F(2,33) = 8.099$ ,  $p = 0.001$ ) in WT mice. *Post hoc* analysis revealed that diazepam caused a significant and dose-dependent increase in percent time spent in open arms (2 mg/kg:  $p < 0.05$ ; 10 mg/kg:  $p < 0.001$ ) and in percent entries into the open arms (2 mg/kg:  $p < 0.05$ ; 10 mg/kg:  $p < 0.01$ ), indicative of an anxiolytic-like action in WT mice. In WT mice, diazepam had a significant effect on distance traveled ( $F(2,33) = 6.161$ ,  $p = 0.005$ ) by increasing total distance traveled at 2 mg/kg ( $p < 0.05$ ) but not 10 mg/kg ( $p = 0.983$ ). However, total arm entries were not altered.


Diazepam had a significant effect in DZ $\alpha$ 5 mice on percent time in open arms ( $F(2,33) = 7.679$ ,  $p = 0.002$ ) and percent entries into open arms ( $F(2,33) = 8.944$ ,  $p < 0.001$ ). *Post hoc* analysis demonstrated that 10 mg/kg diazepam caused a significant increase compared with 2 mg/kg diazepam and vehicle in both percent time in ( $p < 0.01$  for both) and percent entries into ( $p < 0.01$  for both) the open arms. Diazepam had a significant effect on total distance traveled ( $H(2) = 11.866$ ,  $p = 0.003$ ), and *post hoc* analysis showed that 10 mg/kg diazepam significantly increased

distance traveled compared with both vehicle and 2 mg/kg diazepam ( $p < 0.05$  for both). Diazepam also had a significant effect on total number of arm entries in DZ $\alpha$ 5 mice ( $F(2,33) = 3.341$ ,  $p = 0.048$ ), and *post hoc* analysis indicated that 10 mg/kg diazepam significantly increased the total number of arm entries compared to vehicle ( $p < 0.05$ ).


In DZ $\alpha$ 3 mice, diazepam had a significant effect on percent time in open arms ( $H(2) = 10.330$ ,  $p = 0.006$ ) and percent entries into open arms ( $F(2,34) = 6.343$ ,  $p = 0.005$ ). *Post hoc* analysis revealed that 10 mg/kg diazepam decreased percent time in open arms ( $p < 0.05$ ) and percent entries in open arms ( $p < 0.01$ ) compared with vehicle. Diazepam also had a significant effect on total distance traveled ( $F(2,34) = 12.176$ ,  $p < 0.001$ ), and *post hoc* analysis revealed that both 2 mg/kg ( $p < 0.05$ ) and 10 mg/kg diazepam ( $p < 0.001$ ) significantly decreased total distance traveled in DZ $\alpha$ 3 mice. Interestingly, there was no change in total arm entries ( $F(2,34) = 1.796$ ,  $p = 0.181$ ), suggesting that diazepam induces locomotor changes in DZ $\alpha$ 3 mice that are different from the nonspecific sedative changes seen in DZ $\alpha$ 1 mice, in which both distance traveled and total arm entries are reduced. DZ $\alpha$ 1 mice displayed a significant decrease in percent time in open arms at both 2 and 10 mg/kg diazepam ( $H(2) = 24.023$ ,  $p < 0.001$ ; *post hoc* Dunn's test  $p < 0.05$  for both), but the percent entries into open arms was not altered ( $H(2) = 3.304$ ,  $p = 0.192$ ). In contrast to DZ $\alpha$ 3 mice, DZ $\alpha$ 1 mice showed significant effects from diazepam in both total distance traveled ( $H(2) = 28.402$ ,  $p < 0.001$ ) and total arm entries ( $H(2) = 24.688$ ,  $p < 0.001$ ) at both 2 mg/kg ( $p < 0.05$ ) and 10 mg/kg diazepam ( $p < 0.05$ ). The strong reductions in total distance traveled and number of total arm entries indicate that the diazepam effect on percent time in open arms is probably an artifact of sedation. Surprisingly, diazepam had no significant effect on the behavioral measurements of DZ $\alpha$ 2 mice in the EPM.

### Novel OF Test


As a third test of unconditioned anxiety, we used a novel OF (Figure 4), where the time spent in the center zone is used as a measure of anxiolysis. In WT mice, 10 mg/kg diazepam significantly increased center zone time ( $H(2) = 10.177$ ,  $p = 0.006$ ) compared with vehicle ( $p < 0.05$ ) and 2 mg/kg



**Figure 3** Elevated plus maze test. Effect of diazepam (2 and 10 mg/kg) in the elevated-plus maze in wild-type (WT) (129X1/Sv) and triple mutant mice on (a) percent time in open arms, (b) percent entries into open arms, (c) distance traveled, and (d) total number of entries into arms. Results are expressed as mean ( $\pm$  SEM);  $n = 11-13$  per treatment group. \* $p < 0.05$ , \*\* $p < 0.01$ , and \*\*\* $p < 0.001$ .



**Figure 4** Novel open-field test. Effect of diazepam (2 and 10 mg/kg) in novel open field in wild-type (WT) (129X1/Sv) and triple mutant mice on (a) center zone duration and (b) distance traveled. The dotted line ('chance level') represents the time in the center zone that would be expected if the distribution of time spent in the center and outside zones was random (408 s in center zone). Results are expressed as mean ( $\pm$  SEM);  $n = 8-13$  per treatment group. \* $p < 0.05$ .



**Figure 5** Stress-induced hyperthermia test. Change in temperature during stress-induced hyperthermia in wild-type (WT) (129X1/SvJ) and triple mutant mice after an injection of vehicle or 3 mg/kg diazepam. Results are expressed as mean ( $\pm$  SEM);  $n = 11$ –16 per treatment group.  $^{**}p < 0.01$ .

diazepam ( $p < 0.05$ ), without a change in total distance traveled, indicative of an anxiolytic-like effect. In *DZα1* mice, diazepam significantly increased center zone duration ( $F(2,31) = 7.131$ ,  $p = 0.003$ ) at both 2 mg/kg ( $p < 0.05$ ) and 10 mg/kg ( $p < 0.01$ ) diazepam. These results may indicate that positive allosteric modulation of  $\alpha 2$ -GABA<sub>A</sub>Rs is sufficient to elicit an anxiolytic-like response in a novel OF. In *DZα2* mice, diazepam significantly increased total distance traveled ( $H(2) = 15.153$ ,  $p < 0.001$ ) at both 2 mg/kg diazepam ( $p < 0.05$ ) and 10 mg/kg diazepam ( $p < 0.05$ ) compared with vehicle. In contrast to WT mice and *DZα2* mice, diazepam had no effect on center zone duration in *DZα3* and *DZα5* mice. In *DZα3* mice, diazepam significantly affected locomotor activity ( $F(2,27) = 3.502$ ,  $p = 0.044$ ) by decreasing total distance traveled at 2 mg/kg diazepam ( $p < 0.05$ ) but not at 10 mg/kg diazepam. Diazepam had no effect on locomotor activity in *DZα5* mice in this test. An increase in the time in the center zone was observed at 10 mg/kg diazepam in *DZα1* mice ( $H(2) = 8.938$ ,  $p = 0.011$ , *post hoc* Dunn's test  $p < 0.05$ ). However, diazepam had a significant dose-dependent effect on distance traveled in *DZα1* mice ( $H(2) = 18.030$ ,  $p < 0.001$ ), and *post hoc* analysis demonstrated that both 2 mg/kg ( $p < 0.05$ ) and 10 mg/kg ( $p < 0.05$ ) diazepam caused significant decreases on total distance traveled. The strong sedation largely precludes meaningful interpretation of the behavior in the *DZα1* mice. The current results show that  $\alpha 1$ -GABA<sub>A</sub>Rs are sufficient for mediating the sedative effect of diazepam, that at least at 10 mg/kg diazepam  $\alpha 3$ -GABA<sub>A</sub>Rs and  $\alpha 5$ -GABA<sub>A</sub>Rs individually do not mediate sedation, and that positive modulation of  $\alpha 2$ -GABA<sub>A</sub>Rs results in a substantial and dose-dependent locomotor stimulation. These findings on locomotor activity are in line with those published in an OF to which animals were habituated (Ralvenius *et al*, 2015).

### Tests of Predictive Validity of Antidepressant-Like Responses and CPP

The high comorbidity between anxiety and mood disorders leads to a strong therapeutic interest in pharmacological agents that can alleviate both anxiety and symptoms of depression. Some benzodiazepines have also been shown to have antidepressant activity in humans (Amsterdam *et al*, 1986; Petty *et al*, 1995), but their sedative action largely precludes testing them in tasks of pharmacological predictability of antidepressant-like responses (also referred to as tests of behavioral despair like the FST and the TST). Thus,

we conducted tests to evaluate whether modulation of specific GABA<sub>A</sub> receptor subtypes, which do not cause sedation ( $\alpha 2$ ,  $\alpha 3$ , and  $\alpha 5$ ), may lead to antidepressant-like effects in the FST and the TST. Surprisingly, we found that diazepam decreased latency to immobility and increased total immobility in *DZα2* mice in the TST, but not the FST, which may thus reflect muscle relaxation (see Supplementary Information and Supplementary Figures S3 and S4 for more details and statistics). Overall, our results in FST and TST do not support a role for modulation of  $\alpha 3$ - or  $\alpha 5$ -GABA<sub>A</sub>Rs in these paradigms. Moreover, modulation of specific GABA<sub>A</sub> receptor subtypes did not result in CPP (Supplementary Table S1).

### Stress-Induced Hyperthermia

The SIH paradigm is used to measure the autonomic component of anxiolytic-like responses. In this paradigm, we used a dose of diazepam (3 mg/kg) that does not cause hypothermia (see  $T_1$  values in Supplementary Figure S5). In WT mice, 3 mg/kg diazepam significantly decreased the SIH ( $T_2 - T_1$ ) ( $t(22) = 2.989$ ,  $p = 0.007$ ). In the *DZα1*, *DZα2*, *DZα3*, and *DZα5* mice, 3 mg/kg diazepam had no significant effect on the stress-induced increase in body temperature (Figure 5). These results indicate that positive allosteric modulation of a single GABA<sub>A</sub> receptor subtype is not sufficient to attenuate SIH at this dose of diazepam that attenuates SIH in WT mice.

### DISCUSSION

In this study, we used a pharmacogenetic 'restriction-of-function' approach that allows diazepam-induced modulation of individual GABA<sub>A</sub> receptor subtypes with very high specificity. Our technique avoids the major limitation of pure pharmacological approaches that no known 'subtype-selective' compound is really specific for any one receptor subtype. Surprisingly, our findings indicate that positive modulation of  $\alpha 5$ -GABA<sub>A</sub>Rs elicits an anxiolytic-like action, as seen in the EPM and LDB tests. While modulation of  $\alpha 2$ -GABA<sub>A</sub>Rs appears to be sufficient to cause anxiolytic-like effects only in the OF, we did not observe an anxiolytic-like effect of positive modulation of  $\alpha 3$ -GABA<sub>A</sub>Rs in any of the tests. The lack of an effect of specific modulation of  $\alpha 2$ -GABA<sub>A</sub>Rs in the EPM and LDB tests is surprising as it has previously been shown (also in 129X1/SvJ mice, then called



129/SvJ mice) that  $\alpha 2$ -GABA<sub>A</sub>Rs are required for anxiolytic-like responses in these two paradigms (Low *et al*, 2000). The current results indicate that  $\alpha 2$ -GABA<sub>A</sub>Rs are not fully sufficient for an anxiolytic-like response in the classical EPM and LDB tests. Possibly, for full anxiolytic-like responses in these paradigms,  $\alpha 2$ -GABA<sub>A</sub>Rs need to be modulated in concert with another GABA<sub>A</sub> receptor subtype ( $\alpha 1$ ,  $\alpha 3$ , or  $\alpha 5$ ), which may facilitate the  $\alpha 2$ -mediated response.

It is a well-known limitation of unconditioned, etiological models of anxiety that changes in locomotor activity may confound interpretation of results (Dawson and Tricklebank, 1995). A recent study has shown that while baseline locomotor activity is indistinguishable between mutant and WT mice in a familiar OF in which the animals were habituated to the test chamber, specific modulation of  $\alpha 1$ -GABA<sub>A</sub>Rs in *Dza1* mice by diazepam substantially decreased the locomotor activity and specific modulation of  $\alpha 2$ -GABA<sub>A</sub>Rs in *Dza2* mice by diazepam substantially increased the locomotor activity, whereas specific modulation of  $\alpha 3$ -GABA<sub>A</sub>Rs and of  $\alpha 5$ -GABA<sub>A</sub>Rs had no effect on locomotor activity in mice (Ralvenius *et al*, 2015). In our experiments, diazepam also had no locomotor effect in the novel OF where conditions are expected to be anxiogenic because of neophobia in *DZa5* mice, however, in the EPM diazepam increased total distance traveled and total arm entries. Interestingly, *Dza2* mice do not show increased locomotion in the EPM, suggesting that the hyperlocomotive effects of diazepam may be context-dependent. Although we consider it unlikely, we cannot exclude the possibility that the anxiolytic-like action of diazepam in the *DZa2* mice in the novel OF is an artefact of the hyperlocomotion induced by diazepam. Finally, the stronger sedative action of diazepam in *Dza1* mice compared with WT may be due to the fact that in contrast to WT mice in *Dza1* mice diazepam does not modulate  $\alpha 2$ -GABA<sub>A</sub>Rs, which increases locomotor activity. In addition to the effects of these nonspecific or neophobia-induced locomotor changes in the findings, the inconsistency of the anxiolytic-like effects between the three paradigms for the *DZa2* and *DZa5* mice, respectively, may also, in part, be due to subtype-specific anxiolytic-like activity not being strong enough to robustly manifest itself across all tests.

Our finding—with the potential limitations discussed above—that diazepam does not elicit anxiolytic-like behavior in the *DZa3* mice in any of the tests used may thus lend credence to the argument that the anxiety-related effects reported with compounds targeting  $\alpha 3$ -GABA<sub>A</sub>Rs may be due to the limited specificity of the compounds. By far the strongest argument for anxiolysis mediated by  $\alpha 3$ -GABA<sub>A</sub>Rs has been made using compound TP003 (Dias *et al*, 2005), which was reported to display a very high selective efficacy for  $\alpha 3$ -GABA<sub>A</sub>Rs, with ~80% efficacy compared with chlordiazepoxide, whereas it had essentially no efficacy at  $\alpha 1$ -,  $\alpha 2$ -, and  $\alpha 5$ -GABA<sub>A</sub>Rs (Dias *et al*, 2005). Based on these *in vitro* data, TP003 was postulated to be an  $\alpha 3$ -selective modulator and had clear anxiolytic-like effects in rats and squirrel monkeys (Dias *et al*, 2005). However, two recent studies failed to replicate the findings of the initial report by Dias *et al* (2005) regarding the subtype selectivity of TP003, and instead demonstrated comparable efficacies at all diazepam-sensitive  $\alpha$ -subunits (Christian *et al*, 2015; de Lucas *et al*, 2015). Moreover, the presumably  $\alpha 3$ -selective

compound SB-205384 (4-amino-7-hydroxy-2-methyl-5,6,7,8-tetrahydrobenzo[*b*]thieno[2,3-*b*]pyridine-3-carboxylic acid, but-2-ynyl ester), which was shown to have anxiolytic-like actions (Navarro *et al*, 2006), was later found to be a positive modulator at  $\alpha 5$ - and  $\alpha 6$ -GABA<sub>A</sub>Rs in addition to  $\alpha 3$ -GABA<sub>A</sub>Rs (Heidelberg *et al*, 2013). Similarly, while the anxiogenic compound  $\alpha 3$ IA (6-(4-pyridyl)-5-(4-methoxyphenyl)-3-carbomethoxy-1-methyl-1*H*-pyridin-2-one) displays some selectivity for  $\alpha 3$ -GABA<sub>A</sub>Rs, it also possesses some efficacy at  $\alpha 2$ -GABA<sub>A</sub>Rs (Atack *et al*, 2005). Finally, the observation that L-838,417 (3-(2,5-difluorophenyl)-7-(1,1-dimethylethyl)-6-[(1-methyl-1*H*-1,2,4-triazol-5-yl)methoxy]-1,2,4-triazolo[4,3-*b*]pyridazine), an  $\alpha 2/\alpha 3/\alpha 5$ -selective partial positive allosteric modulator, has an anxiolytic-like action in the conditioned emotional response test in  $\alpha 2$ (H101R) mice was also interpreted as evidence for a role for  $\alpha 3$ -GABA<sub>A</sub>Rs in anxiolysis (Morris *et al*, 2006). However, it has not been shown that the  $\alpha 2$ (H101R) point mutation abolishes modulation of  $\alpha 2$ -GABA<sub>A</sub>Rs by L-838 417. Furthermore, L-838 417 also modulates  $\alpha 5$ -GABA<sub>A</sub>Rs (McKernan *et al*, 2000), which—as our results may suggest—could mediate the observed anxiolytic-like effects. Thus, our findings combined with these studies suggest that anxiolytic effects cannot be achieved through the positive modulation of  $\alpha 3$ -GABA<sub>A</sub>-Rs.

In the SIH paradigm, chlordiazepoxide effectively reduced SIH because of cage-change stress in  $\alpha 2$ (H101R) mice (Dias *et al*, 2005), indicating that  $\alpha 2$ -GABA<sub>A</sub>Rs are not required for attenuation of SIH. In our study, we found that while diazepam reduced the SIH caused by restraint and the insertion of the rectal probe during the first temperature measurement in WT mice, the highly subtype-specific modulation of  $\alpha 1$ -,  $\alpha 2$ -,  $\alpha 3$ -, or  $\alpha 5$ -GABA<sub>A</sub>Rs was not sufficient to reduce the SIH, suggesting that the concerted modulation of two or more diazepam-sensitive GABA<sub>A</sub> receptor subtypes may be required to reduce this autonomic stress response.

The current finding that positive modulation of  $\alpha 5$ -GABA<sub>A</sub>Rs results in anxiolysis is most surprising, as we reported previously that modulation of  $\alpha 5$ -GABA<sub>A</sub>Rs is not required for the anxiolytic-like action of diazepam (Crestani *et al*, 2002). Presumably, in  $\alpha 5$ (H105R) mice diazepam exerts anxiolysis via  $\alpha 2$ -GABA<sub>A</sub>Rs, perhaps facilitated by  $\alpha 1$ -GABA<sub>A</sub>Rs or  $\alpha 3$ -GABA<sub>A</sub>Rs. However, the question arises why—if positive modulation of  $\alpha 5$ -GABA<sub>A</sub>Rs is sufficient for anxiolysis—diazepam is not anxiolytic in  $\alpha 2$ (H101R) mice (Low *et al*, 2000), in which it modulates  $\alpha 1$ -,  $\alpha 3$ -, and  $\alpha 5$ -GABA<sub>A</sub>Rs. One potential interpretation is that simultaneous modulation of  $\alpha 5$ -GABA<sub>A</sub>Rs and of  $\alpha 1$ - and  $\alpha 3$ -GABA<sub>A</sub>Rs in the  $\alpha 2$ (H101R) mice results in interactions between different GABA<sub>A</sub>R subtypes, which mask the anxiolytic-like effect.  $\alpha 1$ (H101R)/ $\alpha 2$ (H101R) double mutant mice in which diazepam modulates only  $\alpha 3$ -GABA<sub>A</sub>Rs and  $\alpha 5$ -GABA<sub>A</sub>Rs are also resistant to the anxiolytic-like effect of diazepam (Koester *et al*, 2013), indicating that simultaneous positive modulation of  $\alpha 3$ - and  $\alpha 5$ -GABA<sub>A</sub>Rs may not be sufficient to induce anxiolysis. Based on our findings that the modulation of  $\alpha 3$ -GABA<sub>A</sub>Rs reduces open-arm time in the EPM, which is consistent with an anxiogenic-like effect, one possibility is that the diazepam action on  $\alpha 3$ -GABA<sub>A</sub>Rs nullifies the anxiolytic-like action mediated by  $\alpha 5$ -GABA<sub>A</sub>Rs. It is also noteworthy that in *DZa5* mice the anxiolytic-like effect in the EPM requires a higher dose compared with WT mice. It is thus conceivable that a higher receptor occupancy



of  $\alpha 5$ -GABA<sub>A</sub>Rs is required for anxiolysis if these receptors are the only ones modulated by diazepam, and that this receptor subtype may not necessarily contribute to anxiolysis at low doses of diazepam in WT mice.

$\alpha 5$ -GABA<sub>A</sub>Rs are strongly expressed in the hippocampus (Figure 1 and Supplementary Figure 2). As the ventral hippocampus has been linked to emotion and stress, whereas the dorsal hippocampus has been linked primarily to cognitive functions (Fanselow and Dong, 2010),  $\alpha 5$ -GABA<sub>A</sub>Rs in the ventral hippocampus may be involved in anxiolysis. Furthermore, a recent study showed that conditional genetic deletion of  $\alpha 5$  in the central nucleus of the amygdala (CEA) leads to anxiogenic-like effects and increased fear generalization (Botta et al, 2015). The study concluded that extrasynaptic  $\alpha 5$ -GABA<sub>A</sub>Rs in CEA PKC $\delta^+$  neurons control anxiety. Taken together, our results and the study by Botta et al (2015) suggest the possibility that the anxiolytic-like effects of the specific modulation of  $\alpha 5$ -GABA<sub>A</sub>Rs reported here may be due, at least in part, to the positive modulation of the  $\alpha 5$ -GABA<sub>A</sub>Rs in this specific neuronal population.

In conclusion, our findings reveal a role of  $\alpha 5$ -GABA<sub>A</sub>Rs in mediating the affective component of benzodiazepine-induced anxiolysis and no evidence for a role of  $\alpha 3$ -GABA<sub>A</sub>Rs in anxiolysis. Compounds targeting  $\alpha 5$ - and/or  $\alpha 2$ -GABA<sub>A</sub>Rs should lack both the sedative (Rudolph et al, 1999) and addictive (Tan et al, 2010) properties of benzodiazepines, which have been attributed to the  $\alpha 1$ -GABA<sub>A</sub>Rs. Such compounds may provide additional value in the treatment of conditions such as anxious depression, as positive modulation of  $\alpha 2$ -GABA<sub>A</sub>Rs may enhance reward states (Engin et al, 2014; Reynolds et al, 2012) and positive modulation of  $\alpha 5$ -GABA<sub>A</sub>Rs might reduce cognitive problems commonly observed in psychiatric disorders, such as issues with memory interference and cognitive rigidity (Engin et al, 2015).

## FUNDING AND DISCLOSURE

This work was supported by Award Numbers R03MH094834, R01MH080006 and R01MH095905 from the National Institute of Mental Health to UR and Award Number R03DA033491 from the National Institute on Drug Abuse to UR. EE was supported by a National Alliance for Research on Schizophrenia and Depression Young Investigator Award, an Eleanor and Miles Shore Harvard Medical School Fellowship, and an Andrew P Merrill Memorial Research Fellowship. RSB was supported by a Rappaport Mental Health Research Scholar Award. The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Institute on Drug Abuse, the National Institute of Mental Health or the National Institutes of Health. The funders had no role in design, execution, and publication of the work. In the past 3 years, UR has received compensation for professional services from Concert Pharmaceuticals. The other authors declare no conflict of interest.

## AUTHOR CONTRIBUTIONS

LMB, RAF, JL, DB, EE, and AJN performed experiments and analyzed data, RSB and EE supervised LMB, RAF, JL, and AJN in these activities. BKY performed initial experiments

with triple mutant mice, and HUZ provided the four triple mutant mouse lines used and characterized baseline behavior of these lines. UR conceived, designed, and supervised the work that led to this publication. LMB, EE, and UR drafted the manuscript, and all authors revised the manuscript and approved the final version.

## REFERENCES

- Amsterdam JD, Kaplan M, Potter L, Bloom L, Rickels K (1986). Adinazolam, a new triazolobenzodiazepine, and imipramine in the treatment of major depressive disorder. *Psychopharmacology* **88**: 484–488.
- Atack JR, Hutson PH, Collinson N, Marshall G, Bentley G, Moyes C et al (2005). Anxiogenic properties of an inverse agonist selective for  $\alpha 3$  subunit-containing GABA<sub>A</sub> receptors. *Br J Pharmacol* **144**: 357–366.
- Benke D, Fakitsas P, Roggenmoser C, Michel C, Rudolph U, Mohler H (2004). Analysis of the presence and abundance of GABA<sub>A</sub> receptors containing two different types of  $\alpha$  subunits in murine brain using point-mutated  $\alpha$  subunits. *J Biol Chem* **279**: 43654–43660.
- Botta P, Demmou L, Kasugai Y, Markovic M, Xu C, Fadok JP et al (2015). Regulating anxiety with extrasynaptic inhibition. *Nat Neurosci* **18**: 1493–1500.
- Christian EP, Snyder DH, Song W, Gurley DA, Smolka J, Maier DL et al (2015). EEG- $\beta/\gamma$  spectral power elevation in the rat: a translatable biomarker elicited by GABA<sub>A</sub> $\alpha 2/3$ -positive allosteric modulators at non-sedating anxiolytic doses. *J Neurophysiol* **113**: 116–131.
- Crestani F, Keist R, Fritschy JM, Benke D, Vogt K, Prut L et al (2002). Trace fear conditioning involves hippocampal  $\alpha 5$  GABA<sub>A</sub> receptors. *Proc Natl Acad Sci USA* **99**: 8980–8985.
- Dawson GR, Tricklebank MD (1995). Use of the elevated plus maze in the search for novel anxiolytic agents. *Trends Pharmacol Sci* **16**: 33–36.
- de Lucas AG, Ahring PK, Larsen JS, Rivera-Arconada I, Lopez-Garcia JA, Mirza NR et al (2015). GABA<sub>A</sub>  $\alpha 5$  subunit-containing receptors do not contribute to reversal of inflammatory-induced spinal sensitization as indicated by the unique selectivity profile of the GABA<sub>A</sub> receptor allosteric modulator NS16085. *Biochem Pharmacol* **93**: 370–379.
- Dias R, Sheppard WF, Fradley RL, Garrett EM, Stanley JL, Tye SJ et al (2005). Evidence for a significant role of  $\alpha 3$ -containing GABA<sub>A</sub> receptors in mediating the anxiolytic effects of benzodiazepines. *J Neurosci* **25**: 10682–10688.
- Engin E, Bakhurin KI, Smith KS, Hines RM, Reynolds LM, Tang W et al (2014). Neural basis of benzodiazepine reward: requirement for  $\alpha 2$  containing GABA<sub>A</sub> receptors in the nucleus accumbens. *Neuropsychopharmacology* **39**: 1805–1815.
- Engin E, Smith KS, Gao Y, Nagy D, Foster RA, Tsvetkov E et al (2016). Modulation of anxiety and fear via distinct intrahippocampal circuits. *eLife* **5**: e14120.
- Engin E, Zarnowska ED, Benke D, Tsvetkov E, Sigal M, Keist R et al (2015). Tonic inhibitory control of dentate gyrus granule cells by  $\alpha 5$ -containing GABA<sub>A</sub> receptors reduces memory interference. *J Neurosci* **35**: 13698–13712.
- Fanselow MS, Dong HW (2010). Are the dorsal and ventral hippocampus functionally distinct structures? *Neuron* **65**: 7–19.
- Heidelberg LS, Warren JW, Fisher JL (2013). SB-205384 is a positive allosteric modulator of recombinant GABA<sub>A</sub> receptors containing rat  $\alpha 3$ ,  $\alpha 5$ , or  $\alpha 6$  subunit subtypes coexpressed with beta3 and gamma2 subunits. *J Pharmacol Expl Ther* **347**: 235–241.
- Koester C, Rudolph U, Haenggi T, Papilloud A, Fritschy JM, Crestani F (2013). Dissecting the role of diazepam-sensitive  $\gamma$ -aminobutyric acid type A receptors in defensive behavioral reactivity to mild threat. *Pharmacol Biochem Behav* **103**: 541–549.

- Korpi ER, den Hollander B, Farooq U, Vashchinkina E, Rajkumar R, Nutt DJ *et al* (2015). Mechanisms of action and persistent neuroplasticity by drugs of abuse. *Pharmacol Rev* **67**: 872–1004.
- Low K, Crestani F, Keist R, Benke D, Brunig I, Benson JA *et al* (2000). Molecular and neuronal substrate for the selective attenuation of anxiety. *Science* **90**: 131–134.
- McKernan RM, Rosahl TW, Reynolds DS, Sur C, Wafford KA, Atack JR *et al* (2000). Sedative but not anxiolytic properties of benzodiazepines are mediated by the GABA<sub>A</sub> receptor  $\alpha$ 1 subtype. *Nat Neurosci* **3**: 587–592.
- Morris HV, Dawson GR, Reynolds DS, Atack JR, Stephens DN (2006). Both  $\alpha$ 2 and  $\alpha$ 3 GABA<sub>A</sub> receptor subtypes mediate the anxiolytic properties of benzodiazepine site ligands in the conditioned emotional response paradigm. *Eur J Neurosci* **23**: 2495–2504.
- Navarro JF, Buron E, Martin-Lopez M (2006). Anxiolytic-like activity of SB-205384 in the elevated plus-maze test in mice. *Psicothema* **18**: 100–104.
- Petty F, Trivedi MH, Fulton M, Rush AJ (1995). Benzodiazepines as antidepressants: does GABA play a role in depression? *Biol Psychiatry* **38**: 578–591.
- Ralvenius WT, Benke D, Acuna MA, Rudolph U, Zeilhofer HU (2015). Analgesia and unwanted benzodiazepine effects in point-mutated mice expressing only one benzodiazepine-sensitive GABA<sub>A</sub> receptor subtype. *Nat Commun* **6**: 6803.
- Reynolds LM, Engin E, Tantillo G, Lau HM, Muschamp JW, Carlezon WA Jr. *et al* (2012). Differential roles of GABA<sub>A</sub> receptor subtypes in benzodiazepine-induced enhancement of brain-stimulation reward. *Neuropsychopharmacology* **37**: 2531–2540.
- Rudolph U, Crestani F, Benke D, Brunig I, Benson JA, Fritschy JM *et al* (1999). Benzodiazepine actions mediated by specific  $\gamma$ -aminobutyric acid<sub>A</sub> receptor subtypes. *Nature* **401**: 796–800.
- Rudolph U, Knoflach F (2011). Beyond classical benzodiazepines: novel therapeutic potential of GABA<sub>A</sub> receptor subtypes. *Nat Rev Drug Discov* **10**: 685–697.
- Shader RI, Greenblatt DJ (1993). Use of benzodiazepines in anxiety disorders. *N Engl J Med* **328**: 1398–1405.
- Sigel E, Steinmann ME (2012). Structure, function, and modulation of GABA<sub>A</sub> receptors. *J Biol Chem* **287**: 40224–40231.
- Smith KS, Engin E, Meloni EG, Rudolph U (2012). Benzodiazepine-induced anxiolysis and reduction of conditioned fear are mediated by distinct GABA<sub>A</sub> receptor subtypes in mice. *Neuropharmacology* **63**: 250–258.
- Tan KR, Brown M, Labouebe G, Yvon C, Creton C, Fritschy JM *et al* (2010). Neural bases for addictive properties of benzodiazepines. *Nature* **463**: 769–774.
- Vollenweider I, Smith KS, Keist R, Rudolph U (2011). Antidepressant-like properties of  $\alpha$ 2-containing GABA<sub>A</sub> receptors. *Behav Brain Res* **217**: 77–80.

Supplementary Information accompanies the paper on the Neuropsychopharmacology website (<http://www.nature.com/npp>)