

Modelling Hepatic Endoderm Development: Highly Efficient Differentiation of Human Embryonic Stem Cells to Functional Hepatic Endoderm Requires ActivinA and Wnt3a Signalling.

***David C. Hay¹, Judy Fletcher¹, Catherine Payne¹, John D. Terrace¹, Ronald C. J. Gallagher¹, Jan Snoeys⁴, Jim Black¹, Davina Wojtacha¹, Kay Samuel¹, Zara Hannoun¹, Anne Pryde¹, Celine Filippi¹, Ian S. Currie, Stuart J. Forbes², James A. Ross¹, Philip Newsome³⁺, John P. Iredale²⁺**

¹Scottish Centre for Regenerative Medicine, University of Edinburgh, 49 Little France Crescent, Edinburgh, EH16 4SB, U.K.

²Centre for Inflammation Research, University of Edinburgh, 49 Little France Crescent, Edinburgh, EH16 4TJ, U.K.

³Institute of Biomedical Research, University of Birmingham, Wolfson Drive, Edgbaston. Birmingham, B15 2TT, U.K.

⁴ Johnson & Johnson Pharmaceutical Research & Development, Division of Janssen Pharmaceutica N.V., Turnhoutseweg 30, Building 017 Room 105, B-2340 Beerse, Belgium

⁺ Joint Senior Authors, *Corresponding Author

Dr David C Hay,

Scottish Centre for Regenerative Medicine, University of Edinburgh,

49 Little France Crescent, Edinburgh, EH16 4SB,

Scotland, U.K., Email: davehay@talktalk.net

Running title: Highly Efficient Hepatocyte Differentiation from hESCs

ABSTRACT

Human embryonic stem cells (hESCs) are a valuable source of pluripotential primary cells. However, their homogeneous cellular differentiation to specific cell types *in vitro* has proven difficult thus far. Wnt signalling has been shown to play important roles in coordinating development and we demonstrate that Wnt3a is differentially expressed at critical stages of human liver development *in vivo*. The essential role of Wnt3a in hepatocyte differentiation from hESCs is paralleled by our *in vitro* model, demonstrating the importance of a physiological approach to cellular differentiation. Our studies provide compelling evidence that Wnt3a signaling is important for coordinated hepato-cellular function *in vitro* and *in vivo*. In addition, we demonstrate Wnt3a facilitates clonal plating of hESCs capable of hepatic endoderm differentiation. These studies represent an important step forward toward the use of hESC-derived hepatocytes in biomedical applications and has opened the door to high through-put metabolic analysis of human liver function.