Osmotic stress and water isotope effects in kinesin-1 gliding motility assays

Steve Koch, U. New Mexico Dept. Physics and
Center for High Technology Materials (CHTM)

**Thermodynamics and
kinetics of molecular
motors workshop**
Santa Fe, May 20, 2010

Acknowledgments

Larry Herskowitz
Physics Ph.D. Student

Andy Maloney
Physics Ph.D. Student

Anthony Salvagno
Physics Ph.D. Student

Brian Josey
Physics B.S. Student

Pranav Rathi
Optics Ph.D. Student

Collaborations

Susan Atlas—Lead of the DTRA project
UNM Physics / Cancer Center / Director of CARC

Haiqing Liu (G. Mantano lab)—Microdevice applications of kinesin
LANL & Center for Integrated Nanotechnology (CINT)

Postdoc training

George Bachand—Microdevice applications of kinesin
Sandia & Center for Integrated Nanotechnology (CINT)

Funding

DTRA—DTRA CB Basic Research Program under Grant No. HDTRA1-09-1-008

1. Overview of our gliding assay approach

2. Preliminary results isotope and osmotic stress effects in kinesin-1 gliding assays.

Kinesin / MT introduction

‘Kiney’

Gliding motility assay

Andy Maloney
Gliding motility assays
Caseins
Heavy Water
Osmotic Stress

Nature Precedings : doi:10.1038/npre.2010.4479.2 : Posted 28 May 2010

Glass surface

Assay details

Dmk401 30 ug / ml

MTs 29% TRITC cytoskeleton bovine

Room temp. (See Andy's poster for more details)

Video: Beta casein
6x speed; false-colored

← 110 microns →

Gliding motility assay

Nature Precedings · doi:10.1038/npre.2010.4479.2 · Posted 28 May 2010

Andy can finish one sample in
1 hour from start to finish

30 minutes of data, ~9,000 images

Primary measurement: speed

Also other qualitative measurements

Advantage: lots of data quickly;
robust assay

We've developed an open-source, automated microtubule tracking and speed analysis platform

Larry Herskowitz
Image tracking
Stochastic Simulation
Image Simulation

1. Automated Tracking of MTs

MTs identified with NI Vision 7.1
Segmenting Routines

Position and Angle of MT ends
found via image
pattern matching

X, Y position versus time recorded
for all MTs in folder heirarchy

30 minutes of raw data processed
in approx.
1 hour using 4 cores

See Larry's poster today for details

Template for
pattern matching

2. Automated Speed Analysis

A. Track smoothing

X, Y data are noisy; dominated by Brownian motion of MT ends
Smoothed with Gaussian-weighted 2 second window

Nature Precedings : doi:10.1038/npre.2010.44790v2 May 28, 2010

2. Automated speed analysis

B. Instantaneous speed

$$\text{Speed}_i = |\Delta r_i| / \Delta t_i$$

Alternative to fitting paths to simple shapes

Careful for smoothing errors

What is gliding speed anyway?

Nature Precedings : doi:10.1038/npre.2010.4479.2 : Posted 28 May 2010

2. Automated speed analysis

C. Kernel density estimation of speed probability distribution function (PDF)

Gaussian kernel, bandwidth 50 nm / s

Silverman 1986 Density Estimations

Nature Precedings : doi:10.1038/npre.2010.4479.2 : Posted 28 May 2010

2. Automated speed analysis

C. Kernel density estimation of speed probability distribution function (PDF)

We checked our algorithm using simulated microtubule image series*
constant gliding speed w/ pause of random duration

* Herskowitz and Koch (2010) Nature Precedings <http://dx.doi.org/10.1038/npre.2010.4374.1>

2. Automated speed analysis

C. Kernel density estimation of speed probability distribution function (PDF)

We checked our algorithm using simulated microtubule image series*
constant gliding speed w/ pause of random duration

Calculated speed 805.4 +/- 0.2 nm/s (N=15) Actual gliding speed 807 nm/s

Pauses probably bias data a bit ... more significant with wider bandwidth

2. Automated speed analysis

C. Speed PDF analysis not greatly affected by pauses or tracking errors

Permits automated speed determination for pooled MT tracks for given condition

Nature Precedings : doi:10.1038/npre.2010.4479.2 : Posted 28 May 2010

2. Automated speed analysis

C. Speed PDF analysis not greatly affected by pauses or tracking errors
Permits automated speed determination for pooled MT tracks for given condition

Alpha casein passivation

N=710 MT tracks

~30 minutes assay time

Analysis time ~10 minutes on one core

2. Automated speed analysis

Speed gradually increases with time for all casein varieties

Intrinsic speed differences between varieties

Nature Precedings : doi:10.1038/npre.2010.4479.2 : Posted 28 May 2010

Beta casein highest speed

Differences in active motor density?

We chose alpha casein and these analysis methods for subsequent isotope and osmotic stress studies

See Andy Maloney's poster for more details

See also: Verma, V., Hancock, W. O., & Catchmark, J. M. (2008). *Journal of biological engineering*, 2, 14. doi: 10.1186/1754-1611-2-14.

Varying the properties of water may be a useful knob for studying kinesin

Water isotope effects

Heavy-hydrogen water (D_2O)

Heavy-oxygen water ($H_2^{18}O$)

Osmotic stress

Betaine, Future work: other osmolytes: proline, sucrose, etc.

Future: denaturants

May be a method for probing large-surface area interactions between kinesin and tubulin

Osmotic stress dramatically increases lifetime of bound molecular complexes

An example from protein-DNA interactions

No kinesin studies yet;
A few myosin studies

Sidorova and Rau
Biopolymers, Vol. 53, 363–368 (2000)

$$\frac{d[\ln(k_{\text{off}})]}{d[\text{osmolal}]} = -\frac{\Delta N_w}{55.6}$$

Off rate reduced by >100 fold

Heavy-hydrogen water (D₂O) has been used to probe the actomyosin system

Nature Precedings : doi:10.1038/npre.2010.4479.2 : Posted 28 May 2010

Fig. 1. Dependence of the actin sliding velocity on pH (●) and pD (○) of the H₂O and D₂O assay solution, respectively. Each data point represents the average velocity of 11 different filaments, and the vertical bar shows the standard deviation. The curves were drawn by eye.

The authors list among many possible explanations, an effect on myosin-actin affinity.

How should water isotopes affect kinesin motility?

Answers from molecular dynamics?

Chaen, S., et al (2001). Effect of deuterium oxide on actomyosin motility in vitro. *Biochimica et biophysica acta*, 1506(3), 218-23. Pub med: 11779555.

Another reason we're looking at water effects is our partnership with Atlas / Valone labs

Susan Atlas (UNM; PI of our DTRA project) and Steve Valone (LANL) are developing the "Charge-Transfer Embedded-Atom Model (CT-EAM)" force field for MD simulations of kinesin and other biomolecules.

Incorporates "correct" quantum mechanics so that charge transfer can be handled.

One of the first benefits of CT-EAM will be better simulation of water dynamics...including for differing isotopes.

Big disconnect between length and time scales for gliding assay and MD

But both can look at water dynamics.

MD: How many hydrating water molecules?

How do water dynamics change with isotope?

Stochastic simulation can help us understand how water might affect kinesin motility

We can ask question:
“What will happen if
kinesin-MT off rate
decreases?”

The answers aren't
always obvious when
pathway can branch.

Larry Herskowitz has developed an agent-based stochastic simulation to help us interpret and guide experiments
more info at his poster

Stochastic simulation can help us understand how water might affect kinesin motility

- Default Parameters
- 2 times Default on rate; 1/2 times Default off rate
- 3 times Default on rate; 1/3 times Default off rate
- 5 times Default on rate; 1/5 times Default off rate
- 10 times Default on rate; 1/10 times Default off rate
- 1/2 times Default on rate; 2 times Default off rate

Stochastic simulations tentatively indicate that increased osmotic pressure or increased water mass would decrease kinesin speed.

Heavy-hydrogen water reduces MT gliding speed.

As with regular water, speed gradually increases with time

Mean and SEM for last 8 points of each set.

Heavy-hydrogen water reduces MT gliding speed.

Speed steadily decreases as deuterium conc. Increases.

Also, noticed other features (preliminarily):

Reduced photobleaching

Longer-lasting MTs

MT-MT sticking

Other features seen in D2O assays

Also, noticed other features (preliminarily):

Reduced photobleaching

Longer-lasting MTs

MT-MT sticking

It turns out there is existing literature to support all of these observations

Heavy-hydrogen water has significantly different properties from “regular” water

Naturally abundant 1 / 6600 hydrogen molecules is deuterium

17 mM deuterium in “standard mean ocean water”

11% denser than H₂O. Freezes at 3.8C. D-bonds stronger.

Toxic to eukaryotes. The toxic effects are similar to chemotherapeutic drugs.

D₂O has been used to stabilize viral vaccines.

D₂O stabilizes tubulin and microtubules.

D₂O stimulates tubulin assembly formation.

(Other fascinating factoids...)

Effects on kinesin motility has not yet been studied

Interesting ... but very complicated

Heavy-oxygen water

Nature Precedings : doi:10.1038/npre.2010.4479.2 : Posted 28 May 2010

Surprisingly, heavy-oxygen water has similar effect on gliding speed

Nature Precedings : doi:10.1038/npre.2010.4479.2 : Posted 28 May 2010

Chemistry of oxygen-18 water similar to regular water

Viscosity of oxygen-18 water only 5% higher than regular water (23% for D₂O)*.

Heavy-O water much less confusing, possibly more tractable theoretically. Expensive, though (\$1,000 / gram)

Possible that Heavy-O water is a good probe of kinesin-MT surface interaction

*Viscosity: Kudish et al. *J. Chem. Soc., Faraday Trans. 1*, 1972, **68**, 2041 - 2046,

Betaine (osmolyte) substantially reduces gliding speed

Unsure how important viscosity increase is.

Need to repeat with other osmolytes

Current conclusions from water and other experiments

Heavy-oxygen water looks like promising probe of hydration effects

Heavy-hydrogen water may also be interesting probe, but more complicated.
May have benefits for stabilization of protein in the lab or in microdevices.

Betaine has strong effect on gliding speed. Viscosity a confounding variable.
Need to compare with other osmolytes.

Need a better grasp of solution thermodynamics and kinetics and effect on
motor proteins.

Type of surface passivation affects gliding speed. Gliding speed also increases
with time.

Open Data, Open-source Software

All data presented here is CC0-licensed and available online:

http://openwetware.org/wiki/Koch_Lab:Data/MT_Gliding_Assay_Readme_File

We're still working on the organization and how to best share. In the meantime:

If useful to you, get in touch and we can help you with the software and / or figuring out the data sets!

Acknowledgments

Larry Herskowitz
Physics Ph.D. Student

Andy Maloney
Physics Ph.D. Student

Anthony Salvagno
Physics Ph.D. Student

Brian Josey
Physics B.S. Student

Pranav Rathi
Optics Ph.D. Student

Collaborations

Susan Atlas—Lead of the DTRA project
UNM Physics / Cancer Center / Director of CARC

Haiqing Liu (G. Mantano lab)—Microdevice applications of kinesin
LANL & Center for Integrated Nanotechnology (CINT)

Postdoc training

George Bachand—Microdevice applications of kinesin
Sandia & Center for Integrated Nanotechnology (CINT)

Funding

DTRA—DTRA CB Basic Research Program under Grant No. HDTRA1-09-1-008