

## Tracing blastomere fate choices of early embryos in single cell culture

2

Mingyou Li<sup>1\*</sup>, Ni Hong<sup>1\*</sup>, Zhendong Li<sup>1\*</sup>, Meisheng Yi<sup>1</sup>, Rong Liu<sup>1</sup>, Haobin Zhao<sup>1</sup>,  
4 Yan Yan<sup>1,2</sup>, Yongming Yuan<sup>1</sup>, Changming Li<sup>2</sup>, Ruowen Ge<sup>1</sup>, Jianxin Song<sup>1\*\*</sup>,  
Yunhan Hong<sup>1\*\*</sup>

6

<sup>1</sup>Department of Biological Sciences, National University of Singapore, 14 Science  
8 Drive 4, Singapore 117543; <sup>2</sup>Division of Bioengineering, School of Chemical and  
Biomedical Engineering, Nanyang Technological University, 70 Nanyang Drive,  
10 Singapore 637457

12

\*These authors contributed equally to this work

14

\*\*Correspondence:

16

Professor Yunhan Hong ([dbshyh@nus.edu.sg](mailto:dbshyh@nus.edu.sg))

Professor Jianxin Song ([jianxin\\_song@nuhs.edu.sg](mailto:jianxin_song@nuhs.edu.sg))

18

Department of Biological Sciences

National University of Singapore

20

14 Science Drive 4

Singapore 117543

22

Fax: +65 6779 2486; Tel: +65 6516 2915

24

**Key words:** blastomere, cleavage, fate tracing, neural crest, pigment, single cell  
culture, yolk syncytium

26

28

**Blastomeres of early vertebrate embryos undergo numerous fate choices for division, motility, pluripotency maintenance and restriction culminating in various cell lineages. Tracing blastomere fate choices at the single cell level in vitro has not been possible because of the inability to isolate and cultivate early blastomeres as single cells. Here we report the establishment of single cell culture system in the fish medaka, enabling the isolation and cultivation of individual blastomeres from 16- to 64-cell embryos for fate tracing at the single cell level in vitro. Interestingly, these blastomeres immediately upon isolation exhibit motility, lose synchronous divisions and even stop dividing in  $\geq 50\%$  cases, suggesting that the widely accepted nucleocytoplasmic ratio controlling synchronous divisions in entire embryos does not operate on individual blastomeres. We even observed abortive division, endomitosis and cell fusion. Strikingly,  $\sim 5\%$  of blastomeres in single cell culture generated extraembryonic yolk syncytial cells, embryonic stem cells and neural crest-derived pigment cells with timings mimicking their appearance in embryos. We revealed the maternal inheritance of key lineage regulators and their differential expression in cleavage embryos. Therefore, medaka blastomeres possess the accessibility for single cell culture, previously unidentified heterogeneity in motility, division, gene expression and intrinsic ability to generate major extraembryonic and embryonic lineages without positioning cues. Our data demonstrate the fidelity and potential of the single cell culture system for tracking blastomere fate decisions under defined conditions in vitro.**

24

Numerous cell fate choices occur throughout the animal life. Early in development of diverse animal species, fertilization between a sperm and an egg leads to the totipotent zygote, which undergoes continuous divisions and lineage restrictions, producing an increasing cell number and creating many different cell types. A mammalian embryo undergoes two cell fate decisions until the blastocyst stage, generating two extraembryonic lineages, the trophectoderm (TE) and primitive endoderm (PE), and an embryonic lineage, the epiblast for the future body<sup>1</sup>. The epiblast is pluripotent and generates three germ layers (ectoderm, mesoderm and endoderm), neural crest (NC) and germline. It is a fundamental challenge in developmental biology when and how different cell fates are precisely determined and regulated. Fate tracing in vivo is powerful for analyzing cell fate choices in developing embryos and adult tissues<sup>2</sup>. Recently, in vitro fate tracing has been developed by using either whole embryos<sup>3</sup> or pooled cell populations<sup>4-6</sup>. In no organism has it so far been possible to culture single blastomeres for extended period to trace cell fate choices. Here we chose medaka (*Oryzias latipes*) to establish a single cell culture system to trace blastomere fate choices in vitro.

### 18 **Medaka embryos are accessible for blastomere isolation**

Medaka is an excellent lower vertebrate model for embryonic development<sup>7</sup>, stem cell culture<sup>2,8-13</sup> and has a unique embryology favoring blastomere isolation. In many fish species including zebrafish, early blastomeres undergo meroblastic cleavages atop but not through the yolk<sup>14</sup>, which are incomplete cell divisions and thus prevent blastomere isolation. In medaka, we found a seemingly intact yolk

membrane from the 2-cell stage onwards after in situ hybridization with antisense  
2 RNA probes against various genes such as *boule* and *dazl*<sup>15</sup>, *mitf1*  
(Supplementary Figure 1a-c) and *vasa* (Supplementary Figure 1g and h), which  
4 is easily separable together with the blastoderm from the yolk (Supplementary  
Figure 1d-f and i). To determine when the yolk membrane establishes its  
6 intactness, a fluorescent dye was microinjected into different positions of freshly  
fertilized eggs or one cell of the 2-cell embryos and the dye distribution was  
8 monitored at the 64-cell stage. The dye injected into the yolk near the cytoplasm  
at the animal pole was transported to the future blastoderm (Figure 1a), whereas  
10 the dye injected at the vegetal yolk was not (Figure 1b), and the dye injected into  
one cell of 2-cell embryos was restricted to daughter cells of the injected  
12 blastomere (Figure 1c). Taken together, a prominent yolk membrane is formed  
and acquires its structural and physiological integrity at the first cleavage, which  
14 allows for blastomere isolation in subsequent stages.

### **Tracing fate choices of blastomere divisions in culture**

16 We chose medaka strain HB32C for cell isolation and culture, a permissive strain  
for blastula cell culture in gelatin-coated multiwell plates<sup>10</sup>. We first determined  
18 the developmental stages for the possibility and efficiency of single cell isolation  
and cultivation. Isolated cleavage blastomeres were capable of survival and/or  
20 proliferation in single cell culture, producing an efficiency of 26%, 49% and 67%  
for 16-, 32- and 64-cell blastomeres, respectively (Table 1). Thus, blastomeres  
22 can reliably be isolated from 16- to 64-cell stages for single cell culture under  
defined conditions.

We began with the 16-cell stage (Figure 1e) to examine cell divisions as  
2 the first fate choice of blastomeres. In many animals, both invertebrates and  
vertebrates including *Xenopus*<sup>16</sup> and zebrafish<sup>14</sup>, all blastomeres until the  
4 midblastula transition undergo 10~12 abbreviated cycles of rapid synchronous  
cell divisions without G1 and G2 phases and motility<sup>16</sup>. The cause and  
6 mechanisms for cell synchrony of abridged divisions have been a mystery since  
long. In both *Xenopus* and zebrafish, rapid cleavages have been thought of as  
8 being controlled by a cytoplasmic clock, which operates independently of the  
nucleus but dependent on the nucleocytoplasmic ratio<sup>14,16</sup>. We found that  
10 medaka 16-cell blastomeres displayed considerable differences in morphology,  
with their diameters ranging from 50  $\mu\text{m}$  to 100  $\mu\text{m}$  (Figure 1f and Supplementary  
12 Figure 2a). Strikingly, blastomeres immediately upon isolation lost synchronous  
cell divisions and even stopped dividing, and exhibited motility as evidenced by  
14 movements and pseudopodia (Supplementary Figure 2; Supplementary Movie 1).  
Notably, motility was seen also in actively dividing blastomeres (Supplementary  
16 Figure 2; Supplementary Movies 1 and 2). Within the first 2 hour of culture, we  
observed abortive cell divisions (Supplementary Figure 2a; Supplementary Movie  
18 1) and cell fusions between dividing and non-dividing blastomeres  
(Supplementary Figure 2e-h; Supplementary Movie 2). More importantly, we  
20 observed fast dividing, slow dividing and non-dividing blastomeres, with the fast  
dividing blastomere having completed four cycles of divisions within 2 h  
22 (Supplementary Figure 2e-h; Supplementary Movie 2), comparable to rapid  
cleavages in intact embryos<sup>17</sup>. Taken together, rapid division is maintained in one

or few blastomeres but lost in the remainder. The nucleocytoplasmic ratio  
controlling rapid synchronous divisions in entire cleavage embryos does not  
appear to operate on isolated single blastomeres in culture.

We then seeded individual blastomeres in gelatin-coated 96-well plates for  
single cell culture (Figure 1d) to tracking behaviors for extended periods of time  
at the single cell level. To observe nuclear behaviors in more detail, we explored  
zygotic microinjection of RNA encoding a fusion between the histone 2B and  
green fluorescent protein (H2Bgfp) to visualize nuclei. This led to the  
identification of three major classes of 16-cell blastomeres (Table 1): 74%  
belonged to class I, which ceased cell divisions even after completion of one  
nuclear division (Figure 1g), 6% was class II that underwent endomitosis  
(Supplementary Figure 3), and 20% fallen into class III, which divided three or  
more times within 24 h of culture and produced daughter cells of both  
heterogeneous (Figure 1h) and homogeneous sizes (Figure 1i and j). We also  
observed incomplete divisions (Supplementary Figure 3a). In particular, certain  
blastomeres continued rapid divisions and formed a cluster of ~100 cells at 24 h  
of culture (Figure 1j). Clearly, blastomeres have the heterogeneity in division,  
suggesting again the presence of novel mechanisms in medaka other than the  
nucleocytoplasmic ratio.

#### **Tracing fate choices of blastomere lineage restriction in single cell culture**

We tested the usefulness of single cell culture for tracing blastomere fate choices  
in lineage restriction to the yolk syncytial layer (YSL), pluripotent embryonic stem  
(ES) cells and pigment cells. A medaka embryo at the morula stage has two cell

populations, the envelope layer (EVL) and deep cells. The deep cells are  
2 equivalent to the inner cell mass in mouse, which is then separated into the deep  
cell layer (DCL) and YSL during blastulation<sup>17</sup>. Therefore, a medaka blastula  
4 embryo is similar to a mouse blastocyst embryo in having three blastula lineages:  
EVL, DCL and YSL, which are equivalent to the mouse TE, epiblast and PE,  
6 respectively (Fig. 2a). Around 6.5 h post fertilization at the early blastula stage  
with ~1000 cells, YSL is seen as 4-5 layers of nuclei<sup>17</sup>, which became easily  
8 visible upon nuclear labeling by H2B<sub>gfp</sub> RNA injection (Fig. 2b). The DCL  
contributes to the future embryo body and is capable of generating diploid and  
10 even haploid ES cell cultures<sup>10,11,13</sup>. Pigment cells originate from the NC, a  
transient population arising from the neural plate in vertebrate embryos, which  
12 migrate throughout the body to generate many other cell types<sup>18</sup>. NC is elusive  
for analyses of fate choices because of a transient and migratory nature. In  
14 medaka, pigment cells become visible at day 3 post fertilization<sup>8,11,17</sup>, which  
comprise black-pigmented melanophore and other chromatophores including the  
16 ridophore<sup>19</sup>, which in medaka is autofluorescent<sup>8,11</sup>.

We found that daughter cells of certain 16-cell blastomeres were capable  
18 of generating YSL in single cell culture. As early as 4 h of culture, YSL precursors  
appeared as round and large-sized (~50  $\mu$ m in diameter) cells, in which the  
20 syncytial cytoplasm was located peripherally and the nuclei were positioned  
centrally (Figure 2c). Subsequently, nuclei moved to the periphery and the  
22 cytoplasm moved to the center until 6 h of culture (Figure 2d). They developed  
into YSL cells until day 1, which had multiple prominent nuclei in the periphery

surrounding the centrally residing cytoplasm (Fig. 2e). The 16-cell blastomere-derived YSL cells were indistinguishable from those from midblastula embryos in culture (Figure 2f). Interestingly, the YSL cytoplasm actively formed pseudopodia (Supplementary Figure 4; Supplementary Movie 3). Moreover, 16-cell blastomeres produced a cluster of actively dividing cells at day 1 post culture (Figure 1j), which eventually developed into compacted cells at day 3 (Figure 2g), phenotypically resembling ES cells<sup>10,13</sup>. At day 3, melanocytes began to appear, which exhibited the characteristics of NC-derived pigment cells, including a flat shape, pigmented granules and dendritic processes (Figure 2g). At day 4 of culture, iridocytes became clearly visible by autofluorescence (Figure 2h). We obtained similar results with 32-cell blastomeres (Supplementary Figure 5) and 64-cell blastomeres (Supplementary Figure 6). Although the frequency of non-dividing blastomeres decreased to ~50% for the 32- and 64-cell stages, the proportion of blastomeres capable of lineage restriction remained 4~6% from 16- to 64-cell stages (Table 1). Collectively, single 16- to 64-cell blastomeres possess the ability to generate extraembryonic YSL cells, ES cells and NC-derived pigment cells.

In single cell culture, YSL cells appeared at day 1 and pigment cells appeared at day 3, reminiscent of the timing of their appearance in developing embryos. Interestingly, only proliferative blastomeres were capable of generating ES cells, YSL and pigment cells; In contrast, non-dividing blastomeres did survive for up to 5 days of culture without showing any sign of differentiation (Supplementary Movie 4). These results point to the fidelity of in vitro lineage



tracing in single cell culture. Convincingly, the ability to generate major blastula lineages and differentiated pigment cells of the NC origin is intrinsic to blastomeres and present already at the 16-cell stage, prior to the separation into the three blastula lineages until the 1000-cell stage and far before the organogenesis stage when pigment cells become visible.

## 6 **Expression of key lineage regulators**

In order to determine the molecular basis of the inherent potential for ES cells, YSL and pigment cells, we analyzed the RNA expression of *nanog*, *oct4*, *sox17* and microphthalmia-associated transcription factor (*mitf*). *nanog* and *oct4* are key pluripotency regulators in mouse<sup>20,21</sup> and exhibit pluripotent expression in medaka<sup>22-24</sup>. Mouse *sox17* is a key regulator of PE development<sup>25</sup>. Fish *mitf* is a master regulator of pigment cells of NC lineage<sup>26</sup>. In mice, only *oct4* is maternally supplied<sup>21</sup>, whereas *nanog*<sup>20</sup>, *sox17*<sup>25</sup> and *mitf*<sup>27</sup> commence their expression at the 8-cell stage, 32-cell stage and around embryonic day 10, respectively. We found that the transcripts of medaka *oct4* and *nanog* were maternally supplied and expressed in cleavage embryos (Supplementary Figure 7), consistent with their reported expression<sup>22-24</sup> and the ES cell formation from cleavage blastomeres in single cell culture. Surprisingly, the transcripts of *sox17* and *mitf1* (one of the two medaka *mitf* genes due to an ancient whole genome duplication event in the fish lineage) were also maternal and persistent in cleavage embryos (Supplementary Figures 1 and 8). More intriguingly, *sox17* was predominant in central blastomeres of cleavage embryos (Supplementary Figure 8a and b), and became preferentially expressed in YSL cells when they are formed at the early

blastula stage (Supplementary Figure 8c) and marginal blastomeres that produce  
2 the embryonic endoderm<sup>28</sup>, whereas *mitf1* distributed preferentially in peripheral  
blastomeres (Supplementary Figure 8d-f). Differences in temporospatial  
4 distribution becomes more evident after two-color fluorescent in situ hybridization  
(Figure 3), reinforcing the heterogeneity of blastomeres at the molecular level.  
6 Therefore, medaka 16-cell blastomeres already exhibit differential expression of  
lineage markers, in contrast to the mouse situation where all 16-cell blastomeres  
8 retain the ability to contribute to any of the three blastocyst lineages<sup>29</sup> and show a  
similar gene expression profile<sup>30</sup>. Collectively, the intrinsic potential of early  
10 blastomeres to produce YSL, ES cells and pigment cells accompanies the  
expression of respective lineage markers.

## 12 **Discussion**

In this study, the establishment of a single blastomere culture system and its  
14 exploitation in tracing cell fate choices lead to twelve important findings that are  
novel or different from previous results obtained in whole embryos. First, we  
16 show in medaka that embryonic cells as early as the 16-cell stage can be  
robustly isolated without losing their viability and developmental potential.  
18 Conceptually, meroblastic cleavages in fish are thought to be incomplete due to  
the lack of an intact yolk membrane and thus prevent isolation of intact and viable  
20 blastomeres. Three lines of evidence point to the intactness of yolk membrane in  
early medaka embryos, leading to our choice for this organism for single  
22 blastomere isolation. In situ hybridization reveals that maternal RNAs often do  
not completely enter the cellular blastodisc but leave a substantial level in the

yolk to form an area beneath the blastodisc, indicating the presence of a barrier  
2 between the yolk and blastodisc as early as the 2-cell stage. Furthermore, the  
yolk membrane together with the blastoderm is easily separable from the yolk.  
4 Convincingly, fluorescent dye injected in the vegetal yolk completely remains in  
the yolk, demonstrating the physiological integrity of the yolk membrane.  
6 Therefore, medaka is unique among organisms with meroblastic cleavages in its  
accessibility for blastomere isolation at the cleavage stages. In this regard,  
8 mammalian embryos must be also accessible for blastomere isolation because of  
complete cleavages. Second, we show that early blastomeres can survive and  
10 divide in single cell culture in the absence of any feeder cells. Third, we reveal  
that early blastomeres manifest motility upon isolation, in contrast to previously  
12 described absence of motility in blastomeres until the midblastula stage in many  
egg-laying species including *Xenopus*<sup>31</sup> and zebrafish<sup>14</sup>. Notably, only a subset of  
14 isolated 16-cell blastomeres exhibit pseudopod formation, while the remainder  
does not show this measure of motility during whole period of culture. It appears  
16 that isolation and cultivation allows for manifestation and easy detection of  
motility, rather than stimulating or inducing motility. Even after midblastula  
18 transition in zebrafish, pseudopodia appear in interphase but do not form during  
mitosis<sup>14</sup>. In medaka, pseudopodial formation has not been described before the  
20 midblastula stage. It is likely that rapid cleavage divisions and positioning in the  
embryo prevent pseudopodial formation and cell movement. In this study, even  
22 dividing blastomeres formed pseudopodia and moved round in culture,  
suggesting that mitosis does not prevent cellular motility in vitro. Fourth, we

demonstrate the previously unidentified heterogeneity of blastomeres in morphology, motility, division, gene expression and more importantly, developmental potential in single cell culture. Fifth, our finding that  $\geq 50\%$  of 16~64-cell blastomeres maintain a large size but stop dividing immediately after isolation indicates the presence of unknown mechanisms that regulate synchronous blastomere cleavages. In invertebrates and lower vertebrates including fish, it is widely accepted that all embryonic cells undergo rapid and synchronous divisions until the midblastula stage of  $\sim 2000$  cells when the midblastula transition begins, and cell synchrony of continuous divisions has widely been thought of as being determined by the nucleocytoplasmic ratio<sup>14</sup>. It remains unknown why this ratio does not operate on isolated blastomeres. It is likely that the synchrony of rapid divisions requires cell-cell communications. Intriguingly, there are certain blastomeres that continue rapid divisions in single cell culture. Possibly, such blastomeres may initiate and orchestrate synchronous cleavage divisions via cell-cell interactions. Sixth, we have also observed abortive division, endomitosis and cell fusion in isolated blastomeres, which have not yet been described in early cleavage embryos of lower vertebrates. All these processes may each lead to higher ploidy levels in certain embryonic cells. We have previously reported the invariant presence of  $\sim 20\%$  tetraploid subpopulation in medaka diploid ES cells derived from fertilization midblastula embryos<sup>10</sup> and the presence of  $\sim 20\%$  diploid subpopulation in medaka haploid ES cells derived from gynogenetic midblastula embryos<sup>13</sup>. Whether these variations are causative for the appearance of higher ploidy levels in ES cells remains unknown. Seventh,

we show in medaka that the extraembryonic YSL cells can intrinsically be  
2 specified from cleavage blastomeres. In mouse, the two earliest events leading to  
the separation of TE and PE have been found under the multifaceted regulation  
4 including cell polarity, position and gene expression<sup>1,29</sup>. Eighth, the single cell  
culture system revealed the dynamic process of YSL cell formation and the  
6 motility of YSL cytoplasm. In zebrafish, some marginal blastomeres are confluent  
with the yolk cell cytoplasm resulting from incomplete division collapse and  
8 deposit their nuclei and cytoplasm into the cytoplasmic cortex of the yolk cell,  
thereby forming the YSL, whose nuclei then undergo three to five rounds of  
10 endomitosis without cytokinesis<sup>14,32</sup>. In medaka, the YSL originates also from  
marginal blastomeres and its nuclei undergo endomitosis<sup>17</sup>, but apparently in the  
12 absence of cell fusion with the yolk, because the yolk membrane acquires its  
integrity already at the 2-cell stage and prevents the direct interaction, as we  
14 have shown in this study. The YSL is elusive for study due to difficulties in  
interfering specifically with YSL formation and morphogenesis<sup>32</sup>. Our  
16 observations make medaka model for experimental analyses of YSL formation  
and its separation from the pluripotent DCL. In addition, YSL formation in single  
18 cell culture will provide an excellent system to elucidate the mechanisms  
underlying endomitosis and its role in cell fate specification. Ninth, we show that  
20 certain 16-cell blastomeres have the potential to generate pluripotent ES cells, far  
before the blastula stage when deep cells are formed. This is different from the  
22 situation in mouse, where a pluripotent ground state is established in the inner  
cell mass and epiblast until the blastocyst stage<sup>33</sup>. Tenth, the fact that medaka

blastomeres even at as early as the 16-cell stage already possess the intrinsic  
2 ability to specify pigment cells as the representative NC derivatives suggest the  
cell-autonomous NC specification in this organism, which is in contrast to the  
4 widely accepted notion that NC is induced from the neural tube by BMP and Wnt  
signalings<sup>19,34</sup>. Eleventh, we show that key lineage regulators are maternally  
6 supplied in medaka, in contrast to their zygotic expression in mouse<sup>27</sup>. Finally,  
our observations that lineage formation in single cell culture is dependent on cell  
8 divisions and timings similar to developmental programs of a developing embryo  
strongly suggest the fidelity of the single cell culture system to recapitulate  
10 blastomere fate choices in vivo.

Taken together, we have successfully developed a novel ability to isolate  
12 and cultivate individual blastomeres of early medaka embryos and investigated  
their fate choices. It is evident that the single cell culture system is ideal for fate  
14 tracing in vitro under defined culture conditions without an intact embryonic  
environment. Our success is ascribed to the blastomere accessibility for isolation  
16 and well-defined culture conditions<sup>10,13,35</sup>. We believe that this single cell culture  
system can be developed also in other organisms to trace the behaviors/fates at  
18 the single cell level in culture, offering invaluable lights on the mode, process and  
mechanisms of various cell fate choices.

## 20 **METHODS**

**Embryo microinjection and cell culture.** Embryos were microinjected with 50  
22 ng of H2B<sub>gfp</sub> RNA at the 1-cell stage as described<sup>36</sup>. Embryo manipulation, cell  
isolation and culture were done essentially as described<sup>8,10-13</sup>. Briefly, embryos

1 were treated with proteinase K (10 mg/ml) for 60 min at 28°C to remove the  
2 attachment filaments, rinsed twice in phosphate-buffered saline (PBS) and  
sterilized in PBS-0.1% bleach for 2 min, and rinsed 5 times in PBS. Embryos  
4 were incubated in PBS and monitored for developmental stages under aseptic  
conditions. The chorion was manually torn with a pair of fine forceps at the  
6 vegetal half. For embryos at 32- and 64-cell stages, cells were dissociated by  
gentle pipetting. For embryos at the 16-cell stage, cells were individually  
8 dissociated by using a fine forceps to prevent damage. Healthy cells were  
seeded by pipetting into gelatin-coated 96-well plates containing 150 µl of ES cell  
10 culture medium ESM2. Within 10 min of seeding, the plates were monitored  
under an invert microscope to ensure that one cell was present each well. Cell  
12 growth, attachment, proliferation and differentiation were monitored at regular  
intervals of culture at 28°C in air.

14 Procedures for fish maintenance, gene cloning, RNA synthesis and in situ  
hybridization were performed as described in Supplementary Methods.

16 **Full methods** and associated references are available in the online version of  
the paper at <http://www.nature.com/naturecellbiology/>.

18 **Supplementary Information** is linked to the online version of the paper at  
<http://www.nature.com/naturecellbiology/>.

## 20 **Acknowledgments**

We thank J. Deng for breeding fish, Drs. Peng Li and Shengcai Lin (China) for  
22 discussion. This work was supported by the Biomedical Research Council of

Singapore (07/1/21/19/493, R-08-1-21-19-585 & SBIC-SSCC C-002-2007) and  
2 the Ministry of Education of Singapore.

### Author Contributions

4 M.L., N.H., Z.L., M.Y., R.L., H.Z., Y.Y. and Y.Y performed research; M.L. and N.H.  
analyzed the data; C.L., J.S. and Y.H. designed research, R.G. and Y.H. wrote  
6 the paper.

**Author Information** Reprints and permissions information is available at

8 [www.nature.com/naturecellbiology/reprints](http://www.nature.com/naturecellbiology/reprints). Correspondence and requests for  
materials should be addressed to Y. H. ([dbshyh@nus.edu.sg](mailto:dbshyh@nus.edu.sg)) and J.S.  
10 ([jianxin\\_song@nuhs.edu.sg](mailto:jianxin_song@nuhs.edu.sg)).

1 Zernicka-Goetz, M., Morris, S. A. & Bruce, A. W. Making a firm decision:  
12 multifaceted regulation of cell fate in the early mouse embryo. *Nat Rev Genet* **10**,  
467-477 (2009).

14 2 Nakamura, S., Kobayashi, K., Nishimura, T., Higashijima, S. & Tanaka, M.  
Identification of germline stem cells in the ovary of the teleost medaka. *Science*  
16 **328**, 1561-1563 (2010).

3 Ohinata, Y. *et al.* A signaling principle for the specification of the germ cell  
18 lineage in mice. *Cell* **137**, 571-584 (2009).

4 Barker, N. *et al.* Lgr5(+ve) stem cells drive self-renewal in the stomach and build  
20 long-lived gastric units in vitro. *Cell Stem Cell* **6**, 25-36 (2010).

5 Ohinata, Y. *et al.* Blimp1 is a critical determinant of the germ cell lineage in mice.  
22 *Nature* **436**, 207-213 (2005).

6 Sato, T. *et al.* Single Lgr5 stem cells build crypt-villus structures in vitro without  
24 a mesenchymal niche. *Nature* **459**, 262-265 (2009).

7 Wittbrodt, J., Shima, A. & Scharl, M. Medaka--a model organism from the far  
26 East. *Nat Rev Genet* **3**, 53-64 (2002).

8 Hong, N. *et al.* Accessibility of host cell lineages to medaka stem cells depends on  
28 genetic background and irradiation of recipient embryos. *Cell Mol Life Sci* **67**,  
1189-1202 (2010).

9 Hong, Y. *et al.* Establishment of a normal medakafish spermatogonial cell line  
30 capable of sperm production in vitro. *Proc Natl Acad Sci U S A* **101**, 8011-8016  
32 (2004).

10 Hong, Y., Winkler, C. & Scharl, M. Pluripotency and differentiation of  
34 embryonic stem cell lines from the medakafish (*Oryzias latipes*). *Mech Dev* **60**,  
33-44 (1996).



- 11 Hong, Y., Winkler, C. & Schartl, M. Production of medakafish chimeras from a  
2 stable embryonic stem cell line. *Proc Natl Acad Sci U S A* **95**, 3679-3684 (1998).
- 12 Hong, Y., Winkler, C. & Schartl, M. Efficiency of cell culture derivation from  
4 blastula embryos and of chimera formation in the medaka (*Oryzias latipes*)  
6 depends on donor genotype and passage number. *Dev Genes Evol* **208**, 595-602  
(1998).
- 13 Yi, M., Hong, N. & Hong, Y. Generation of medaka fish haploid embryonic stem  
8 cells. *Science* **326**, 430-433 (2009).
- 14 Kane, D. A. & Kimmel, C. B. The zebrafish midblastula transition. *Development*  
10 **119**, 447-456 (1993).
- 15 Xu, H., Li, Z., Li, M., Wang, L. & Hong, Y. Boule is present in fish and  
12 bisexually expressed in adult and embryonic germ cells of medaka. *PLoS One* **4**,  
e6097 (2009).
- 14 16 Newport, J. & Kirschner, M. A major developmental transition in early *Xenopus*  
16 embryos: I. characterization and timing of cellular changes at the midblastula  
stage. *Cell* **30**, 675-686 (1982).
- 17 Iwamatsu, T. Stages of normal development in the medaka *Oryzias latipes*. *Mech*  
18 *Dev* **121**, 605-618 (2004).
- 18 Nikitina, N., Sauka-Spengler, T. & Bronner-Fraser, M. Chapter 1. Gene  
20 regulatory networks in neural crest development and evolution. *Curr Top Dev*  
*Biol* **86**, 1-14 (2009).
- 22 19 Kelsh, R. N. *et al.* The Tomita collection of medaka pigmentation mutants as a  
24 resource for understanding neural crest cell development. *Mech Dev* **121**, 841-859  
(2004).
- 20 Cavaleri, F. & Scholer, H. R. Nanog: a new recruit to the embryonic stem cell  
26 orchestra. *Cell* **113**, 551-552 (2003).
- 21 Nichols, J. *et al.* Formation of pluripotent stem cells in the mammalian embryo  
28 depends on the POU transcription factor Oct4. *Cell* **95**, 379-391 (1998).
- 22 Camp, E. *et al.* Nanog regulates proliferation during early fish development. *Stem*  
30 *Cells* **27**, 2081-2091 (2009).
- 23 Hong, Y., Winkler, C., Liu, T., Chai, G. & Schartl, M. Activation of the mouse  
32 Oct4 promoter in medaka embryonic stem cells and its use for ablation of  
spontaneous differentiation. *Mech Dev* **121**, 933-943 (2004).
- 34 24 Thermes, V. *et al.* Medaka simplet (FAM53B) belongs to a family of novel  
36 vertebrate genes controlling cell proliferation. *Development* **133**, 1881-1890  
(2006).
- 25 Niakan, K. K. *et al.* Sox17 promotes differentiation in mouse embryonic stem  
38 cells by directly regulating extraembryonic gene expression and indirectly  
antagonizing self-renewal. *Genes Dev* **24**, 312-326 (2010).
- 40 26 Bejar, J., Hong, Y. & Schartl, M. Mitf expression is sufficient to direct  
42 differentiation of medaka blastula derived stem cells to melanocytes.  
*Development* **130**, 6545-6553 (2003).
- 27 Hou, L., Arnheiter, H. & Pavan, W. J. Interspecies difference in the regulation of  
44 melanocyte development by SOX10 and MITF. *Proc Natl Acad Sci U S A* **103**,  
9081-9085 (2006).

- 28 Kobayashi, D., Jindo, T., Naruse, K. & Takeda, H. Development of the endoderm  
2 and gut in medaka, *Oryzias latipes*. *Dev Growth Differ* **48**, 283-295 (2006).
- 29 Rossant, J. & Tam, P. P. Blastocyst lineage formation, early embryonic  
4 asymmetries and axis patterning in the mouse. *Development* **136**, 701-713 (2009).
- 30 Guo, G. *et al.* Resolution of cell fate decisions revealed by single-cell gene  
6 expression analysis from zygote to blastocyst. *Dev Cell* **18**, 675-685 (2010).
- 31 Newport, J. & Kirschner, M. A major developmental transition in early *Xenopus*  
8 embryos: II. Control of the onset of transcription. *Cell* **30**, 687-696 (1982).
- 32 Carvalho, L. & Heisenberg, C. P. The yolk syncytial layer in early zebrafish  
10 development. *Trends Cell Biol*, doi:S0962-8924(10)00130-3 (2010).
- 33 Silva, J. *et al.* Nanog is the gateway to the pluripotent ground state. *Cell* **138**, 722-  
12 737 (2009).
- 34 Harris, M. L. & Erickson, C. A. Lineage specification in neural crest cell  
14 pathfinding. *Dev Dyn* **236**, 1-19 (2007).
- 35 Yi, M., Hong, N. & Hong, Y. Derivation and characterization of haploid  
16 embryonic stem cell cultures in medaka fish. *Nat Protoc* **5**, 1418-1430 (2010).
- 36 Li, M. *et al.* Medaka *vasa* is required for migration but not survival of primordial  
18 germ cells. *Mech Dev* **126**, 366-381 (2009).

## 20 **Figure legends**

### **Table 1 | Efficiency and differentiation of single blastomere culture**

22 **Figure 1 | Single cell culture.** (a-c) Merged micrographs of live embryos. The  
physiological integrity of the yolk membrane (ym) is evidenced by the prevented  
24 distribution of injected red fluorescent dye from yolk to the cell mass. Vg  
transgenic embryos were injected with red fluorescent dye at the positions  
26 (arrows) and photographed at 32-cell stage. The GFP signal is from maternal  
expression from the *vasa* promoter. (a and b) Lateral views following yolk  
28 injection 5 min after fertilization near the blastoderm (bd; a) or at the vegetal pole  
(b). (c) Top view of one-cell injection at the 2-cell stage, showing the restricted  
30 dye distribution into descendents of the injected blastomere. ch, chorion; od, oil  
droplet; yk, yolk. An embryo is 1 mm in diameter. (d) Schematic illustration of  
32 single cell isolation and culture. Shown here is a 16-cell embryo for manual  
dissociation, and cells are individually seeded into 96-well plate, one cell one well.

Cell behaviors are regularly monitored. (e) 16-cell blastoderm. (f) Isolated 16-cell  
2 blastomeres, showing differences in size and transparency. (g-j) Heterogeneity of  
16-cell blastomeres. Embryos were microinjected with H2B<sub>gfp</sub> RNA at the 1-cell  
4 stage for labeling nuclei (green). Three major classes of blastomeres are seen.  
Class I is yolky, actively forms pseudopodia (asterisks) without cell division (g).  
6 Class II undergoes endomitosis (see Supplementary Figure 3). Class III exhibits  
active divisions and produce daughter cells of heterogeneous sizes (h) and a  
8 homogeneous size (i and j) often capable of attachment (arrow) until day 1 of  
culture. Embryos are 1 mm in diameter. Scale bars, 50 μm.

10 **Figure 2 | Lineage tracing in vitro.** (a) Medaka blastula lineages and mouse  
blastocyst lineages. (b) Early medaka blastula following zygotic injection of  
12 H2B<sub>gfp</sub> RNA, showing YSL nuclei (green). Oil droplets (od) depict the vegetal  
pole. (c) YSL precursor at 4 h of culture. (d) Developing YSL cells at 6 h of  
14 culture. (e) YSL cells at day 1 of culture. Three nuclei (nu) are seen in a large-  
sized cytoplasmic syncytium. (f) Control YSL cells from midblastula embryos at  
16 day 3 of culture. (g) ES cells and black pigmented melanocytes (mc) at day 3. (h)  
Yellow fluorescent iridocytes (ic). Scale bars, 50 μm.

18 **Figure 3 | RNA Expression of key lineage regulators.** After whole mount in  
situ hybridization with antisense RNA probes, the signals were detected by green  
20 (*oct4* and *mitf1*) and red fluorescence (*nanog* and *sox17*), and the blastoderm  
were separated from the yolk for photography. Nuclei were stained blue with  
22 DAPI. (a) *oct4* and *nanog* expression. (b-d) *mit1* and *sox17* expression at the  
16~32-cell stage (b), 32-cell stage (c) and 64-cell stage (d). Scale bars, 50 μm.

**Table 1 | Efficiency and differentiation of single blastomere culture<sup>1)</sup>**

Stage	Blastomere, n (%)		Class, n (%) <sup>3)</sup>			Daughter cells, n(%) <sup>4)</sup>		
	Seeded	Survived <sup>2)</sup>	I	II	III	ES	Pig	YSL
16-cell	194	50 (26±10.7)	37 (74±12.9)	3 (6±3.9)	10 (26±10.9)	3 (6)	3 (6)	3 (6)
32-cell	206	100 (49±14.5)	50 (50±12.3)	4 (4±2.3)	46 (46±12.3)	5 (5)	4 (4)	4 (4)
64-cell	224	150 (67±12.4)	77 (51±16.7)	6 (4±4.8)	67 (45±13.8)	9 (6)	6 (4)	6 (4)

<sup>1)</sup> HB32 embryos with and without injection with H2B<sub>gfp</sub> RNA at the 1-cell stage were dissociated for single cell culture at 16-, 32- and 64-cell stages. Data are presented as means ± s.d. from four independent experiments.

<sup>2)</sup> Survival rate at 1 day post culture was derived by comparison to the number of cells seeded.

<sup>3)</sup> Classes I, II and III are non-dividing, endomitotic and dividing blastomeres until 1 day post culture. Percentages are derived by comparisons to the number of cells survived.

<sup>4)</sup> Blastomeres that produced daughter cells containing embryonic stem (ES) cells, yolk syncytial layer (YSL) cells, and pigment cells (Pig; melanocytes and/or iridocytes) at 1-4 dpc. Percentages were derived by comparisons to the number of cells survived.





Figure 1 | Single cell culture.



**Figure 2 | Lineage tracing in vitro.**




Figure 3 | RNA Expression of lineage markers.