Can Plant-Made Copper Chaperones Heal Early Alzheimer's Disease?

^{a*}Bernd Kastenholz, ^bBasil Horst, and ^cJürgen Horst

BK, BH and JH equally contributed to this paper.

1 **Institute of origin:**

2 ^aIBG-2: Plant Sciences, Research Centre Juelich, 52425 Juelich, Germany

3 ^bColumbia University, Department of Dermatology, New York, NY 10032, USA

4 ^cWestfälische Wilhelms-Universität Münster, Institut für Humangenetik, 48149 Münster,
5 Germany

6

7 **Corresponding author:**

8 Bernd Kastenholz, ^aIBG-2: Plant Sciences, Research Centre Juelich, 52425 Juelich, Germany

9 Tel.: +49 (0) 2461 614825;

10 Fax: +49 (0) 2461 612492;

11 E-mail: b.kastenholz@fz-juelich.de

12

13 **Running title:**

14 Phytopharmaceuticals for restoring biometal homeostasis in human cells

15

16 **Key words:**

17 Alzheimer's disease (AD), recombinant plant-made pharmaceuticals (PMPs), CCS, *Ginkgo*
18 *biloba*, SOD, A β , molecular chaperones

19

20 **Abbreviations:**

21 CCS: copper chaperone for superoxide dismutase

22 SOD-1: Cu,Zn-superoxide dismutase

23 A β : amyloid β

24 AD: Alzheimer's disease

25 ROS: reactive oxygen species

26 PMPs: plant-made pharmaceuticals

Abstract:

Therapeutic recombinant plant-made copper chaperone for superoxide dismutase (CCS) derived from *Ginkgo biloba* leaves may establish and maintain physiologic Cu levels through restoration and modulation of biometal metabolism in organ systems of younger Alzheimer patients (> 50 years). Medications developed from plant-made copper chaperone proteins may delay progression during early disease stages or even be a basis for a possible causal treatment of preclinical stages of Alzheimer's disease by restoring cellular function of the CCS-SOD-1 mechanism and by preventing formation of A β plaques, a major putative factor involved in Alzheimer's disease pathogenesis.

1 INTRODUCTION

2 The dysregulation of biometal (Cu, Zn, Fe) homeostasis and oxidative stress in brain cells
 3 have been found to impact on accumulation of amyloid β ($A\beta$), a major putative factor
 4 involved in early Alzheimer's disease (AD) pathogenesis [1]. The regulation of metal ion
 5 homeostasis in the cytoplasm is strongly influenced by copper chaperone for superoxide
 6 dismutase (CCS) and Cu,Zn-superoxide dismutase (SOD-1) [2]. The dynamic interplay of
 7 properly folded CCS and SOD-1 guarantees that free Cu and Zn ions are being complexed by
 8 these metal proteins and do not catalyze oxidation processes of proteins, lipids, DNA, and
 9 other molecules in the cells [2]. When these physiological complexation mechanisms do not
 10 function properly, oxidative stress and dys-homeostasis of Cu and Zn metabolism may give
 11 rise to misfolding, accumulation and aggregation of amyloid β peptides [2]. The outcome of
 12 these pathological processes may lead to incurable chronically progressive neurodegenerative
 13 diseases such as Alzheimer's disease [1,2]. Several therapeutic strategies and nearly all
 14 medications used or suggested as $A\beta$ inhibitors, including metal-chelating agents or radical
 15 scavengers, at present time, aim at the treatment of AD symptoms only and may either be
 16 toxic, lack specificity or have unknown mechanisms of action *in vivo* [2,3].

17
 18 The aim of this article is to give a short review on the interaction of metal ions with novel and
 19 early herbal compounds derived from *Ginkgo biloba*, and their possible role in the treatment
 20 of early Alzheimer's disease. Though *Ginkgo biloba* leaf extracts are generally administered
 21 to treat dementia syndroms in older AD patients (> 65 years old), no data is available on
 22 possible effects of recombinant plant-made copper chaperones from *Ginkgo biloba* in younger
 23 AD patients (> 50 years old) with preclinical stages of disease.

25 INTERACTION OF HERBAL DRUGS WITH METAL IONS

26 The medical community, including the pharmaceutical industry, as well as AD patients, have
 27 become aware of the well-known antioxidant effects of ancient herbal medications, namely
 28 *Ginkgo biloba* leaf extracts [4]. A complex mixture of flavonoids and terpenoids are thought
 29 to represent the main bioactive compounds of these plant extracts [5]. For example, the
 30 flavonol quercetin, a major compound of *Ginkgo* extract, was found to interact with Cu^{2+} and
 31 Fe^{3+} [6]. However, an ideal therapeutic drug to dissolve $A\beta$ peptides would involve a
 32 compound selective for Cu^+ , Zn^{2+} and Fe^{3+} [7]. A meta-analysis by Birks and Evans revealed
 33 that commercially available medicinal plant extracts (EGb 761[®] *Ginkgo biloba* leaf extract)
 34 have no consistent pattern of any clinical benefit associated with *Ginkgo biloba* for people

with dementia or cognitive impairment [4]. Furthermore, the results from the study of He and colleagues suggested that high doses of herbal remedies can even be toxic to cells [8]. Thus, *Ginkgo biloba* extracts may induce unwanted side-effects and may also lack specificity as to the binding of Cu^+ and Zn^{2+} in the cytoplasm. Likely, these extracts neither reduce metal-based oxidative stress efficiently nor contribute to the homeostatic control of biometals in human cells, though *in vitro* studies have shown that ginkgolides may protect against the synapse damage and cognitive loss seen during the early stages of AD [5]. However, medicinal plants may contain other, more efficient bioactive molecules apart from the well-known flavonoids and terpenoids, namely metallochaperones [9].

EFFICACY OF PLANT-MADE COPPER CHAPERONES

As a basis for a new drug development involving metal-chelating agents, the following facts may be important. Endogenous biomolecules such as Cu,Zn-superoxide dismutase, are one of the major means by which cells counteract the deleterious effects of reactive oxygen species (ROS). For proper functioning SOD-1 has to be activated by the metallochaperone protein, copper chaperone for SOD [9-12]. Copper ions are required for enzymatic activity whereas the zinc ion helps to stabilize the enzyme [10]. Studies by Choi and colleagues suggest that recombinant human CCS molecules produced in bacteria provide a potential strategy for therapeutic delivery of these compounds in various human diseases related to ROS and SOD [10]. Drugs derived from recombinant proteins potentially have greater efficacy and fewer side-effects than small organic molecules (e.g., Cu orotate, quercetin), because their action can be more precisely targeted towards the A β plaque formation as a major putative factor in the pathogenesis of Alzheimer's disease rather than the treatment of AD symptoms [11]. The dysfunction of the CCS-SOD-1 interplay may be one specific but major mechanism in the pathogenesis of Alzheimer's disease. It is a well-known fact that biometals (Fe, Zn, Cu) are accumulated in the brain with normal ageing [13]. Important factors affecting the balance between metal ion accumulation and deficiency are, for example, environmental exposure, ageing or drug interaction [14]. The inability of the human organism to maintain the metal ion homeostasis due to improperly folded CCS and SOD molecules in brain cells is suggested as causal for preclinical stages, development and progression of AD and other neurodegenerative diseases [9,11,15].

For the restoration and modulation of metal ion homeostasis in the treatment of AD, we have proposed a novel class of pharmacologically active plant ingredients as antioxidants: copper

chaperone for superoxide dismutase derived from medicinal plants (e.g., *Ginkgo biloba*) [9,11]. In molecular farming approaches transgenic plants (e.g., tobacco) may serve as an efficient production platform for medications in regard to protein yield, quality and stability [16]. Recombinant CCS proteins produced in transgenic plants may cross the blood-brain barrier and are relatively free from side-effects [9,11]. Being properly-folded, plant-made CCS may have the ability to bind and deliver $\text{Cu}^+/\text{Cu}^{2+}$, Zn^{2+} and Fe^{3+} ions and to normalize the SOD-1 activity via specific protein-protein interactions in the central nervous system and peripherally [9,11].

In contrast to plant-made pharmaceuticals (PMPs), organisms like yeast, mammals or bacteria used for molecular farming approaches, express properly folded as well as improperly folded recombinant therapeutic proteins. These medications may lack stability and pharmacological efficiency in protein-misfolding diseases, e.g., Alzheimer's disease [16]. Furthermore, PMPs may have several advantages in terms of cost scalability or safety issues compared to the other genetically modified organisms [16].

Because of their specific biochemical behavior, plant-made CCS from medicinal plants may be efficient in the treatment of patients with preclinical stages of AD. The endogenous levels of this essential copper protein may be important, since a mild copper deficiency has been described in AD patients [17]. Furthermore, the expression level of CCS has been found to reflect the Cu status of patients and thus, may serve as a marker for *in vivo* copper levels [18]. The metal levels in patients with Alzheimer's disease compared with healthy individuals are important parameters for developing drugs that may restore the intracellular metal ion metabolism [14]. It is anticipated that plant-made CCS may establish physiologic copper levels through restoration and modulation of biometal metabolism in diseased organ systems of AD patients [9,11]. Yet, to our knowledge, no systematic studies have addressed the possible role of plant-made CCS in copper homeostasis.

The effects of recombinant CCS and a possible role in the treatment of AD can be characterized as follows: unfolded SOD-1 in diseased blood involving denatured proteins and peptides [19], may be activated by copper ion incorporation via specific CCS-SOD interaction [9,11]. Copper-demetalated CCS and Cu cofactor-containing CCS complexes may pass through the blood-brain barrier and activate unfolded SOD-1 by Cu ion transfer or bind free Cu, Zn and Fe ions in brain cells, respectively [9,11]. Since copper homeostasis is regarded as

an important possible factor in the complex pathogenesis of AD [20] restoring homeostasis of Cu metabolism may positively affect early disease stages of AD patients by balancing oxidative and anti-oxidative processes and by reducing protein-misfolding processes leading to formation of A β plaque deposits [9, 11, 20]. The interactions of plant-derived CCS medications with unfolded human SOD-1 and free metal ions (Cu, Zn, Fe) may trigger a cascade of other biochemical reactions, such as the degradation of A β plaque deposits by molecular chaperones and the ubiquitin proteasome system [21].

Further studies to determine the dose, bioavailability and mechanisms of action of plant-made pharmaceuticals seem warranted in younger AD patients (> 50 years) as previously proposed [9, 11].

CONCLUSION

For the restoration and modulation of metal ion homeostasis and for balancing intracellular pro-oxidative and antioxidative processes in the treatment of Alzheimer's disease, plant-made copper chaperone for superoxide dismutase (CCS) proteins potentially have greater efficacy and fewer side-effects compared to small organic molecules (e.g., quercetin) from medicinal plant extracts or therapeutic recombinant proteins produced in bacteria, fungi or mammals. Possibly suitable for use in younger AD patients (> 50 years), CCS proteins derived from medicinal plants may be targeted more towards prevention of A β plaque formation as a major putative factor involved in AD pathogenesis, rather than the treatment of AD symptoms in older patients (> 65 years). We suggest that recombinant plant-made CCS derived from *Ginkgo biloba* leaves might be promising in the treatment of patients suffering from preclinical symptoms of Alzheimer's disease.

REFERENCES

1. Wesson, D.W.; Nixon, R.A.; Levy, E.; Wilson, D.A. Mechanisms of Neural and Behavioral Dysfunction in Alzheimer's Disease. *Mol. Neurobiol.*, **2011**, in press.
2. Crichton, R.R.; Ward, R.J. *Metal-Based Neurodegeneration: From Molecular Mechanisms to Therapeutic Strategies*, 1st ed.; John Wiley & Sons Ltd.: Chichester, England, **2006**.
3. Estrada, L.D.; Yowtak, J.; Soto, C. Protein misfolding disorders and rational design of antimisfolding agents. *Methods Mol. Biol.*, **2006**, *340*, 277-293.

- 1 4. Birks, J.; Evans, J.G. Ginkgo biloba for cognitive impairment and dementia. *Cochrane*
2 *Database Syst. Rev.*, **2009**, DOI:10.1002/14651858.CD003120.pub3.
- 3 5. Bate, C.; Tayebi, M.; Williams, A. Ginkgolides protect against amyloid-beta (1-42)-
4 mediated synapse damage in vitro. *Mol. Neurodegen.*, **2008**, 3, Article No.: 1.
- 5 6. Kuo, S.-M.; Leavitt, P.S.; Lin, C.-P. Dietary flavonoids interact with trace metals and
6 affect metallothionein level in human intestinal cells. *Biol. Trace Elem. Res.*, **1998**,
7 62(3), 135-153.
- 8 7. Cuajungco, M.P.; Faget, K.Y.; Huang, X.; Tanzi, R.E.; Bush A.I. Metal chelation as a
9 potential therapy for Alzheimer's disease. *Ann. N.Y. Acad. Sci.*, **2000**, 920, 292-304.
- 10 8. He, J.; Lin, J.; Li, J.; Zhang, J.-H.; Sun, X.-M.; Zeng, C.-M. Dual effects of Ginkgo
11 biloba leaf extract on human red blood cells. *Basic Clin. Pharmacol. Toxicol.*, **2009**,
12 104(2), 138-144.
- 13 9. Kastenholtz, B.; Garfin, D.E. Medicinal plants: a natural chaperones source for treating
14 neurological disorders. *Protein Pept. Lett.*, **2009**, 16(2), 116-120.
- 15 10. Choi, S.H.; Kim, D.W.; Kim, S.Y.; An, J.J.; Lee, S.H.; Choi, H.S.; Sohn, E.J.; Hwang,
16 S.-I.; Won, M.H.; Kang, T.-C.; Kwon, H.J.; Kang, J.H.; Cho, S.-W.; Park, J.; Eum,
17 W.S.; Choi, S.Y. Transduced human copper chaperone for Cu,Zn-SOD (PEP-1-CCS)
18 protects against neuronal cell death. *Mol. Cells*, **2005**, 20(3), 401-408.
- 19 11. Kastenholtz, B.; Garfin, D.E.; Horst, J.; Nagel, K.A. Plant metal chaperones: a novel
20 perspective in dementia therapy. *Amyloid J. Prot. Fold. Disord.*, **2009**, 16(2), 81-83.
- 21 12. Sekhon, B.S. Metallochaperones - an overview. *Curr. Chem. Biol.*, **2010**, 4(2), 173-
22 186.
- 23 13. Smith, D.G.; Cappai, R.; Barnham, K.J. The redox chemistry of the Alzheimer's
24 disease amyloid beta peptide. *Biochim. Biophys. Acta - Biomembr.*, **2007**, 1768(8),
25 1976-1990.
- 26 14. Zatta, P.; Drago, D.; Bolognin, S.; Sensi S.L. Alzheimer's disease, metal ions and
27 metal homeostatic therapy. *Trends Pharmacol. Sci.*, **2009**, 30(7), 346-355.
- 28 15. Bolognin, S.; Messori, L.; Zatta, P. Metal ion physiopathology in neurodegenerative
29 disorders. *Neuromol. Med.*, **2009**, 11(4), 223-238.
- 30 16. Nagel, K.A.; Kastenholtz, B.; Gilmer, F.; Schurr, U.; Walter, A. Novel detection
31 system for plant protein production of pharmaceuticals and impact on conformational
32 diseases. *Protein Pept. Lett.*, **2010**, 17(6), 723-731.
- 33 17. Pajonk, F.-G.; Kessler, H.; Supprian, T.; Hamzei, P.; Bach, D.; Schweickhardt, J.;
34 Herrmann, W.; Obeid, R.; Simons, A.; Falkai, P.; Multhaup, G.; Bayer, T.A. Cognitive

- decline correlates with low plasma concentrations of copper in patients with mild to moderate Alzheimer's disease. *J. Alz. Dis.*, **2005**, 8(1), 23-27.
18. Wang, T.; Guo, Z.J. Copper in medicine: homeostasis, chelation therapy and antitumor drug design. *Curr. Med. Chem.*, **2006**, 13(5), 525-537.
19. Kang, M.N.; Park, W.; Kang, S.M.; Lim, K.; Lee, B.; Kim, G.J.; Kim, S.Y.; Yang, Y.S.; Na, H.R.; Youn, Y.C.; An, S.S.A.; Ju, Y.R.; Kim, S. Presence of amyloid-beta (A β) oligomers in blood from patients with Alzheimers disease. *Prion*, **2010**, 4(3), 224-224.
20. Lin, C.-J., Huang, H.-C., Jiang, Z.-F. Cu(II) interaction with amyloid- β peptide: a review of neuroactive mechanisms in AD brains. *Brain Res. Bull.*, **2010**, 82(5-6), 235-242.
21. Luo, G.R.; Le, W.D. Collective roles of molecular chaperones in protein degradation pathways associated with neurodegenerative diseases. *Curr. Pharm. Biotechnol.*, **2010**, 11(2), 180-187.