

Standard operating procedures (SOP) in experimental stroke research:

SOP for middle cerebral artery occlusion in the mouse

Ulrich Dirnagl, and the members of the MCAO-SOP group

Dept. of Experimental Neurology and Center for Stroke Research (CSB),

Charite University Medicine Berlin, Germany

Address correspondence to:

Prof.Dr. Ulrich Dirnagl

Dept. of Experimental Neurology and Center for Stroke Research (CSB)

Charite University Medicine Berlin

Charitepl.1

10098 Berlin Germany

Tel. +49-30-45056013

ulrich.dirnagl@charite.de

Members of the MCAO-SOP group: Odilo Engel, Dr. Tracy Farr, Dr. Karen Gertz, Dr.

Denise Harhausen, Prof. Dr. Ute Lindauer, Dr. Vincent Prinz, PD Dr. Andre Rex

Acknowledgements: Funded by the Bundesministerium für Bildung und Forschung (Center for Stroke Research Berlin and the European Community's Seventh Framework Programme (FP7/2007-2013) under grant agreement n° 201024 and n° 202213 (European Stroke Network) .

Thus far, the translation of promising results from preclinical stroke research into effective clinical therapy has not met with success (1). Among the numerous possible reasons for this failure, quality problems in some of the basic research or preclinical studies have to be considered. False-positive results, inflated effect sizes, and marginal reproducibility may have overestimated or even affected the potential of novel stroke therapeutics (2). Systematic reviews have found quantitative evidence that low study quality may have introduced a bias into preclinical stroke research (3,4,5). As opposed to many other causes of the 'translational roadblock', study quality is fully under the control of the researcher, and thus a prime target for improvement. Increasingly, funding bodies and review boards overseeing animal experiments are taking a proactive stance, and demand auditable measures of quality control in preclinical research (6). The Stroke Therapy Academic Industry Roundtable (STAIR) recently updated its recommendations for the evaluation of preclinical data on neuroprotective drugs (7) to include good laboratory practice (GLP) issues (8).

Monitoring, auditing, and standard operating procedures (SOPs) are key elements of quality control in randomized clinical trials (RCTs). It has been proposed that experimental stroke research adapt some of the tools used in clinical stroke research. In particular, stroke laboratories should set up and publish their SOPs (e.g., on their institutional websites), and guarantee that their studies adhere to these standards (9). This is all the more important, as a certain portion of their experiments, evaluations, etc. are not performed by professionals, but rather by students in training who are unaware of these issues.

A standard operating procedure (SOP) is a set of instructions with the force of a directive covering those features of operations that lend themselves to a definite or standardized procedure without loss of effectiveness. The primary purpose of an SOP in experimental stroke research is to guide and standardize working procedures in order to ensure data reliability and integrity. It is crucial that researchers, students and technicians read and follow

the SOPs. If this is not the case, SOPs will not only fail to achieve their goal; they will also engender a false sense of security. Failures are often due to technical shortcomings in the SOPs themselves. SOPs should be written by the user, as they must convey a clear instruction. The user must not only understand the instruction but also be prepared to carry it out.

We in the following introduce for the first time an SOP in experimental stroke research.

Write down what you do, do what is written down!

SOP for middle cerebral artery occlusion (MCAO)

Title	Middle cerebral artery occlusion (MCAO) in the mouse (intraluminal suture)
Date	Version 4: 3.4.2012
Version History	Version 1: 28.07.2009 Version 2: 17.8.2010 Version 3: 12.7.2011
Name of Author	Vincent Prinz, Janett König, Shengbo Ji, Ute Lindauer, Andre Rex, Ulrich Dirnagl
Purpose	Experimental induction of focal cerebral ischemia after occlusion of the middle cerebral artery
Scope and Applicability	Applies to a procedure in a standard lab equipped and certified for in vivo experimentation in rodents (including anesthesia with volatile anesthetics). Experimental procedures require approval by the relevant committees (in Germany: <i>Tierversuchsgenehmigung, LAGeSo</i>)
Introduction	Experimental focal ischemia is most commonly studied after permanent or transient occlusion of the middle cerebral artery (MCA) in rodents. Proximal MCA occlusion can be induced by an intraluminal suture (so-called filament model) and causes injury to cortex and deeper brain structures (striatum). Distal MCA occlusion (the so-called 'Brint' or 'Tamura'-models) is usually produced by placing a vascular clip on a pial vessel or by cautery. Distal occlusion typically spares the striatum and primarily involves the neocortex. Pannecrosis develops in

the territory supplied by the respective artery with glial and endothelial cell death. If recirculation is established early (2 hrs or less) outcome is better (transient MCA occlusion). In some ways, the reperfused brain imitates restoration of blood flow after spontaneous lysis of a thromboembolic clot in humans, even though reperfusion after clot lysis is certainly more complex than an on-off phenomenon as modeled by placement and retraction of an intravascular filament.

This SOP describes a mouse model of transient proximal MCAO (30 – 60 minutes occlusion time).

References

Carmichael, S.T. (2005) Rodent models of focal stroke: size, mechanism, and purpose *NeuroRx*. **2(3)**, 396-409. Review.

Dirnagl, U. (2006) Bench to bedside: The quest for quality in experimental stroke research *J Cereb Blood Flow Metab* **26**, 1465-78.

Dirnagl, U., Iadecola, C., Moskowitz, M.A. (1999) Pathobiology of ischaemic stroke: an integrated view *Trends Neurosci.* 22(9), 391-7.

Engel O, Kolodziej S, Dirnagl U, Prinz V (2011) Modeling Stroke in Mice - Middle Cerebral Artery Occlusion with the Filament Model. *J Vis Exp*:e2423

Shah, Z.A., Namiranian, K., Klaus, J., Kibler, K., Dore, S. (2006) Use of an optimized transient occlusion of the middle cerebral artery protocol for the mouse stroke model *J Stroke Cerebrovasc Dis.* **15**, 133-8.

Materials and Supplies

- filament USP 4/0 or 6/0 Suprama
- 8.0 nylon monofilament for coating with Xantopren M and Activator Universal Plus or commercially available filaments (e.g. Docol)
- Xantopren M Mucosa (Heraeus Kulzer)
- Activator Universal Plus (Heraeus Kulzer, for Xantropren)

	<ul style="list-style-type: none"> • surgical needle and thread for suture
Instrumentation	<ul style="list-style-type: none"> • dissecting microscope (max. x 40) • temperature feedback controlled heating plate • surgical instruments <ul style="list-style-type: none"> • forceps (Dumont Medical #5 straight tip 0,05mm x 0,02 mm) • surgical scissors (skin cut) • forceps for skin handling and suture (e.g. standard anatomical) • vascular spring scissors (Vannas) • 2 hemostats (Hartmann) • Micro vascular clamp (e.g. S&G B1-V) and applying forceps • Needle holder (Olsen-Hegar or other) • anesthesia system for isoflurane and nitrous oxide • heated recovery cage
Cautions	<p>Maintain a body temperature of 36.5 +/- 0.5 °C also after reperfusion (for 2 hours). See also Appendix below.</p> <p>Ensure proper pain relief in the perioperative and postoperative period, e.g. by repeated topical application of a long-acting local anaesthetic, like bupivacaine ointment serving as an absorption depot.</p> <p>Surgical procedures should be carried out under clean conditions. (Sterile surgical instruments and materials, clean gown, gloves, etc.). See also Appendix below.</p>
Personnel	<i>In general:</i>
Qualifications	Surgeons need:

- general supervision and instruction,
- the appropriate certification according to FELASA (A/B),
- official registration.

Also required in Germany:

- official registration (*Personenanzeige*)
- in Berlin: *Landesamt für Gesundheit und Soziales LAGeSo*,

For internal use:

- If you have questions contact Andre Rex (andre.rex@charite.de)
- <http://141.42.165.178/expneurointern/anweisungen/einweischaemie.html>
- In the initial training phase, review our video demonstration of the MCAO procedure

New surgeons must have passed practical qualification (see Appendix below)

<p>Names of SOP</p> <p>Reviewers</p>	<p>Ulrich Dirnagl (ulrich.dirnagl@charite.de)</p> <p>Berlin, 02.04.2012</p>
<p>Protocol</p>	<ol style="list-style-type: none"> 1. Mice are anaesthetized with 1.5% Isoflurane and maintained in 1.0% Isoflurane with 2/3 N₂O and 1/3 O₂ using a vaporizer. 2. Disinfect the skin of the ventral neck with an alcohol based skin disinfectant (e.g. Octenisept). 3. A midline neck incision is made and the soft tissues are pulled apart. 4. The left common carotid artery (LCCA) is carefully dissected free from the surrounding nerves (without harming the vagus nerve) and a ligature is made using 6.0/7.0 string.

5. Then the left external carotid artery (LECA) is separated and a loose thread (6.0/7.0) is looped around it above the occipital artery bifurcation and secured externally with a pair of hemostats.
6. Next, the left internal carotid artery (LICA) is isolated and a knot is prepared with a 6.0 filament.
7. After obtaining good view of the left internal cerebral artery (LICA) and the left pterygopalatine artery (LPA), the LICA is clipped.
8. A small hole is cut in the LCCA before it bifurcates to the LECA and the LICA. A monofilament made of 8.0 nylon coated with silicon hardener mixture (see below) is then introduced into the artery.
9. The clipped arteries are opened while the filament is inserted about 9 mm into the LICA to occlude LMCA.
10. The third knot on the LICA is closed to fix the filament in position and the thread around the LECA is removed.
11. For pain relief, Bupivacaine gel is topically applied in the wound, and the wound is temporally closed with a Michel suture clip or with an adaptive suture. The mice receive saline 0.5 ml subcutaneously as volume replenishment.
12. The mice are allowed to recover in the heated recovery cage for the duration of the occlusion.
13. After X min/hours occlusion, the mice are re-anaesthetized and the third knot is opened and the filament withdrawn (if reperfusion is intended)
14. All animals receive second volume replenishment as described above.
15. The remaining sutures are cut and the skin is adapted with a surgical suture
16. The body temperature of the mice during surgery is maintained at $36.5^{\circ}\text{C} \pm$

0.5°C using a heating plate.

17. I think it is very, very important to release the ligature on the LECA..

For sham operations the filament is inserted to occlude LMCA and withdrawn immediately to allow instant reperfusion (8.). The subsequent operation is identical to the animals undergoing cerebral ischemia (9.-14.).

A video is available which demonstrates the above described procedure. (see references)

Monofilament construction:

- 8.0 nylon filament is cut into pieces of 11 mm length under the microscope
- the filament tip must be coated over a distance of 8 mm completely and evenly with a hardener mixture of Xantopren M Mucosa and Activator Universal Plus

Appendix to SOP:

1. Entry qualification experiment for mouse MCAO surgeons
2. Randomized selection of animals from cage and concealment of treatment allocation
 - 2 a. Pharmacological study
 - 2 b. Genetically manipulated animals
3. Temperature control
4. Outcome assessment
5. Physiological parameters
6. Quasi-sterile surgery

1. Entry qualification experiment for mouse MCAO surgeons:

New surgeons need to demonstrate in a series of experiments that they perform the MCAO-operation within 15 minutes. Reproducibility is verified by induction of a certain infarct volume within a standard deviation of 40%. Mortality must not exceed 10% within 24 hours.

2. Randomized selection of animals from cage and concealment of treatment allocation

2 a. Pharmacological study:

Animals in cage are marked with bar/dot code at the beginning of the procedure.

Computer program (random number generator) selects animal, and assigns it to concealed treatment arm ('A', 'B', etc.).

Stock solution or pharmaceutical ready for application is prepared by assistant and randomly assigned code ('A', 'B', etc.).

2 b. Genetically manipulated animals:

Animals in both cages (e.g. knockout / wildtype) are marked with bar/dot code at the beginning of the procedure. Computer program (random number generator) selects animal and assigns it to concealed experimental arm ('A', 'B', etc.). Blinded intervention whenever possible.

3. Temperature control:

The body temperature of mice during surgery is maintained at $36.5^{\circ}\text{C} \pm 0.5^{\circ}\text{C}$ using a temperature controlled heating plate. Maintain a body temperature of $36.5 \pm 0.5^{\circ}\text{C}$ also after reperfusion (for 2 hours) using a heated recovery box.

4. Outcome assessment

Infarct volume should be evaluated blinded. Functional outcome (*more than Bederson Score!*) should be assessed as well. Mortality and exclusion of animals have to be reported, including specific causes for exclusion.

Exclusion criteria:

- no stroke,
- Problems during induction of MCAo (excessive bleeding, prolonged operation time \geq 15 min, thread placement).

CAVE:

Especially in genetically manipulated animals, be aware of vascular alterations, which might directly affect stroke outcome

5. Physiological parameters:

MABP, HR, blood gases, CBF should be measured in selected animals.

6. Quasi-sterile surgery

Prior to surgery, the surgeon has to scrub his hands. It is advisable to wear clean gown, and non-sterile gloves at all times the animal is being handled. The surgeon has to wear a clean gown, cap and mask during surgery. Surgical gloves ought to be worn. If gloves cannot be used, a surgical hand scrub from tips to elbows must precede every operation. The necessary components of aseptic techniques in rodents include also sterile instruments, and separate surgical and animal prep areas. The use of glass bead sterilizer for re-sterilization of instruments during for repetitive procedures is recommended.

- All instruments used must be sterilized prior to each group of surgeries.
- Instruments must be kept on sterile non-porous drapes during use.
- Separate instruments should be used for skin and tissue handling.
- Instruments must be cleaned of blood and debris by brushing or wiping with sterile water or saline and sterile gauze sponges between surgeries. (Best: distilled water)
- If contamination has occurred, instruments must be placed in 70% ethanol or a glass bead sterilizer for the appropriate period of time for the method used to be effective (or the instrument pack replaced by a new sterile instrument pack) between animals.
- If 70% ethanol is used, instruments must be rinsed with sterile water or saline before being used on the next animal.
- Surgical gloves and blades should be changed after contamination.
- Following surgery all instruments must be thoroughly cleaned and rinsed.

7. Postoperative care:

After surgery the animals must be weighted and checked daily for signs of discomfort.

The mice could show some weight loss post-surgery. To avoid dehydration the mice receive daily s.c. injections as volume replenishment. The mice receive mashed food in a petri-dish placed on the floor to encourage eating. The food is replaced daily for seven days.

References:

1. Endres, M., Engelhardt, B., Koistinaho, J., Lindvall, O., Meairs, S., Mohr, J.P., Planas, A., Rothwell, N., Schwaninger, M., Schwab, M.E., Vivien, D., Wieloch, T., Dirnagl, U. (2008) Improving outcome after stroke: overcoming the translational roadblock *Cerebrovasc Dis.* **25**(3), 268-78.
2. Dirnagl, U., Macleod, M.R. (2009) Stroke research at a road block: The streets from adversity should be paved with meta-analysis and good laboratory practice *Br J Pharmacol* (in press).
3. Bath, P.M.W., Gray, L.J., Bath, A.J.G., Buchan, A., Miyata, T., Green, A.R., on behalf of the NXY-059 Efficacy Meta-analysis in individual Animals with Stroke (NEMAS) investigators (2009) Effects of NXY-059 in experimental stroke: an individual animal meta-analysis *Br J Pharmacol* (in press).
4. Crossley, N.A., Sena, E., Goehler, J., Horn, J., van der Worp, B., Bath, P.M., Macleod, M., Dirnagl, U. (2008) Empirical evidence of bias in the design of experimental stroke studies: a meta-epidemiological approach *Stroke* **39**(3), 929-34.

5. Macleod, M.R., van der Worp, H.B., Sena, E.S., Howells, D.W., Dirnagl, U., Donnan, G.A. (2008) Evidence for the efficacy of NXY-059 in experimental focal cerebral ischaemia is confounded by study quality *Stroke* **39**, 2824-9.
6. Macleod, M.R., Fisher, M., O'Collins, V., Sena, E.S., Dirnagl, U., Bath, P.M., Buchan, A., van der Worp, H.B., Traystman, R., Minematsu, K., Donnan, G.A., Howells, D.W. (2009) Good Laboratory Practice: Preventing Introduction of Bias at the Bench *Stroke* **40**, e50-2, published as reprint in *J Cereb Blood Flow Metab* **29**, 221-3.
7. Stroke therapy academic industry roundtable (Fisher, M., Chair) (1999) Recommendations for standards regarding preclinical neuroprotective and restorative drug development *Stroke* **30**, 2752-8.
8. Fisher, M., Feuerstein, G., Howells, D.G., Hurn, P.D., Kent, TA., Savitz, S.I., Lo, E. (2009) Update of the Stroke Therapy Academic Industry Roundtable (STAIR) preclinical recommendations *Stroke* **40**, 2244-50.
9. Dirnagl, U. (2006) Bench to bedside: The quest for quality in experimental stroke research *J Cereb Blood Flow Metab* **26**, 1465-78.