

LEUKAEMIA

Fusion cooking

The mixed lineage leukaemia (*MLL*) gene can be fused to one of 30 different gene partners as a result of translocations involving the long arm of chromosome 11, but it is unclear whether this mutation alone can induce acute myeloid leukaemia (AML). Ono and colleagues now suggest that the leukaemogenesis that is mediated by *MLL* fusion proteins is a multistep process.

MLL is a nuclear protein with domains that implicate it as a transcriptional regulator as well as being involved in chromatin-modifying supercomplexes. Although diverse, the *MLL* fusion proteins are all oncogenic, apparently promoting leukaemia through similar mechanisms — the aberrant activation of homeobox genes. The fusion partners of *MLL* are either transcriptional activators (such as *ENL*), making activation of *MLL* target genes independent of *MLL* homodimerization, or proteins with oligomerization domains (such as *SEPT6*), causing unregulated dimerization and constitutive activation of *MLL*. However, expression of *MLL* fusion proteins in animals does not induce AML with the rapidity that is seen in the human disease. So Ono and colleagues investigated the idea that secondary genetic events might be involved.

The authors focused on *MLL*–*SEPT6*. *SEPT6* is one of the septins, cytoplasmic proteins that are required

for cytokinesis. Expression of the retrovirally expressed *MLL*–*SEPT6* in mouse haematopoietic progenitor cells *in vitro* resulted in immortalization through a block in cell differentiation and increased self-renewal. Transplantation of these cells into a mouse model induced, after a long latency, a lethal myeloproliferative disease, but not AML.

MLL–*SEPT6*-expressing cells isolated from these animals require interleukin-3 (*IL-3*) for survival. Given that *IL-3* independence is associated with leukaemic progression, the authors reasoned that *IL-3* independence might be a cooperating lesion. They co-infected the *MLL*–*SEPT6*-expressing cells with a retrovirus expressing the mutant form of FMS-like receptor tyrosine kinase 3 (*FLT3*), which has also been implicated in AML and might substitute for the tyrosine-kinase survival signal normally supplied by *IL-3*. Co-expression of this gene led to *IL-3*-independent growth *in vitro* and the induction of AML with short latency *in vivo*.

Given that mutant *FLT3* also cooperated with *MLL*–*ENL*, the authors surmise that *MLL* fusion proteins impair differentiation and enhance the self-renewal capacity of haematopoietic progenitors, and activation of *FLT3* confers a proliferative and/or survival capacity.

Nicola McCarthy

 **References and links**

ORIGINAL RESEARCH PAPER Ono, R. *et al.* Dimerization of *MLL* fusion proteins and *FLT3* activation synergize to induce multi-lineage leukemogenesis. *J. Clin. Invest.* 10 Mar 2005 (doi:10.1172/JCI200522725)



IN BRIEF

STEM CELLS

Skin carcinoma arising from donor cells in a kidney transplant recipient.

Aractingi, S. *et al. Cancer Res.* **65**, 1755–1760 (2005)

Cutaneous tumours arising in female patients who had received a male kidney were examined by quantitative polymerase chain reaction and fluorescence *in situ* hybridization. Male cells were detected in many cases. A particular carcinoma showed a pattern of male cells that seemed to have arisen from the clonal expansion of a single donor cell. So, stem cells from a grafted organ might migrate to the skin, differentiate into, or fuse with, keratinocytes and then, rarely, undergo neoplastic transformation.

CIRCADIAN RHYTHMS

Circadian sensitivity to the chemotherapeutic agent cyclophosphamide depends on the functional status of the *CLOCK*/*BMAL1* transactivation complex.

Gorbacheva, V. Y. *et al. Proc. Natl Acad. Sci. USA* **102**, 3407–3412 (2005)

How does the time of day regulate the body's sensitivity to chemotherapeutic agents? Gorbacheva *et al.* showed that mice lacking functional copies of two main circadian-rhythm-regulating genes, *CLOCK* and *BMAL1*, are highly sensitive to treatment with cyclophosphamide at any time of day, unlike controls. This is due to a decrease in the recovery and survival rate of B cells after treatment.

LEUKAEMIA

A sumoylation site in *PML/RARA* is essential for leukemic transformation.

Zhu, J. *et al. Cancer Cell* **7**, 143–153 (2005)

The *PML*–*RARA* oncogene causes acute promyelocytic leukaemia (APL) by repressing myeloid differentiation. Zhu *et al.* find that sumoylation of *PML* is required for transformation *ex vivo*. Expression in mice of a *PML*–*RARα* mutant that cannot be sumoylated leads to myeloid hyperplasia, but not to leukaemia. The authors show that sumoylation allows recruitment of the potent transcriptional repressor *DAXX*, indicating a mechanism by which *PML*–*RAR* might initiate APL.

TUMORIGENESIS

The *RET/PTC*–*RAS*–*BRAF* linear signaling cascade mediates the motile and mitogenic phenotype of thyroid cancer cells.

Melillo, R. M. *et al. J. Clin. Invest.* 10 Mar 2005 (doi:10.1172/JCI200522758)

Activating mutations of the oncogenes *BRAF* or *RAS*, or fusions of the *RET* tyrosine receptor kinase are all found in papillary thyroid carcinomas, but never together. Melillo *et al.* reveal that this is because they function in a linear oncogenic signalling pathway, where *RET* fusions stimulate *RAS*-dependent activation of *BRAF*. This cascade initiates a transcriptional programme in thyroid cells that eventually stimulates proliferation and invasion.