CIRCADIAN RHYTHMS

Clocking on

Women who work at night have an increased risk of developing breast or colon cancer, which could be evidence that disruption of our circadian rhythm upsets more than just our sleep patterns. The genes that regulate the circadian clock in mammals have recently been identified, and Sigal Gery and colleagues now show that one of these genes, period 1 (*PER1*), might function as a tumour-suppressor gene.

Given the established link between circadian rhythms and the DNA-damage response, the authors investigated the response to ionizing radiation (IR) of colorectal cancer cell lines that overexpress *PER1*. The authors showed that overexpression of *PER1* induced high expression levels of p53 that were only slightly increased after treatment with IR. Despite p53 expression, these cells did not have increased expression of p21, the cyclin-dependent-kinase inhibitor that mediates p53-induced cell-cycle arrest after DNA damage. This is because, after IR, *PER1* overexpression results in the increased expression of the oncoprotein MYC, which is known to inhibit p21 expression and induce apoptosis. Therefore, cells that expressed *PER1* were less able to arrest in G2/M and had higher rates of apoptosis than control cells.

Given that *PER1* seems to be important for the cellular response to IR, does it interact with other proteins that regulate this pathway? Co-immunoprecipitation experiments showed that *PER1* can interact with the DNA-damage-response kinase, ataxia telangiectasia mutated (*ATM*), and the checkpoint protein *CHK2*. The authors propose that *PER1* might facilitate the activation of *ATM* or might function as an adaptor protein that recruits other proteins that are required for the *ATM*-mediated response to DNA damage.

Overexpression of *PER1* in various cancer cell lines inhibited clonogenic growth as well as anchorage-independent growth, which further indicates that *PER1* might function as a tumour suppressor. So the authors investigated whether *PER1* expression is altered in human cancer. Reverse transcriptase-PCR analyses showed that 70% of human non-small-cell lung cancer samples and 46% of human breast cancer samples had reduced expression levels of *PER1* compared with controls.

These results further increase our preliminary understanding of how the circadian clock regulates pathways that are crucial for the suppression of malignant growth.

Nicola McCarthy

ORIGINAL RESEARCH PAPER Gery, S. *et al.* The circadian gene *Per1* plays an important role in cell growth and DNA damage control in human cancer cells. *Mol. Cell* 22 375–382 (2006)

TUMORIGENESIS

Nuclear bodies

Tumorigenesis that is driven by loss of the tumour suppressor PTEN is known to activate the proto-oncogene *AKT* in many human cancers. Activation of cytoplasmic *AKT* has been thought to be the most relevant. However, Pier Paolo Pandolfi and colleagues now show that additional deficiency of the tumour suppressor *PML* — which forms dynamic intranuclear structures (nuclear bodies) that are involved in apoptosis, proliferation and senescence — leads to an increase in the nuclear localization of *AKT*, which accelerates tumorigenesis.

First the authors studied the effect of *Pml* loss on *AKT* activation in *Pten^{+/-}* mice. *AKT* activation increased by 50% in normal colon and prostate cells, particularly in the nuclear compartment, and led to the development of colon and prostate cancers. To investigate whether the change in

phospho-*AKT* (pAKT) localization was an inherent feature of *Pml* loss, Pandolfi and colleagues looked at the *AKT* status of *Pml^{+/+}* and *Pml^{-/-}* mouse embryonic fibroblasts (MEFs). After the addition of serum to the cell culture, *AKT* activation — especially in the nucleus — was sustained in *Pml^{-/-}* MEFs. If *Pml* was transduced into *Pml^{-/-}* cells, *AKT* activation decreased. These findings were confirmed by confocal laser scanning microscopy.

So, does *Pml* loss increase the function of nuclear pAKT? The forkhead transcription factor *FOXO3a* is a nuclear target of pAKT, and phosphorylation of *FOXO3a* leads to its inactivation and export from the nucleus. Decreased nuclear *FOXO3a* was seen after serum stimulation in *Pml^{-/-}* MEFs compared with *Pml^{+/+}* MEFs. In addition, the *FOXO3a*

POPULATION GENETICS

Associated risk

Family history and ethnicity are known risk factors for prostate cancer, but little is known about the genetic factors behind them. Amundadottir *et al.* have now shown that a common allele is associated with prostate cancer, and that it is more common in men of African descent, who are at greater risk of prostate cancer, than in men of European descent.

Variation at microsatellite loci might have functional consequences, and can also be used to search for regions of the genome that are associated with disease risk. The authors genotyped European men with prostate cancer and found that one particular microsatellite allele, DG8S737–8, on chromosome 8q24 is associated with disease risk.

This could be because the allele itself confers risk, or because it is close to the locus that does and most individuals who carry the microsatellite allele also carry the risk allele. This would be the case if there had been little recombination between the two loci since the risk allele arose by mutation.

Significantly, the authors found that the DG8S737–8, which also increases the severity of prostate cancer, was more common in men of African descent and was also closely associated with prostate cancer risk in this population. First, this could explain why prostate cancer is more common and more severe in men of African descent; and second, it shows that the area of the chromosome in which the risk factor lies is very close to the microsatellite, as all but the closest associations would

target genes — tumour suppressor *Cdkn1b* (which encodes p27), the pro-apoptotic factor *Bim* and DNA repair protein *Gadd45* — were downregulated before the development of prostate tumours.

What is the mechanism by which *Pml*-loss regulates AKT? It could be either stimulation of AKT activation or prevention of pAKT inactivation. The only known AKT Thr308 phosphatase — PP2A — was shown to specifically bind PML and was enriched in PML nuclear bodies, as was pAKT. These results indicate that *Pml*-deficiency must antagonize PP2A function by preventing the recruitment of both pAKT and PP2A into PML bodies.

These data show that PML controls the nuclear tumour-suppressor network for the inactivation of nuclear pAKT, and highlights the importance of AKT localization in human cancer cells.

Ezzie Hutchinson

ORIGINAL RESEARCH PAPER Trotman, L.C. *et al.* Identification of a tumour suppressor network opposing nuclear Akt function. *Nature* 7 May 2006 (doi: 10.1038/nature04809)

be expected to have broken down by recombination since African and European populations diverged.

It is therefore likely that the microsatellite variant itself confers cancer risk, but the authors also considered two other candidate risk factors. First, they found an unknown transcript that is expressed in differently spliced forms in normal and prostate cell lines. Second, although *MYC* is too far away on the chromosome to be associated with the microsatellite, it is near enough for remote expression elements or genomic instability near the microsatellite to affect its expression. Although the actual risk factor has not been firmly identified, these results are a significant step towards doing so.

Patrick Goymer

ORIGINAL RESEARCH PAPER Amundadottir, L.T. *et al.* A common variant associated with prostate cancer in European and African populations. *Nature Genet.* 7 May 2006 (doi:10.1038/ng1808)

HEPATOCELLULAR CARCINOMA

Putting p53 in context

Hepatocellular carcinoma is one of the most prevalent cancers worldwide and is lethal in about 75% of cases. To understand the aetiology of this disease more clearly, Ron DePinho and colleagues have examined chemically induced liver damage in mouse models and found that genomic instability that is fuelled by telomere dysfunction and reduced p53 function can cooperate to induce hepatocellular carcinoma progression.

Although hepatocellular carcinomas have various aetiologies, one common factor could be cycles of hepatocyte necrosis and regeneration that might promote genetic instability and create a favourable microenvironment for carcinoma progression. Mice that have been treated with the hepatocarcinogen carbon tetrachloride (CCl_4), which mimics key aspects of human chronic liver disease, readily develop hepatocellular carcinomas. The authors observed that levels of expression of telomerase reverse transcriptase (TERT) are increased in most CCl_4 -induced hepatocellular carcinomas, and that TERT levels increased with disease progression. Given that telomere dysfunction and p53 inactivation are frequently seen in human hepatocellular carcinomas, DePinho and colleagues used *Tert*-null mice and *Trp53*-heterozygous mice to investigate further.

As expected, *Tert*-null mice had shorter telomeres with each generational intercross, with telomere dysfunction observed by the third or fourth generation. *Tert*-heterozygous mice were crossed with *Trp53*-heterozygous mice to obtain *Trp53*^{+/+} and *Trp53*^{+/-} mice that were also *Tert*^{+/-} (intact telomeres). Intercrossing of these genotypes and further successive generational

mating resulted in mice with dysfunctional telomeres.

In mice with intact telomeres, loss of one *Trp53* allele did not affect the incidence of CCl_4 -induced early neoplastic lesions or hepatocellular carcinoma. However, telomere dysfunction significantly suppressed the tumorigenesis that was observed in *Trp53*^{+/-} mice, and this suppression was reversed by loss of a single *Trp53* allele. The authors also examined the histopathological grade of the hepatocellular carcinomas and found that mice with telomere dysfunction and intact p53 had complete suppression of aggressive disease, whereas the *Trp53* heterozygotes developed aggressive hepatocellular carcinomas with no deletion or mutation of the wild-type *Trp53* allele. There was also a trend toward accelerated progression in *Trp53* heterozygotes with dysfunctional rather than intact telomeres.

DePinho and colleagues conclude that the effect of p53 loss on hepatocarcinogenesis that is induced by CCl_4 is dependent on cellular context — intact or dysfunctional telomeres — and they hypothesize that loss of p53 in hepatocellular carcinoma that is associated with chronic liver disease might function predominantly to enable hepatocyte survival in the presence of telomere dysfunction. In addition, viral transduction of *Tert* in this mouse model might permit the creation of a more accurate model of human hepatocellular carcinomas, which show telomerase reactivation.

Sarah Seton-Rogers

ORIGINAL RESEARCH PAPER Farazi, P.A. *et al.* Cooperative interactions of p53 mutation, telomere dysfunction, and chronic liver damage in hepatocellular carcinoma progression. *Cancer Res.* 66, 4766–4773 (2006)