

# BCR–ABL and *CDKN2A*: a dropped connection

Richard T. Williams\* and Charles J. Sherr†

In their recent Review of chronic myeloid leukaemia (CML) stem cell biology (Savona, M. & Talpaz, M. Getting to the stem of chronic myeloid leukemia. *Nature Rev. Cancer* 8, 341–350)<sup>1</sup>, Savona and Talpaz focus their attention exclusively on deregulated self-renewal mechanisms in Philadelphia chromosome (Ph)<sup>+</sup> hematopoietic stem cells (HSCs) in chronic-phase CML and on the role of  $\beta$ -catenin abnormalities in granulocyte–macrophage progenitors in myeloid blast crisis. However, a review of genomic abnormalities in lymphoid blast crisis provides insights into alternative mechanisms of progenitor self-renewal in BCR–ABL-driven diseases. Deletion of the *CDKN2A* locus (encoding the *INK4A* and *ARF* tumour suppressor genes) frequently occurs in the progression of CML from chronic-phase to lymphoid, but not myeloid, blast crisis<sup>2</sup>, and a recent genome-wide analysis<sup>3</sup> (published concurrently with this CML review) has reinforced the conclusions of earlier studies. *INK4A* and *ARF* share common second and third exons, are translated in alternate reading frames, and exhibit no sequence similarity<sup>4</sup>. Whereas p16<sup>INK4A</sup> is a *bona fide* cyclin-dependent kinase inhibitor, p14<sup>ARF</sup> (p19<sup>Arf</sup> in mice) is not, but instead regulates p53 tumour suppressor function through its interaction with MDM2 (REF. 5). BCR–ABL induces p19<sup>Arf</sup> *in vivo* to oppose tumour development, but

the combination of BCR–ABL expression and *Arf* inactivation is sufficient to generate leukaemia-initiating cells in a murine model of Ph<sup>+</sup> acute lymphocytic leukaemia<sup>6</sup>. Therefore, in BCR–ABL-driven lymphoid leukaemias — specifically, CML lymphoid blast crisis and Ph<sup>+</sup> acute lymphocytic leukaemia — deletion of *INK4A* and *ARF* in committed Ph<sup>+</sup> lymphoid blasts is expected to simultaneously disrupt both the RB and p53 tumour suppressor networks, thereby facilitating leukaemia-initiating cell self-renewal and enhancing targeted therapeutic resistance *in vivo*.

\*Departments of Oncology and of Genetics & Tumor Cell Biology and †Howard Hughes Medical Institute, St. Jude Children's Research Hospital, 262 Danny Thomas Place, Memphis, Tennessee 38105, USA.

Correspondence to R.T.W.

e-mail: richard.williams@stjude.org

1. Savona, M. & Talpaz, M. Getting to the stem of chronic myeloid leukemia. *Nature Rev. Cancer* 8, 341–350 (2008).
2. Calabretta, B. & Perrotti, D. The biology of CML blast crisis. *Blood* 103, 4010–4022 (2004).
3. Mullighan, C. G. *et al.* BCR–ABL1 lymphoblastic leukaemia is characterized by the deletion of Ikaros. *Nature*, 453, 110–114 (2008).
4. Quelle, D. E. *et al.* Alternative reading frames of the *INK4a* tumor suppressor gene encode two unrelated proteins capable of inducing cell cycle arrest. *Cell* 83, 993–1000 (1995).
5. Sherr, C. J. & McCormick, F. The RB and p53 pathways in cancer. *Cancer Cell* 2, 103–112 (2002).
6. Williams, R. T., den Besten, W. & Sherr, C. J. Cytokine-dependent imatinib resistance in mouse BCR–ABL<sup>+</sup>, *Arf*-null lymphoblastic leukemia. *Genes Dev.* 21, 2283–2287 (2007).