CELL DEATH

1+1≠2

The order and combination in which cancer drugs are given can determine whether or not tumour cells respond. Michael Yaffe and colleagues now show that pretreatment with one targeted drug for a specific period of time can increase the sensitivity of triple-negative breast cancer (TNBC) cells to standard chemotherapy.

By adopting a systems biology approach to examine the changes that are imposed by drugs on signalling networks in breast cancer cell lines, these authors found that a 4-hour pretreatment of a TNBC cell line (BT-20) with erlotinib (an inhibitor of the epidermal growth factor receptor (EGFR)), followed by doxorubicin significantly increased cell death compared with either drug alone or with both drugs given at the same time. This drug combination and timing was only synergistic in TNBC cells and not in ERBB2-expressing breast cancer cells for example, and extended to other inhibitors of EGFR and other DNA-damaging drugs.

How does pretreatment with an EGFR inhibitor increase sensitivity to a DNA-damaging agent? Yaffe and colleagues investigated changes in gene expression and found that a substantial number of genes were differentially expressed in BT-20 cells in response to erlotinib compared with breast cancer cells that are not sensitized to doxorubicin by erlotinib. GeneGO pathway annotation showed that these differentially expressed genes were most often involved in apoptosis, inflammation and differentiation pathways. Interestingly, gene set enrichment analyses showed that the expression of genes known to be associated with TNBC status was reduced by chronic erlotinib treatment and that the cells increased the expression of genes associated with a luminal A subtype of breast cancer, a less aggressive disease with a good overall prognosis.

To further understand the effect of EGFR inhibition, the authors measured time-dependent signalling events and constructed a data-driven mathematical model (based on *in vitro* changes in protein expression and activity) that related signalling 'inputs' to phenotypic 'outputs'. This model could accurately predict phenotypic outcomes, such as apoptosis or autophagy, following treatment, and it indicated that caspase 8 was likely to be a key determinant of sensitivity induced by erlotinib. Indeed, knockdown of caspase 8 in BT-20 cells prevented the synergism between erlotinib and doxorubicin. *In vitro* analyses of an additional ten TNBC cell lines indicated that nine of the ten responded to erlotinib followed by doxorubicin, but that only four lines had a synergistic response. In all four lines this correlated with caspase 8 cleavage and high levels of basal EGFR phosphorylation, but not with expression levels of EGFR. Treatment of BT-20 xenografts with doxorubicin alone resulted in tumour regression followed by regrowth within 14 days. However, pretreatment with erlotinib 8 hours before doxorubicin treatment reduced tumour volume and prevented regrowth within 14 days.

The authors further extended this pretreatment paradigm to show that ERBB2-overexpressing breast cancer cell lines can be sensitized to doxorubicin by pretreatment with lapatinib (an ERBB2 and EGFR inhibitor). Moreover, lung cancer cell lines with active EGFR also respond synergistically to erlotinib pretreatment and doxorubicin, and this was inhibited by caspase 8 knockdown.

These results indicate that an assessment of changes in signalling and gene expression networks in response to specific drug combinations could be used to determine useful sequences of standard chemotherapies and targeted agents.

Nicola McCarthy

ORIGINAL RESEARCH PAPER Lee, M. J. et al. Sequential application of anticancer drugs enhances cell death by rewiring apoptotic signalling networks. *Cell* **149**, 780–794 (2012)

“
knockdown
of caspase 8
in BT-20 cells
prevented the
synergism
between
erlotinib and
doxorubicin